

## Fréchet derivative.

Let  $D$  be a doubly connected domain in  $\mathbb{R}^2$  with smooth boundary  $\Gamma$  of class  $C^2$ . We assume that  $\Gamma$  consists of two disjoint curves  $\Gamma_1$  and  $\Gamma_2$ , meaning  $\Gamma = \Gamma_1 \cup \Gamma_2$ , with  $\Gamma_1 \cap \Gamma_2 = \emptyset$ , such that  $\Gamma_1$  is contained in the interior of  $\Gamma_2$ .

We consider the integral operator

$$(A\phi)(x) = \int_{\Gamma_1} \phi(y) \Phi(x, y) ds(y), \quad x \in \Gamma_2. \quad (0.1)$$

The exterior curve has the parametric representation

$$\Gamma_2 = \{x_2(s) = (x_{12}(s), x_{22}(s)), s \in [0, 2\pi]\}.$$

For simplicity, we consider starlike interior curve

$$\Gamma_1 = \{x_1(s) = r(s)(\cos s, \sin s) : s \in [0, 2\pi]\}, \quad (0.2)$$

where  $r : \mathbb{R} \rightarrow (0, \infty)$  is a  $2\pi$  periodic function representing the radial distance from the origin.

The parametrized operator has the form

$$(A\phi)(s) = \int_0^{2\pi} \phi(\sigma) \Phi(x_2(s), x_1(\sigma)) d\sigma.$$

Let  $q$  is the radial function of the perturbed interior boundary.

The Fréchet derivative  $\mathcal{A}$  of the integral operator  $A$  with respect to  $x_1$  as a linear operator on  $q$ . The integral operator  $\mathcal{A}$  can be obtained by formal differentiation of the kernel  $\Phi$  with respect to  $x_1$

$$\mathcal{A}[\phi, r; q](s) = \int_0^{2\pi} q(\sigma) \phi(\sigma) \Psi(s, \sigma) d\sigma$$

with kernel

$$\Psi(s, \sigma) = \text{grad}_{x_1} \Phi(x_2(s), x_1(\sigma)) \cdot (\cos \sigma, \sin \sigma).$$

For example, let

$$\Phi(x, y) = \ln \frac{1}{|x - y|}, \quad x \neq y.$$

Then

$$\Psi(s, \sigma) = \frac{(x_2(s) - x_1(\sigma)) \cdot (\cos \sigma, \sin \sigma)}{|x_2(s) - x_1(\sigma)|^2}.$$