

# Solutes and water cycling in catchments

Prof. Kate Heal  
School of GeoSciences

Soil, Water & Atmospheric Processes

# Carbon and Nitrogen Cycling in the Forest Water Cycle

Prof. Kate Heal, Nick Forrest, Daniele  
Ferraretto

School of GeoSciences, The University of Edinburgh

Prof. Xiangqing Ma, Dr Xiaolong Hou  
Forestry College

Fujian Agriculture and Forestry University, China



# Acknowledgements

- Funders: China Scholarship Council, NERC, Elizabeth Sinclair Irvine Bequest, Forest Research
- Dr Robert Clement, Prof. Maurizio Mencuccini, Dr Lisa Wingate, Prof. Dave Reay, Prof. John Moncrieff, Prof. Paul Jarvis (University of Edinburgh)
- Dr Sirwan Yamulki, Dr Nadeem Shah (Forest Research)
- Technical support at the University of Edinburgh: Robert Howard, Michael Spencer, Emanuel Blei, Philip Stack, Andrew Gray, John Morman
- Dr Andy Stott (NERC Life Sciences Mass Spectrometry Facility)
- Tillhill Economic Forestry Ltd. for site access



**UPM** TILHILL



# Background

- Climate change; increasing atmospheric CO<sub>2</sub> concentrations => carbon (C) sequestration in biomass
- **Forests apparent C sinks BUT sink strength affected by:**
  - Release from soil
  - Loss of C through water flux
  - Nutrient availability
  - Forest age and management



# Research questions

- How do C losses in water compare with C uptake by forest?
- What is the relationship between C sequestration and forest nitrogen cycling?
- How to scale from point measurements to forest-wide fluxes?

# Study site: Griffin Forest (56.6°N, 3.8°W)

- Cool temperate climate
- Altitude: 275-532 m above sea level
- Catchment area 4.47 km<sup>2</sup>
- Average slope 6°
- Geology: Dalradian schist overlain by glacial till
- Soils: 0.7 m depth humic gley
- Forest planted 1980-1981
- Sitka spruce dominant
- C sequestration measured since 1997






# Field measurements of hydrological flux

Logging raingauges



Storage raingauges



[www.met-office.gov.uk](http://www.met-office.gov.uk)

Stream outflow




Throughfall  
& stemflow  
samplers



[www.bangor.ceh.ac.uk](http://www.bangor.ceh.ac.uk)

## Ratings curve for Cultullich Burn, Griffin Forest 2000

$$\log_{10} (Q) \text{ (l/s)} = - 6.853 + 3.655 \log_{10} (\text{reading})$$



—	Regression
- - -	95% CI
S	0.107552
R-Sq	97.8%
R-Sq(adj)	97.5%



Water depth readings every 15 minutes recorded electronically by data logger



# Other measurements

- Chemical analyses of water samples
  - pH
  - $\text{NO}_3\text{-N}$ ,  $\text{NH}_4\text{-N}$
  - Total carbon (TC), dissolved organic carbon (DOC), total inorganic carbon ( $\text{TIC} = \text{TC} - \text{DOC}$ )
- Forest characterisation
  - tree DBH (diameter at breast height) surveyed




Image credit: Michael Spencer

# Measurement of C uptake by the forest



- Measured as flux of  $\text{CO}_2$  up and down from forest
- **Eddy covariance**  
(micrometeorological method)
- Movement of air parcels measured in 3 dimensions using a **sonic anemometer**
- Air analysis  $\Rightarrow$  flux of  $\text{CO}_2$  calculated
- Other micrometeorological measurements to check energy balance and calculate evapotranspiration


# Rainfall amount results



No spatial variation in rainfall totals in catchment

No relationship between rainfall and altitude

# Rainfall and streamflow 2000




# Annual evapotranspiration estimates 2000

- Catchment water balance method

$E = \text{precipitation} - \text{streamflow}$

- Eddy covariance

$$E = \frac{\text{Measured latent energy}}{\text{Latent heat of energy of evaporation of water}}$$



Close agreement between evapotranspiration estimates => confidence in hydrological flux measurements


(Nick Forrest)




# Problems of scaling up throughfall/stemflow fluxes

Need to take account of factors such as:

- Depth/volume vs. tree properties relationships



# Relationship between N in rainfall and altitude



Increased N deposition with altitude  
well-documented

# Streamwater carbon and nitrogen flux calculations

- Mass flux = concentration x flow
- Frequent flow measurements **BUT infrequent concentration measurements** => different flux calculation methods

## 1) Mean value

Flux = flow volume (from mean flow) x mean concentration for whole study period


## 2) Individual time periods

$$\text{Flux} = \sum_{t=1}^{t=n} (\text{flow volume (from mean flow)} \times \text{mean concentration})$$

## 3) Continuous


Where relationship between concentration and flow, concentration estimated for every streamflow measurement

# Flow-concentration relationships for TIC in stream water



At low flows TIC inputs from mineral weathering in deeper soil horizons less diluted


# Flow-concentration relationships for DOC in stream water



DOC flushed from organic near-surface  
soil horizons at high flows



# Annual hydrological N and C fluxes, 2000



Net nitrogen gain by catchment ( $0.027 \text{ t ha}^{-1} \text{ a}^{-1}$ )

Net carbon loss from catchment ( $0.026 \text{ t ha}^{-1} \text{ a}^{-1}$ )

# Comparison of C losses in water to C sequestration

Parameter	Study site		
	Griffin Forest (Scotland)	Hokkaido (N Japan)	Moor House (N England)
<i>Reference</i>	<i>This research</i>	<i>Shibata et al. (2005)</i>	<i>Worrall et al. (2003)</i>
Vegetation	Sitka spruce plantation	Temperate deciduous forest	Heather (dwarf shrub)
C loss in water (t ha <sup>-1</sup> a <sup>-1</sup> )	-0.026	-0.04 (incl. POC)	-0.348 (-0.149 without POC)
C sequestered (t ha <sup>-1</sup> a <sup>-1</sup> )	6.1	2.6	0.55
% sequestered C lost in water	0.43	1.5	63 (27 without POC)

Carbon losses in river water small % of C sequestered in forests,  
**BUT** may be more significant for shrub vegetation cover

# Ongoing research focus on role of the forest canopy in nitrogen cycling

Rainfall and  
cloud water  
collection x 2




Stemflow  
collection  
x 22

Throughfall  
and litter  
collection x 18



Stream flow gauging x 4

# Preliminary results for forest canopy N processing



Confirms c.70% N deposition intercepted by canopy  
Stemflow insignificant for N fluxes

# Interaction between atmospheric N and C sequestration by forests

- Possible explanations for atmospheric N uptake by forest canopy
  1. Foliar uptake by needles
  2. Cation exchange for  $\text{NH}_4\text{-N}$
  3. Uptake by microfauna & algae in canopy (sometimes associated with aphid infestation)
- C sequestration per unit added N ( $\Delta\text{C}:\Delta\text{N}$ )
  - 210 (Magnani et al., 2007) – considering atmospheric N deposition (onto forest canopy)
  - 46 (De Vries et al., 2006) – considering N application onto forest soil
  - c.170 in 2000 at Griffin Forest
- Although atmospheric N deposition may enhance C sequestration it may => acidification of soils and disruption of nutrient cycling



# Recommended papers with further information on this topic

Clement, R. J., Jarvis, P. G. and Moncrieff, J. B. (2012). Carbon dioxide exchange of a Sitka spruce plantation in Scotland over five years. *Agricultural and Forest Meteorology*, 153, 106-123.

Dail, D. B., Hollinger, D. Y., Davidson, E. A., Fernandez, I., Sievering, H. C., Scott, N. A. and Gaige, E. (2009). Distribution of nitrogen-15 tracers applied to the canopy of a mature spruce-hemlock stand, Howland, Maine, USA. *Oecologia*, 160, 589-599.

Magnani, F. et al. (2007). The human footprint in the carbon cycle of temperate and boreal forests. *Nature*, 447, 848-850.