

Notes on the solution to 1D Schrödinger equation for piecewise potentials

bjd

Facultat de Física, Universitat de Barcelona, Diagonal 645, 08028 Barcelona, Spain.

Brief rendition of the way to solve the Schrödinger equation for piecewise potentials

We consider the time independent 1D Schrödinger equation, (in natural units)

$$-\frac{1}{2}\partial_x^2\psi(x) + V(x)\psi(x) = E\psi(x). \quad (1)$$

Where the potential is taken to be piecewise constant, that is,

$$V(x) = V_k \quad x_k < x < x_{k+1}. \quad (2)$$

where $\{x_k\}$ is a partition of the interval $[0, L]$, and $k = 0, \dots, N$. For simplicity we take the utmost left and right potentials to be infinite, that is, the wave function is zero at $x = 0$ and $x = L$. For each interval, $[x_k, x_{k+1}]$ the general solution of the Schrödinger equation reads,

$$\psi_k(x) = A_k e^{i\kappa_k x} + B_k e^{-i\kappa_k x} \quad (3)$$

where $\kappa_k = \sqrt{2(E - V_k)}$.

First we consider the first and last intervals,

$$\begin{aligned} \psi_0(x) &= A_0 e^{i\kappa_0 x} + B_0 e^{-i\kappa_0 x} \\ \psi_N(x) &= A_N e^{i\kappa_N x} + B_N e^{-i\kappa_N x} \end{aligned} \quad (4)$$

the boundary conditions imply,

$$\begin{aligned} \psi_0(x) &= A_0 (e^{i\kappa_0 x} - e^{-i\kappa_0 x}) = A_0 2i \sin(\kappa_0 x) \\ \psi_N(x) &= A_N e^{i\kappa_N L} (e^{i\kappa_N (x-L)} - e^{-i\kappa_N (x-L)}) = A_N 2i \sin(\kappa_N (x-L)) \end{aligned} \quad (5)$$

If we consider two inner intervals we have to impose continuity of the wave function and of its first derivative, that reads,

$$\begin{aligned} \psi_k(x_{k+1}) &= \psi_{k+1}(x_{k+1}) \\ \psi'_k(x_{k+1}) &= \psi'_{k+1}(x_{k+1}) \end{aligned} \quad (6)$$

which read,

$$\begin{aligned} A_k e^{i\kappa_k x_{k+1}} + B_k e^{-i\kappa_k x_{k+1}} &= A_{k+1} e^{i\kappa_{k+1} x_{k+1}} + B_{k+1} e^{-i\kappa_{k+1} x_{k+1}} \\ A_k i\kappa_k e^{i\kappa_k x_{k+1}} - B_k i\kappa_k e^{-i\kappa_k x_{k+1}} &= A_{k+1} i\kappa_{k+1} e^{i\kappa_{k+1} x_{k+1}} - B_{k+1} i\kappa_{k+1} e^{-i\kappa_{k+1} x_{k+1}}. \end{aligned} \quad (7)$$

These can be written in matrix form as,

$$\begin{pmatrix} e^{i\kappa_k x_{k+1}} & e^{-i\kappa_k x_{k+1}} \\ i\kappa_k e^{i\kappa_k x_{k+1}} & -i\kappa_k e^{-i\kappa_k x_{k+1}} \end{pmatrix} \begin{pmatrix} A_k \\ B_k \end{pmatrix} = \begin{pmatrix} e^{i\kappa_{k+1} x_{k+1}} & e^{-i\kappa_{k+1} x_{k+1}} \\ i\kappa_{k+1} e^{i\kappa_{k+1} x_{k+1}} & -i\kappa_{k+1} e^{-i\kappa_{k+1} x_{k+1}} \end{pmatrix} \begin{pmatrix} A_{k+1} \\ B_{k+1} \end{pmatrix} \quad (8)$$

which can be written as,

$$\mathcal{M}(\kappa_k, x_{k+1}) \phi_k = \mathcal{M}(\kappa_{k+1}, x_{k+1}) \phi_{k+1} \quad (9)$$

where,

$$\mathcal{M}(\kappa, x) = \begin{pmatrix} e^{i\kappa x} & e^{-i\kappa x} \\ i\kappa e^{i\kappa x} & -i\kappa e^{-i\kappa x} \end{pmatrix} \quad \phi_k = \begin{pmatrix} A_k \\ B_k \end{pmatrix} \quad (10)$$

This allows us to solve the wave function in the $k+1$ interval from the values in the k interval,

$$\phi_{k+1} = \mathcal{M}^{-1}(\kappa_{k+1}, x_{k+1}) \mathcal{M}(\kappa_k, x_{k+1}) \phi_k \quad (11)$$

where the inverse matrix reads,

$$\mathcal{M}^{-1}(\kappa, x) = \begin{pmatrix} \frac{1}{2} e^{-i\kappa x} & \frac{-i}{2\kappa} e^{-i\kappa x} \\ \frac{i}{2} e^{i\kappa x} & \frac{1}{2\kappa} e^{i\kappa x} \end{pmatrix} \quad (12)$$

A. how to find the eigenenergies?

To find the energy quantization we basically write the wave function in the last interval as a function of the wave function in the first interval,

$$\hat{\phi}_N = \mathcal{M}^{-1}(\kappa_N, x_N) \mathcal{M}(\kappa_{N-1}, x_N) \dots \mathcal{M}^{-1}(\kappa_{j+1}, x_{j+1}) \mathcal{M}(\kappa_j, x_{j+1}) \dots \mathcal{M}^{-1}(\kappa_1, x_1) \mathcal{M}(\kappa_0, x_1) \phi_0 \quad (13)$$

with $\phi_0 = \{A_0, -A_0\}$. This should indeed be equal to ϕ_N , that we wrote in Eq. (??).

Now for instance we do,

$$\hat{\phi}_N(1)/\hat{\phi}_N(2) - \phi_N(1)/\phi_N(2) = 0 \quad (14)$$

and look for values of E that solve the above equation. Notice in the above equation the values of A_0 and A_N .
