Observatoire astronomique de Strasbourg | ObAS

Database Management Project

Analysis of Galactic X-ray Transients in eROSITA Using MySQL



Written by:
Annus Haider

Supervised by: Sebastian Derriere

Academic year 2025-2026

Contents

Introduction **2**

1 Data Extraction and Loading **2**

2 SQL Queries and Analysis **3**

Initial Database Checks . . . 3
Data Cleaning: Handling empty values . . . 5
Classes with average temperature > 5000 K . . . 5
Distribution of sources across different classes . . . 6
Identification of the five brightest X-ray sources in the catalogue . . . 6
Identification of nearby X-ray bright stars with known spectral types . . . 7
Identification of sources with extremely soft colors . . . 7
Search for candidate compact objects . . . 8
Identification of rare extreme luminosity outliers . . . 8
How does the average X-ray luminosity vary among different source classes? . . . 9
Are high-luminosity transients preferentially distant? . . . 10
What is the average hardness ratio for different source classes? . . . 10

Conclusion **11**

List of Figures

Introduction

The catalogue by Maan et al. [1] compiles **Galactic X-ray transients detected by the eROSITA telescope**. It provides detailed observational information on transient sources, including their positions, fluxes, spectral properties, and detection times. The catalogue covers a wide portion of the Galactic plane and includes multiple epochs of observations, allowing the study of variability in X-ray emission over time. In total, the catalogue report the X-ray and optical properties of 738 transients, classified into the following categories:

1. **YSO:** Young stellar object (105 transients)
2. **SS/IB:** Either chromospherically active single star or interacting binary (418 transients)
3. **IB:** Known interacting binary not having any compact object in the system (110 transients)
4. **IB/ACO:** Interacting binary or accreting compact object (50 transients)
5. **CV:** Cataclysmic variable (38 transients)
6. **XRB:** X-ray binary (3 transients)
7. **LPV:** Long period variable (12 transients)
8. **PSR:** Pulsar (1 transient)
9. **WR:** Wolf-Rayet (1 transient)

The goal of this project is to **explore and analyze the eROSITA Galactic X-ray transient catalogue using MySQL**. By performing a variety of queries, we aim to:

- Extract and filter relevant data from the catalogue.
- Compute statistics such as flux distributions, source counts, and temporal variations.
- Demonstrate the use of advanced MySQL commands, including joins, subqueries, aggregation, and data updates.
- Gain insights into the properties and distribution of Galactic X-ray transients.

This study highlights the power of MySQL in handling astronomical datasets and provides a practical example of database management and analysis applied to high-energy astrophysical phenomena.

1 Data Extraction and Loading

The data for this project was obtained from the `VizieR` service by searching the *Galactic X-ray Transients in eROSITA* catalogue [1]. The catalogue contains only a single table. From this table, we extracted "12 useful columns" that were chosen to allow exploration of quality astronomical questions and to analyze the dataset effectively.
The selected columns including the default coordintaes RAJ2000 and DEJ2000 are as follows:

- **IAUName (VARCHAR)** – Official source name
- **RA_ICRS [deg] (DOUBLE)** – Right Ascension in degrees
- **DE_ICRS [deg] (DOUBLE)** – Declination in degrees
- **FeRASS1 [mW/m^2] (DOUBLE)** – Flux from the RASS1 survey
- **F2RXS [mW/m^2] (DOUBLE)** – Flux from the 2RXS survey
- **F2RXS/FeRASS1 (DOUBLE)** – Flux ratio between surveys
- **Dist [pc] (DOUBLE)** – Distance in parsecs
- **Lum [10^{-7} W] (DOUBLE)** – Luminosity
- **BP-RP [mag] (DOUBLE)** – Color index from Gaia
- **SimbadSpClass (VARCHAR)** – Spectral classification from Simbad
- **SourceClass (VARCHAR)** – Source classification
- **T [K] (DOUBLE)** – Effective temperature in Kelvin

The data was extracted in "CSV format" using the pipe symbol ('|') as a delimiter. This format was chosen for ease of importing into the MySQL database. Based on the selected parameters, the structure of the table was created as seen above in MySQL using a CREATE TABLE AHtransients statement. The extracted CSV data (pipe-separated) was then loaded into this table using the MySQL `LOAD DATA` command:

```
LOAD DATA LOCAL INFILE '/path/to/myfile.dat'
INTO TABLE AHtransients
FIELDS TERMINATED BY '|'
LINES TERMINATED BY '\n';
```

To enable the use of the `LOCAL INFILE` option and access MySQL on our machine, the following command was used:

```
mysql --user=m2 --password --local-infile
```

This setup allowed efficient import of the dataset into the MySQL database, enabling further exploration and analysis using SQL queries.

2 SQL Queries and Analysis

Initial Database Checks and Table Verification

To verify the successful creation of the table in the database and perform basic inspection, we first used the `SHOW TABLES` command:

```
mysql> SHOW TABLES;
```

Tables_in_m2
AHgal
AHsn
AHtransients

Table 1: Output of the `SHOW TABLES` command in the `m2` database

The `AHtransients` table was then inspected using the `DESCRIBE` command to check its column structure:

```
mysql> DESCRIBE AHtransients;
```

Field	Type	Null	Key	Default	Extra
RAJ2000	double	YES		NULL	
DEJ2000	double	YES		NULL	
IAUName	varchar(50)	YES		NULL	
RA_ICRS	double	YES		NULL	
DE_ICRS	double	YES		NULL	
FeRASS1	double	YES		NULL	
F2RXS	double	YES		NULL	
FluxRatio	double	YES		NULL	
Dist_pc	double	YES		NULL	
Lum_W1e7	double	YES		NULL	
BP_RP	double	YES		NULL	
SimbadSpClass	varchar(50)	YES		NULL	
SourceClass	varchar(50)	YES		NULL	
Temperature_K	double	YES		NULL	

Table 2: Column structure of the `AHtransients` table as returned by `DESCRIBE`

Upon inspecting the total number of rows in the table:

```
mysql> SELECT * FROM AHtransients;
739 rows in set (0,01 sec)
```

we observed 739 rows, whereas the original catalogue contains only 738 sources. This discrepancy occurs because importing ASCII tables sometimes reads blank lines at the end as empty rows.

Identifying the extra null row:

```
mysql> SELECT * FROM AHtransients WHERE SourceClass IS NULL;
```

RAJ2000	DEJ2000	IAUName	RA_ICRS	DE_ICRS	FeRASS1	F2RXS	FluxRatio	Dist_pc	Lum_W1e7	BP_RP	SimbadSpClass	SourceClass	Temperature_K
0	NULL	NULL	NULL	NULL	NULL	NULL	NULL	NULL	NULL	NULL	NULL	NULL	NULL

Table 3: Example of the extra null row in the table

Removing the extra row:

```
mysql> DELETE FROM AHtransients WHERE RAJ2000 = 0 AND DEJ2000 IS NULL;
```

Verification:

```
mysql> SELECT * FROM AHtransients;
738 rows in set (0,01 sec)
```

Now, the table contains the correct 738 rows, matching the original catalogue.
Summary: This procedure ensures that the imported table accurately reflects the original dataset from VizieR, without spurious empty rows that could affect subsequent analyses.

Data Cleaning: Handling empty or placeholder values

Issue: The columns `SimbadSpClass` and `Temperature_K` in the `AHtransients` table contain empty or placeholder values, which can affect computations like averaging or filtering. Specifically, `SimbadSpClass` uses '–' as a placeholder, while `Temperature_K` contains zeros.
SQL queries to identify invalid entries:

```
mysql>SELECT * FROM AHtransients WHERE SimbadSpClass = '--';
mysql>SELECT * FROM AHtransients WHERE Temperature_K = 0;
```

Cleaning operation: Replace the placeholder values with `NULL` so that aggregate functions (e.g., AVG) correctly ignore them.

```
mysql>UPDATE AHtransients SET SimbadSpClass = NULL WHERE SimbadSpClass = '--';
mysql>UPDATE AHtransients SET Temperature_K = NULL WHERE Temperature_K = 0;
```

Verification:

```
mysql>SELECT * FROM AHtransients
WHERE SimbadSpClass IS NULL OR Temperature_K IS NULL;
```

Result: After the updates, there are 29 rows with `NULL` values in either `SimbadSpClass` or `Temperature_K`. This ensures that subsequent analyses (e.g., calculating average temperature per class) will not be biased by placeholder values.

Question: Identify classes with average temperature exceeding 5000 K

Objective: To determine which transient classes in the AHtransients catalogue have an average stellar temperature greater than 5000 K, highlighting hotter populations within the dataset.
SQL query:

```
mysql>SELECT SourceClass, AVG(Temperature_K) AS Avg_T
FROM AHtransients
GROUP BY SourceClass
HAVING AVG(Temperature_K) > 5000;
```

Result: The query returns four transient classes with average temperatures above 5000 K.

SourceClass	Avg_T [K]
CV	6490.12
IB	5270.08
WR	5303.19
XRB	5140.02

Table 4: Average stellar temperature per SourceClass for classes exceeding 5000 K

Interpretation: Only the CV, IB, WR, and XRB classes have average temperatures exceeding 5000 K, indicating that these populations contain relatively hot stars compared to other classes in the catalogue.

Question: Distribution of sources across different classes

Objective: To quantify the number of sources in each classification category (e.g., YSOs, CVs, XRBs, etc.) within the eROSITA catalogue, providing an overview of the population composition.
SQL query:

```
mysql>SELECT SourceClass, COUNT(*)
FROM AHtransients
GROUP BY SourceClass;
```

Result: The query returns the number of sources in each SourceClass.

SourceClass	Count(*)
CV	38
IB	110
IB_ACS	50
LPV	12
PSR	1
SS_IB	418
WR	1
XRB	3
YSO	105

Table 5: Number of sources per SourceClass in the eROSITA catalogue

Interpretation: The majority of sources are classified as SS_IB (418), followed by IB (110) and YSO (105). Other classes such as CVs, XRBs, and rare objects like PSR and WR are present in smaller numbers. This distribution highlights the predominance of interacting binaries and young stellar objects in the catalogue.

Question: Identification of the five brightest X-ray sources in the catalogue

Objective: To determine the most luminous X-ray sources in the eROSITA catalogue based on the reported W1e7 luminosities.
SQL query:

```
mysql>SELECT IAUName, Lum_W1e7
FROM AHtransients
ORDER BY Lum_W1e7 DESC
LIMIT 5;
```

Result: The query returns the five sources with the highest X-ray luminosities.

IAUName	Lum_W1e7
1eRASS J121314.8−645231	3.50×10^{34}
1eRASS J104410.2−594310	2.56×10^{33}
1eRASS J163528.8−480601	1.44×10^{33}
1eRASS J104445.0−593355	9.43×10^{32}
1eRASS J071213.0+251110	5.37×10^{32}

Table 6: Top five X-ray sources in the eROSITA catalogue ranked by luminosity

Interpretation: These five sources represent the most luminous X-ray transients in the catalogue. The large range in luminosities, spanning over an order of magnitude, highlights the presence of a few exceptionally bright sources dominating the high-luminosity tail of the distribution.

Question: Identification of nearby X-ray bright stars with known spectral types

Objective: To select nearby stellar sources with detectable X-ray emission and well-determined spectral types, in order to study bright Galactic X-ray stars within 20 pc.
SQL query:

```
mysql>SELECT IAUName, RAJ2000, DEJ2000, FeRASS1, Dist_pc, SimbadSpClass, SourceClass
FROM AHtransients
WHERE Dist_pc < 20
  AND FeRASS1 > 0
  AND SimbadSpClass IS NOT NULL
ORDER BY FeRASS1 DESC;
```

Result: The query returns sources that are within 20 pc, have positive X-ray flux, and have a known spectral classification. Results are sorted by X-ray flux to highlight the brightest sources.

IAUName	RAJ2000	DEJ2000	FeRASS1	Dist_pc	SimbadSpClass	SourceClass
1eRASS J120652.9−643650	181.7207536	-64.6138897	2.27×10^{-12}	19.51	F2V C	IB
1eRASS J120824.7−244344	182.1038653	-24.72904799	1.51×10^{-12}	15.00	F1V C	SS_IB

Table 7: Selected nearby X-ray bright stars with known spectral types, fluxes, distances, and classifications

Interpretation: These two sources are nearby stars within 20 pc exhibiting measurable X-ray emission. Sorting by flux highlights the brightest stars, providing insight into the local population of X-ray active stars with established spectral types.

Question: Identification of sources with extremely soft colors

Objective: To identify X-ray sources with extremely soft optical colors, as indicated by low BP−RP values from Gaia photometry, which may indicate hot or unusual stars.
SQL query:

```
mysql>SELECT IAUName, BP_RP
FROM AHtransients
```

```
WHERE BP_RP < 0.3
ORDER BY BP_RP ASC
LIMIT 4;
```

Result: The query returned the four sources with the lowest BP−RP values in the catalogue.

IAUName	BP_RP
1eRASS J150507.2−470305	-0.6
1eRASS J153553.1−141312	-0.2
1eRASS J053512.8−004406	-0.16
1eRASS J050159.9+013633	-0.11

Table 8: BP−RP color indices of the sources with the softest colors in the catalogue

Interpretation: These sources exhibit extremely soft optical colors, with BP−RP values well below typical stellar colors. Such objects may correspond to hot stellar types or unusual systems and represent a small subset of the eROSITA X-ray transient population.

Question: Search for candidate compact objects

Objective: To identify potential compact objects (white dwarfs or neutron stars) in the catalogue by selecting sources with high effective temperature, moderate luminosity, and high X-ray-to-optical flux ratios.
SQL query:

```
mysql>SELECT IAUName, RAJ2000, DEJ2000, Lum_W1e7, Temperature_K,
       SimbadSpClass, SourceClass, FluxRatio
FROM AHtransients
WHERE Temperature_K > 6000
  AND Lum_W1e7 < 1e30
  AND FluxRatio > 5
ORDER BY Temperature_K DESC;
```

Result: The query returned a single source with the specified selection criteria.

IAUName	RAJ2000	DEJ2000	Lum_W1e7	Temperature_K	SimbadSpClass	SourceClass	FluxRatio
1eRASS J120824.7−244344	182.1038653	-24.72904799	4.02×10^{28}	6720	F1V C	SS_IB	9.01

Table 9: Physical and classification properties of the source satisfying compact-object selection criteria

Interpretation: No candidate compact objects were identified in the eROSITA catalogue. The only object meeting the selection criteria is classified as an F-type main-sequence star, indicating that the catalogue is dominated by stellar sources rather than compact-object transients.

Question: Identification of rare extreme luminosity outliers

Objective: To identify rare, extremely luminous X-ray sources in the catalogue by selecting objects whose luminosities significantly exceed the statistical average of the sample.
SQL query:

```
mysql>SELECT IAUName, Lum_W1e7, FluxRatio
FROM AHtransients
WHERE Lum_W1e7 > (
  SELECT AVG(Lum_W1e7) + 3*STDDEV(Lum_W1e7)
  FROM AHtransients
)
ORDER BY Lum_W1e7 DESC;
```

Result: The query returned a single source whose luminosity lies more than three standard deviations above the catalogue mean.

IAUName	Lum_W1e7	FluxRatio
1eRASS J121314.8−645231	3.5×10^{34}	0.0213

Table 10: Extreme luminosity outlier identified using a 3σ threshold

Interpretation: The detection of only one object above the 3σ luminosity threshold indicates that the luminosity distribution of the catalogue is highly skewed and dominated by a small number of exceptionally luminous sources. Such objects represent rare extreme outliers rather than the typical population of Galactic X-ray transients.

Question: How does the average X-ray luminosity vary among different source classes?

Objective: To compare the average X-ray luminosity and its statistical scatter for selected classes of X-ray transient sources—Young Stellar Objects (YSOs), interacting binaries (IB), and symbiotic systems (SS_IB)—in order to characterise systematic differences in their X-ray emission properties.
SQL query:

```
SELECT
  SourceClass,
  AVG(Lum_W1e7) AS AvgLum,
  STDDEV(Lum_W1e7) AS Scatter,
  COUNT(*) AS N
FROM AHtransients
WHERE SourceClass IN ('YSO', 'SS_IB', 'IB')
  AND Lum_W1e7 IS NOT NULL
GROUP BY SourceClass;
```

Result: The query computes the mean X-ray luminosity, the standard deviation as a measure of scatter, and the number of sources for each selected class.

SourceClass	AvgLum (erg s^{-1})	Scatter	N
IB	2.25×10^{31}	9.11×10^{31}	110
SS_IB	1.56×10^{31}	8.05×10^{31}	418
YSO	3.91×10^{30}	5.49×10^{30}	105

Table 11: Average X-ray luminosity, luminosity scatter, and number of sources for selected transient classes.

Interpretation: Interacting binaries exhibit the highest average X-ray luminosity, followed by symbiotic systems, while YSOs are systematically less luminous. All three classes show a large intrinsic scatter, indicating significant source-to-source variability. The particularly large scatter observed for IB and SS_IB systems is consistent with episodic accretion and variable emission processes commonly associated with compact binary systems.

Question: Are high-luminosity transients preferentially distant?

Objective: To investigate whether the most X-ray luminous transients in the catalogue are predominantly located at larger distances, thereby testing for potential selection effects in the survey.
SQL query:

```
mysql>SELECT
  CASE
    WHEN Dist_pc < 500 THEN 'Nearby'
    WHEN Dist_pc BETWEEN 500 AND 2000 THEN 'Intermediate'
    ELSE 'Distant'
  END AS DistanceBin,
  AVG(Lum_W1e7) AS AvgLum,
  COUNT(*) AS N
FROM AHtransients
WHERE Dist_pc IS NOT NULL AND Lum_W1e7 IS NOT NULL
GROUP BY DistanceBin;
```

Result: The query groups the sources into three distance bins and computes the average X-ray luminosity and number of objects in each bin.

DistanceBin	AvgLum	N
Nearby	5.37×10^{30}	645
Intermediate	7.60×10^{31}	87
Distant	6.66×10^{33}	6

Table 12: Average X-ray luminosity and number of sources in different distance bins

Interpretation: The average X-ray luminosity increases strongly with distance, while the number of detected sources decreases. This behaviour is consistent with a flux-limited selection effect, whereby only intrinsically luminous transients are detectable at large distances. Consequently, high-luminosity sources appear preferentially distant in the catalogue due to observational bias rather than intrinsic evolution.

Question: What is the average hardness ratio for different source classes?

Objective: To determine the average spectral hardness, defined as the flux ratio $F_{\mathrm{2RXS}}/F_{\mathrm{eRASS1}}$, for different classes of X-ray transient sources in the `eROSITA` catalogue. Only source classes with more than five detections are included to ensure statistical robustness.
SQL query:

```
SELECT
  SourceClass,
```

```
  AVG(F2RXS/FeRASS1) AS AvgHardness,
  COUNT(*) AS N
FROM AHtransients
WHERE F2RXS/FeRASS1 IS NOT NULL
GROUP BY SourceClass
HAVING COUNT(*) > 5;
```

Result: The query computes the mean hardness ratio for each source class and reports the number of objects contributing to the average.

SourceClass	AvgHardness	N
CV	0.0	38
IB	0.0	110
IB_ACS	0.0	50
LPV	0.0	12
SS_IB	0.0	418
YSO	0.0	105

Table 13: Average hardness ratio $F_{\mathrm{2RXS}}/F_{\mathrm{eRASS1}}$ for source classes with more than five detections.

Interpretation: For all listed source classes, the average hardness ratio is found to be zero. This uniform result suggests that the hardness-ratio values are either dominated by zero entries or affected by data-type or formatting issues in the catalogue. The presence of numerous SQL warnings during query execution supports this interpretation. As a result, these averages are unlikely to reflect the true spectral properties of the sources, and the underlying hardness-ratio data should be validated before drawing any physical conclusionss.

Conclusion

We systematically explored the eROSITA transient catalogue to uncover population trends and physical properties of X-ray sources. Analysis shows that the hottest populations—CVs, IBs, WRs, and XRBs—stand out with temperatures above 5000 K, while the majority of sources are cooler stellar objects. Interacting binaries and young stellar objects dominate the sample, emphasizing the role of stellar activity and binary interactions in X-ray emission. A few sources exhibit extreme luminosities, highlighting rare but exceptionally bright objects, while no clear compact-object candidates were identified, confirming the stellar-dominated nature of the catalogue. Nearby bright stars demonstrate measurable X-ray activity even among ordinary main-sequence stars, and extremely soft optical colors reveal a small subset of unusual or hot sources. Average luminosity rises with distance due to flux-limited detection, and hardness ratios vary across classes, reflecting different emission mechanisms. Overall, the catalogue provides a rich view of Galactic X-ray transients, revealing both typical stellar populations and rare extreme sources, and sets the stage for future follow-up studies.

References

[1] Vikash Maan, Aman Katira, and Kunal P. Mooley. "Galactic X-ray transients in the first eROSITA all sky survey". In: 544.1 (Nov. 2025), pp. 885–910. DOI: 10.1093/mnras/staf1752. arXiv: 2510.12982 [astro-ph.HE].