



Ethnomedicinal plants used by traditional healers of North Tripura district, Tripura, North East India



Prakash Roy Choudhury^a, Manabendra Dutta Choudhury^a, Sanjoy Singh Ningthoujam^b,
Dipan Das^a, Deepa Nath^c, Anupam Das Talukdar^{a,*}

^a Department of Life Science & Bioinformatics, Assam University, Silchar, Assam, India

^b Department of Botany, Ghanapriya Women's College, Imphal, Manipur, India

^c Department of Botany & Biotechnology, Karimganj College, Karimganj, Assam, India

ARTICLE INFO

Article history:

Received 24 December 2014

Received in revised form

3 March 2015

Accepted 3 March 2015

Available online 17 March 2015

Keywords:

Ethnomedicine

North Tripura district

Medicinal plant

India

Quantitative analysis

ABSTRACT

Ethnopharmacological relevance: Information about ethnomedicinal plants used by traditional healers in Tripura, India is very much limited. As there is rapid erosion of traditional knowledge in the state, needs has arisen to document the diversity of medicinal plants, their preparation and associated diseases.

Aim of the study: Attempts have been made to explore the herbal medicines used by the traditional healers belonging to the various communities in North Tripura district of Tripura, India and document the associated traditional knowledge on the utilisation of medicinal plants.

Materials and methods: The study was based on ethnomedicinal field survey covering a period of 1 year from February 2012 to February 2013. The ethnomedicinal information was collected by using semi-structured questionnaires from different healers and knowledge holders. Collected data were analysed through informant consensus factor, Index of Specialisation and Relative frequency of citation to determine culturally significant plants.

Results: A total of 75 species of plants under 68 genera belonging to 43 families were collected during the study for the treatment of 15 disease categories. Leaves were the most frequently used plant parts and most of the medicines were prepared in the form of extract and administered orally. F_{IC} values of the present study indicated that there was a high agreement in the use of plants in the treatment of digestive system disorders and respiratory system disorders among the healers. Plants having high ISu are *Justicia adhatoda*, *Pajanelia longifolia*, *Catharanthus roseus* etc.

Conclusion: The present study highlighted certain species having higher priority indices for further phytochemical investigation. Counselling of traditional health practitioners should be taken into consideration in order to smooth continuation and extension of traditional medical knowledge and practice for ensuring safe and effective therapy.

© 2015 Elsevier Ireland Ltd. All rights reserved.

1. Introduction

Ethnomedicinal plants contribute a significant role in accomplishing the primary health care needs of the population in developing countries (Rai et al., 2000), particularly as alternative and complementary *materia medica*. According to World Health Organisation (WHO), 80% of world's population in developing countries rely on herbal medicines, traditional treatments, and traditional practitioners for the main source of health care, and sometimes the only source of care as because it is close to home, accessible and affordable (Akerle, 1984; Shil et al., 2014). In many societies, traditional medicine systems are culturally acceptable

with high acceptance level among the concerned people. The affordability of most traditional medicines makes them attractive at a time of soaring health-care costs and nearly universal austerity (Calixto, 2005). A number of people with diverse cultural backgrounds reside in India and practice their own system of traditional medicine for primary healthcare through generations. Reportedly, 70–80% of the rural Indians use non-allopathic (Ayurveda, Siddha, Unani and Homoeopathy) medicines for their healthcare and herbs constitute a major ingredient of these alternative systems of medicine (Gogtay et al., 2002).

The North East (NE) India, one of the richest reservoirs of plant diversity in India (Mao et al., 2009), encompasses a sizeable ethnic population of about 8 million (Sharma et al., 2014) with its conterminous eight sister states. The state Tripura, one of the states of NE India, with a total geographical area of 10491.69 km² is located between 22° 56' and 24° 32' N latitude and 90° 09' to 92° 20' E longitude. The state

* Corresponding author. Tel.: +91 9401416452; fax: 91 3842 270920.

E-mail address: anupam@bioinfoaus.ac.in (A. Das Talukdar).

is the home for various ethnic groups namely Tripuri, Bengali, Debbarma, Reang, Naotia, Chakma, Bhil, Bhutia, Chaimal, Garo, Halam, Khasia, Kuki, Lepcha, Lushai, Mog, Munda, Orang, Santhal, Uchoi, Kolo and many other smaller groups. Every community in the state have a rich heritage of medicinal plants and their usage along with their own system of preserving this tradition (Deb et al., 2013). These knowledge ranges from the therapeutic treatment of commonly occurring household remedies to specialized treatment unique to certain traditional healers. Traditional healers following Ayurvedic system of medicine are locally called *Kaviraja* while those following community-specific folk medicine are known as *Ojhas* respectively. Although there are rapid erosion of ethnomedicinal knowledge among these communities, with urbanisation and introduction of western medicine system, many of them in different pockets of the state are still practicing the traditional health practices. It has been felt that documentation of the traditional knowledge from different communities would help in conserving these rich heritages of knowledge and also in incorporating scientific knowledge to the practices and validation of such therapies.

Different research workers in different time periods have put tremendous efforts for investigating the ethnobotanical aspects of Tripura's diverse vegetation. Mention may be made of pioneering works such as Deb (1968, 1975a, 1975b, 1975c, 1976, 1978a, 1978b, 1981a, 1981b, 1983), Singh et al. (1997), Devbarma (1976), Majumdar et al. (2006), Shil and Choudhury (2009), Das et al. (2009), Das and Choudhury (2010), Sen et al. (2011), Sharma et al. (2013); Sharma et al. (2014), Shil et al. (2014) among others. In spite of such efforts, considerably less attention has been paid to the North Tripura District of Tripura by these ethnobotanists for exploring the ethnomedicinal resources.

Considering these limitations in exploring the diversity of medicinal plants vis-à-vis to the traditional medicine, the present study was attempted to document the ethnomedicinal plant wealth of North Tripura district of Tripura and their use by different communities inhabiting in the district.

2. Materials and methods

2.1. Study area

The study was conducted in North Tripura district of Tripura (Fig. 1) which has a geographical area of 1422.19 km² and located at latitude of 24° 36' and at longitude of 92° 19'. The district is divided into three sub-divisions, namely, Dharmanagar, Kanchanpur and Panisagar and has a total population of 422,370 (approx). Main ethnic communities of the districts are Bengali, Tripuri, Halam, Reang, Chakma, Manipuri and Debbarma following Hinduism, Christianity and Buddhism.

2.2. Questionnaire survey and ethnobotanical data collection

The ethnomedicinal field survey was carried out over a period of one year in three sub-divisions of North Tripura district, Tripura during February 2012 to February 2013. Semi-structured questionnaires were administered to collect ethnobotanical data from the informants (Ningthoujam et al., 2013). During the survey, 30 informants were interacted out of which 20 are professional. Most of our information were provided by these 20 informants (male: 13; female: 7), out of which 12 were non-healers (they are having the knowledge of medicinal plant but do not practicing it) and rests are *Kavirajas* and *Ojhas*. All the informants belonged to the 7 different communities and were born and reared in their respective villages. Traditional healers were practicing for years and derived their knowledge from their ancestry or learnt through traditional institutions. Documentation about the informants' community, age, experience, occupation (apart from practicing traditional healing), literacy and social acceptability of

their healing ability were also done. Most of the informants were found to be illiterate and school dropouts. Prior informed consent was obtained before collecting the information.

On the basis of information provided by the informants, plant samples were collected from fields and photographs of the plants in flowering/fruitlet conditions were taken. Some specimens were collected from the healers' private collections and home gardens as these plants are no more found in the wild condition during the survey. Plants collected from the fields were cross-checked with the healers. The plants were initially identified at Department of Life Science and Bioinformatics, Assam University by discussing mainly from the "Flora of Tripura State" (Deb, 1981a, 1983) and books covering nearby regions namely Flora of British India (Hooker, 1872–1897) and Flora of Assam (Kanjalil et al., 1934, 1936, 1938, 1940). Plants were again verified with the records of the Botanical Survey of India (BSI), North Eastern Circle, Shillong. The herbarium specimens were deposited at the Department of Life Science and Bioinformatics, Assam University, Silchar. Scientific names were standardised by cross-checking with the IPNI (www.ipni.org) and the Plant List (www.theplantlist.org) databases.

2.3. Analysis

2.3.1. F_{IC} (Informant consensus factor)

Informant Consensus Index was one of popular quantitative approaches for identifying the relative importance of medicinal plants with the ailment categories in a particular culture. The present index was initially developed by Trotter and Logan (1986) and later on modified by Heinrich et al. (1998). The present index calculated on the basis of the following equation:

$$F_{IC} = N_{ur} - N_t / (N_{ur} - 1),$$

where N_{ur} stands for the number of use reports for a particular use category and N_t stands for the number of taxa used for a particular ailment category by all informants. The indices could reflect the homogeneity in the use of plants in the ailment categories among the informants of the study area. As many species may be associated with the same disease, this factor becomes significant tool for determining the most used plant species for treating a particular ailment. A higher F_{IC} value indicates the use of relatively few plants by the informants in the treatment of a particular ailment category whereas a lower F_{IC} value indicates that there are disagreement among the informants with regard to use of a particular plant for treating a particular ailment category.

2.3.2. IS_u (Index of Specialisation unit)

In the present work, evaluation of the ethnobotanical data was performed with modified fidelity level index (Friedman et al., 1986; Bruni et al., 1997) which have been applied in determining the importance of medicinal recipes (Ningthoujam et al., 2013). The primary use of the recipe was used as a fidelity index which is expressed as a percentage of the citation for that recipe (% C_m). The modified index called the "Index of Specialisation by the main use" (IS_u) was used to determine the consensus for the specific medical uses along with type of preparation and application. This index was calculated with the following formula:

$$IS_u = (\%C_m \times \%C_t) / 100$$

where IS_u is the Index of Specialisation by the main use, % C_m is the percentage of citation for a particular drug, and % C_t is the percentage of the total citations.

2.3.3. Relative frequency of citation (RFC)

The RFC is obtained by the equation $RFC = FC/N$ where, FC is the frequency of citation i.e., the number of informants who mention




Fig. 1. Map showing the study area.

the use of the species and N is the total number of informants interviewed in the survey (Tardío and Pardo-de-Santayana, 2008).

3. Results

In the survey, a total of 75 plant species have been reported by 20 traditional healers for treatment of 15 different of disease categories. Herbal recipes of these medicinal plants are tabulated with scientific name and family followed by vernacular name, RFC, disease, parts used, preparation, application, number of citation, number of use types, Ct, %Ct, Cm, %Cm, ISu, other uses and similar use (Table 1).

Among the 20 interviewees, 08 people are traditional health practitioners (THPs) and rests are indigenous non-healers. Major informants were male (65%). There were a high percentage of interviewees who were around 60–70 years (35%) old followed by informants who were more than 70 years (20%) old. The majority of interviewees were illiterate (70%). Experience wise most of the interviewees had 10–20 years of experience (45%) of traditional healing practices (Fig. 2).

In the present study a total of 75 plant species belonging to 43 plant families are identified as having medicinal values used by THPs of North Tripura district. The family Asteraceae and Rutaceae (6 species each) represents the highest number of species, followed by Lamiaceae and Lauraceae (5 species each), Euphorbiaceae (4 species) and Apocynaceae (3 species).

In terms of percentage of plant parts used by the informants, the percentages are as follows, leaf 70.65%, bark 7.6%, fruit 5.43%, root 4.89%, stem 4.34%, whole plant 3.26%, seed 1.63%, latex 1.08%, rhizome and milky juice 0.54% each.

Respiratory system disorders had the highest F_{IC} of 0.85 followed by digestive system disorders (0.84), endocrine system disorders and infection (0.82) (Table 2).

Relative frequency of citation (RFC) of each species has been evaluated from the available information. In the present study RFC of plant species varied between 1 and 0.1. *Litsea glutinosa* of family Lauraceae has shown the highest RFC of 1.0 followed by *Pajanelia longifolia* (0.95), *Catharanthus roseus* (0.9), *Justicia adhatoda* (0.9), and *Ocimum tenuiflorum* (0.9).

The Index of Specialisation (ISu) of medicinal plant species in the present study ranged from 0.12 to 1.78. *J. adhatoda* has shown highest ISu of 1.78 followed by *P. longifolia* (1.65), *C. roseus* (1.52), *Centella asiatica* (1.39), *Oroxylum indicum* (1.39), *Carica papaya* (1.39), *L. glutinosa* (1.14), *Annona reticulata* (1.14), *Citrus lemon* (1.14), *Hyptis suaveolens* (1.14), *Lagerstroemia speciosa* (1.14), *Mimosa pudica* (1.14), *Pandanus foetidus* (1.14), *Justicia gendarussa* (1.01), *Bryophyllum pinnatum* (1.01), *Enhydra fluctuans* (1.01).

4. Discussion

The present study indicated that traditional healers have substantial knowledge on specificities of different medicinal plants according to disease types, which is also corroborated with earlier studies (Das and Choudhury, 2010). Differential values of FIC in different categories reflected the preponderances of the healer's viewpoints as well as common ailments facing the communities covered. Disease categories with highest values of FIC such as Respiratory System Disorders and Digestive System Disorders are associated with common ailments like cough, sinus, indigestion, and stomach ache. Moreover, another category of Infection exhibiting

Table 1
Medicinal plants of North Tripura district of Tripura state against different ailments.

Plant name and family	Vernacular name	Voucher no.	RFC	Main disease	Parts used	Preparation	Application	Number of citations per preparation	No. of uses type	Ct	%Ct	Cm	%Cm	ISu	Other uses and mode of application	Relevant ethnobotanical/ pharmacological citation
<i>Acmella paniculata</i> (Wall. ex DC.) R.K. Jansen	Ansa (H)	AUS2507	0.35	Paralysis	Leaf	Crushed	Rubbed	3	2	4	0.50	3	75	0.38	Eczema (rubbed); Headache (drink)	(Hossain et al., 2014)
Asteraceae				Fever	Leaf	Decoction	Drink	2	1	2	0.25	2	100	0.25		
<i>Aegle marmelos</i> (L.) Correa	Bael (H, B, M)	AUS2508	0.65	Diarrhoea	Fruit	Decoction	Drink	3	3	5	0.63	3	60	0.38	Indigestion (Decoction, drink); weak heart (Decoction, drink)	(Shoba and Thomas, 2001; Brijesh et al., 2009; Arul et al., 2004, 2005)
Rutaceae				Knee Pain	Fruit	Crushed	Rubbed	2	1	2	0.25	2	100	0.25		
				Fever	Fruit	Crushed	Drink	5	1	5	0.63	5	100	0.63		
				Fever	Leaf	Crushed	Drink	1	1	1	0.12	1	100	0.12		
				Asthma	Fruit	Burnt	Inhale	1	1	1	0.12	1	100	0.12		
<i>Aerva javanica</i> (Burm. F.) Juss. ex Schult.	Bishalyakarani (B)	AUS2509	0.40	Joint Pain	Leaf	Crushed	Rubbed	2	2	8	1.01	6	75	0.76		(Qureshi and Bhatti, 2008)
Amaranthaceae				Cut and Wounds	Leaf	Crushed	Rubbed	6								
<i>Ageratum conyzoides</i> (L.) L.	Bon tulsi (B),	AUS2510	0.55	High BP	Leaf	Crushed	Drink	5	2	8	1.01	5	62.5	0.63	Angina (Crushed, drink)	(Ming, 1999; Rajendran et al., 2004)
Asteraceae	Samnama (R)			Fever	Flower	Crushed	Drink	3	1	3	0.38	3	100	0.38		
<i>Alocasia macrorrhizos</i> (L.) G.Don	Anman kachu (B)	AUS2511	0.35	Arthritis	Rhizome	Crushed	Rubbed	7	1	7	0.89	7	100	0.89		(Nadkarni and Basu, 1996; Rahman et al., 2011; Rahmatullah et al., 2010b)
Araceae																
<i>Aloe vera</i> (L.) Burm.f.	Kumari (R), gritkumari (B)	AUS2512	0.40	Jaundice	Stem	Crushed	Drink	5	1	5	0.63	5	100	0.63		(Vogler and Ernst, 1999; Surjushe et al., 2008)
Liliaceae				Liquiria	Leaf	Decoction	Drink	3	1	3	0.38	3	100	0.38		
<i>Alstonia scholaris</i> (L.) R. Br.	Lawthong (H)	AUS2513	0.8	Diarrhoea	Bark	Decoction	Drink	6	2	9	1.14	6	66.6667	0.76		(Shah et al., 2010; Das et al., 2003; Attar-Rahman et al., 1985)
Apocynaceae				Ulcer	Milky Juice	Decoction	Drink	3	1	3	0.38	3	100	0.38		
				Bone Fracture	Leaf	Crushed	Rubbed	4	1	4	0.50	4	100	0.50		
<i>Amaranthus spinosus</i> L.	Katadugi (B)	AUS2514	0.55	Snake Bite Stone	Bark Leaf	Decoction Decoction	Drink Drink	3 7	2	11	1.39	7	63.6364	0.89	Stomachache (Decoction, drink)	(Ballabh et al., 2008)
Amaranthaceae																
<i>Annona reticulata</i> L.	Atafal (B)	AUS2515	0.50	Cuts	Leaf	Crushed	Rubbed	9	1	9	1.14	9	100	1.14		(Chavan et al., 2012)
Annonaceae				Infestation of Cattle	Leaf	Decoction	Vapour	1	1	1	0.12	1	100	0.12		
<i>Archidendron clypearia</i> (Jack) I.C. Nielsen ^a	Loapatto (C)	AUS2516	0.45	Dysentery	Bark	Decoction	Drink	6	2	9	1.14	6	66.6667	0.76	Stomachache (Decoction, drink)	
Mimosaceae																
<i>Averrhoa carambola</i> L.	Kapra (B)	AUS2517	0.60	Jaundice	Leaf	Crushed	Pellet	7	1	7	0.89	7	100	0.89		(Borah et al., 2006; Mia et al., 2007; Ripunjoy, 2013)
Averrhoaceae				Dysentery	Leaf	Decoction	Drink	5	1	5	0.63	5	100	0.63		
<i>Azadirachta indica</i> A. Juss.	Neem (B)	AUS2518	0.65	Indigestion	Leaf	Decoction	Drink	2	3	9	1.14	4	44.4444	0.50	Fever, TB (Decoction, drink)	(Muthu et al., 2006; Khosla et al., 2000; Chattopadhyay et al., 1987; Kar et al., 2003)
Meliaceae				Diabetes	Leaf	Crushed	Drink	3	1	3	0.38	3	100	0.38		
				Allergy	Leaf	Crushed	Rubbed	1	1	1		1	100	0		
	Payan (B)	AUS2519	0.60	Cough	Leaf	Decoction	Drink	3	2	7	0.89	4	57.1429	0.50		

<i>Begonia roxburghii</i> A. DC.																Fever (Decoction, drink)	(Yogendra Kumar, 2008; Gupta, 2006; Srivastava 2010a)
Begoniaceae				Allergy	Leaf	Crushed	Rubbed	5	1	5	0.63	5	100	0.63			
<i>Bryophyllum pinnatum</i> (Lamk.) Oken	Gorol (B)	AUS2520	0.60	Dysentery	Leaf	Crushed	Drink	4	1	4	0.50	4	100	0.50			(Dalziel, 1937; Kamboj and Saluja, 2009)
Crassulaceae				Stone	Leaf	Decoction	Drink	8	1	8	1.01	8	100	1.01			
<i>Cajanus cajan</i> (L.) Millsp.	Arhar (B), domurshumi (C)	AUS2521	0.50	Jaundice	Leaf	Crushed	Pellet	7	1	7	0.89	7	100	0.89			(Kong et al., 2009; Morton, 1976; Akinloye and Olaniyi, 2011)
Papilionaceae				Allergy	Leaf	Crushed	Rubbed	3	1	3	0.38	3	100	0.38			
<i>Calotropis gigantea</i> (L.) Dryand.	Akanda (B)	AUS2522	0.20	Urinary Troubles	Leaf	Decoction	Drink	4	1	4	0.50	4	100	0.50			(Misra et al., 1993)
Apocynaceae																	
<i>Carica papaya</i> L.	Koiphol (B, DB, H)	AUS2523	0.60	Teeth Infection	Leaf	Crushed	Rubbed	9	2	11	1.39	9	81.81	1.14			(Rahmatullah et al., 2010b)
Caricaceae				Bone Fracture	Leaf	Crushed	Rubbed	2						0			
<i>Catharanthus roseus</i> (L.) G. Don.	Nayantara (B)	AUS2524	0.9	Arthritis Diabetes	Root Leaf	Crushed Decoction	Rubbed Drink	1 12	1	12	1.52	12	100	1.52			(Ghosh and Gupta, 1980; Banerjee and Basu, 1991; Singh et al., 2001; Kirtikar and Basu, 1933; Nammi et al., 2003; Rasineni et al., 2010)
Apocynaceae				Cut and Wounds	Leaf	Crushed	Rubbed	6	1	6	0.76	6	100	0.76			
<i>Centella asiatica</i> (L.) Urb.	Thunimankuni (B, M)	AUS2525	0.85	Gastric	Leaf	Crushed	Drink	11	1	11	1.39	11	100	1.39			(Cheng and Koo, 2000; Sairam et al., 2001; Chatterjee et al., 1992; Guo et al., 2004)
Apiaceae				Eye Infection	Leaf	Decoction	Drink	6	1	6	0.76	6	100	0.76			
<i>Cinnamomum verum</i> J. Presl	Dalchini (B, H)	AUS2526	0.15	Asthma	Bark	Crushed	Rubbed	1	1	1	0.12	1	100	0.12			(Khare, 2007; Mukti et al., 2012)
Lauraceae				Asthma	Bark	Decoction	Drink	2	1	2	0.25	2	100	0.25			
<i>Cissus quadrangularis</i> L.	Harkuchla (B)	AUS2527	0.10	Bone Fracture	Leaf	Crushed	Rubbed	2	1	2	0.25	2	100	0.25			(Shirwaikar et al., 2003)
Vitaceae																	
<i>Citrus limon</i> (L.) Osbeck	Lebu (B, H)	AUS2528	0.60	Diabetes	Leaf	Decoction	Drink	9	1	9	1.14	9	100	1.14			(Wagh et al., 2011; Naim et al., 2012)
Rutaceae				Dysentery	Fruit	Crushed	Drink	3	1	3	0.38	3	100	0.38			
<i>Citrus maxima</i> (Burm.) Merr.	Jambura (B, H)	AUS2529	0.60	Indigestion	Fruit	Crushed	Drink	3	2	10	1.27	7	70	0.89			(Bhandurge et al., 2010; Kundusen et al., 2011)
Rutaceae				Diabetes	Leaf	Crushed	Drink	7									
<i>Clausena heptaphylla</i> (Roxb.) Wight and Arn	Kukra (B)	AUS2530	0.35	Allergy Indigestion	Leaf Leaf	Crushed Crushed	Drink Rubbed Pellet	2 5	1 1	2 5	0.25 0.63	2 5	100 100	0.25 0.63			(Nath and Choudhury, 2010)
Rutaceae				Stone	Leaf	Decoction	Drink	2	1	2	0.25	2	100	0.25			
<i>Clerodendrum cordatum</i> D. Don	Bhati (B)	AUS2531	0.45	Jaundice	Leaf	Crushed	Rubbed	4	2	5	0.63	4	80	0.50			(Shukla et al., 2013)
Lamiaceae				Gastric Inflammation	Leaf	Decoction	Drink	3	2	4	0.50	3	75	0.38			
<i>Coccinia grandis</i> (L.) Voigt	Telakuchi (B)	AUS2532	0.40	Inflammation Diabetes	Leaf Leaf	Decoction Crushed	Drink Rubbed Drink	1 1 6	1	6	0.76	6	100	0.76			(Venkateswaran and Pari, 2003; Shibib et al., 1993; Umamaheswari and Chatterjee, 2008)
Cucurbitacea				Cut and Wounds	Leaf	Crushed	Rubbed	2	1	2	0.25	2	100				
<i>Cocos nucifera</i> L.	Narikul (B)	AUS2533	0.55	Teeth Infection	Leaf	Crushed	Rubbed	6	2	11	1.39	6	54.54	0.76			(Rahmatullah et al., 2010a; Yong et al., 2009)
Arecaceae				Inflammation	Leaf	Crushed	Rubbed	5									
<i>Conarus paniculatus</i> Roxb. ^a	Uroshichak (C)	AUS2534	0.30	Diarrhoea	Leaf	Decoction	Drink	5	1	5	0.63	5	100	0.63			

Table 1 (continued)

Plant name and family	Vernacular name	Voucher no.	RFC	Main disease	Parts used	Preparation	Application	Number of citations per preparation	No. of uses type	Ct	%Ct	Cm	%Cm	ISu	Other uses and mode of application	Relevant ethnobotanical/ pharmacological citation
Connaraceae				Stomach-Ache	Leaf	Crushed	Drink	1	1	1	0.12	1	100	0.12		
<i>Croton bonplandianum</i> Baill. ^a	Japur (T)	AUS2535	0.35	Indigestion	Root	Decoction	Drink	7	1	7	0.89	7	100	0.89		
Euphorbiaceae																
<i>Cynodon dactylon</i> (L.) Pers ^a	Durba (B)	AUS2536	0.35	Indigestion	Leaf	Decoction	Drink	4	1	4	0.50	4	100	0.50		
Poaceae				Semen Loss	Leaf	Crushed	Drink	3	1	3	0.38	3	100	0.38		
<i>Cuscuta reflexa</i> Roxb.	Sunnalata (H),	AUS2537	0.65	Jaundice	Leaf	Decoction	Drink	2	5	12	1.52	5	41.66	0.63		(Siwakoti and Siwakoti, 2000; Hossan et al., 2009; Patel et al., 2012)
Convolvulaceae	Kuchilalata (B)			Jaundice	Stem	Decoction	Drink	2								
				Gastric	Leaf	Decoction	Drink	1								
				Urinary Troubles	Stem	Decoction	Drink	5								
<i>Datura stramonium</i> L.	Dhutra (B)	AUS2538	0.40	Ear Pain	Leaf	Decoction	Drink	2								
Solanaceae				Knee Pain	Leaf	Crushed	Rubbed	5	2	8	1.01	5	62.5	0.63		(Gaire and Subedi, 2013)
<i>Dendrophthoe falcata</i> (L.f.) Ettingsh.	Gaschotta (C), amburus (B),	AUS2539	0.45	Inflammation	Leaf	Crushed	Rubbed	3								
Loranthaceae				Cough	Leaf	Decoction	Drink	4	2	7	0.89	4	57.1429	0.50		(Pattanayak and Mazumder, 2010; Rajendran et al., 2004)
				Cough Cut and Wounds	Root	Decoction	Drink	3								
<i>Dichrocephala integrifolia</i> (L.f.) Kuntz. ^a	Bortora (C)	AUS2540	0.25	Joint Pain	Leaf	Crushed	Rubbed	2	1	2	0.25	2	100	0.25		
Asteraceae																
				Skull Hardening	Leaf	Crushed	Rubbed	1								
<i>Diplazium esculentum</i> (Retz.) Sw.	Bon paloi (H, B)	AUS2541	0.35	Piles	Leaf	Crushed	Drink	1	1	1	0.12	1	100			
Athyriaceae				Indigestion	Leaf	Crushed	Drink	1	1	1	0.12	1	100	0.12		(Deb et al., 2009; Angami et al., 2006; Roy et al., 2013)
				Bone Fracture	Whole Plant	Crushed	Rubbed	2	2	6	0.76	4	66.6667	0.50		
<i>Dracaena elliptica</i> Thunb. & Dalm. ^a	Hodorotengait (C)	AUS2542	0.45	Arthritis	Leaf	Crushed	Rubbed	4								
Asparagaceae				Small Pox	Stem	Crushed	Rubbed	7	1	7	0.89	7	100	0.89		
<i>Enhydra fluctuans</i> Lour.	Helenchak (C)	AUS2543	0.75	Body Pain	Stem	Decoction	Drink	2	1	2	0.25	2	100	0.25		
Asteraceae				Indigestion	Leaf	Decoction	Drink	8	2	12	1.52	8	66.6667	1.01		(Sannigrahi et al., 2011; Sannigrahi et al., 2010)
				Nervous Disorder	Leaf	Decoction	Drink	4								
<i>Euphorbia neriifolia</i> L.	Shairapul (H), Shibgach (C)	AUS2544	0.55	Inflammation	Leaf	Crushed	Rubbed	3	1	3	0.38	3	100	0.38		
Euphorbiaceae				Pneumonia	Whole Plant	Decoction	Drink	7	1	7	0.89	7	100	0.89		(Jahan et al., 2011; Pokharen et al., 2011)
<i>Ficus hirta</i> Vahl ^a	Tamangaddu (C)	AUS2545	0.55	Allergy	Latex	Ointment		4	1	4	0.50	4	100	0.50		
Moraceae				Stomach-Ache	Leaf	Decoction	Rubbed	7	1	7	0.89	7	100	0.89		
<i>Ficus pumila</i> L.	Ludisharbuho (C)	AUS2546	0.30	Snake Bite	Root	Crushed	Rubbed	4	2	9	1.14	5	55.5556	0.63		
Moraceae				Diabetes	Leaf	Decoction	Drink	1	1	1	0.12	1	100	0.12		(Leong et al., 2008; Liao et al., 2012)
<i>Floscopa scandens</i> Lour. ^a	Panneihedugi (C)	AUS2547	0.6	Body Pain	Leaf	Crushed	Rubbed	5								
				Indigestion	Leaf	Crushed	Rubbed	7	2	10	1.27	7	70	0.89		

Commelinaceae				Sore Eye	Stem	Crushed	Drop	2	1	2	0.25	100	0.25	
				Bone Fracture	Leaf	Crushed	Rubbed	3						
<i>Hibiscus rosa-sinensis</i> L.	Jaba (B)	AUS2548	0.70	Dysentery	Leaf	Decoction	Drink	7	3	14	1.78	7	50	(Ross, 1999; Kim et al., 2003)
Malvaceae				Arthritis	Root	Decoction	Drink	3						
				Liver Troubles	Flower	Decoction	Drink	4						
<i>Houttuynia cordata</i> Thunb.	Toppaieigkong (M), Kokpanikouo (H)	AUS2549	0.40	Appetite Loss	Leaf	Decoction	Drink	5	2	8	1.01	5	62.5	(Kumar et al., 2014; Srivastava et al., 2010b)
Saururaceae				Body Pain	Leaf	Decoction	Drink	3						
<i>Hyptis suaveolens</i> Piot.	Tukma (B)	AUS2550	0.60	Liver Troubles	Leaf	Decoction	Drink	9	1	9	1.14	9	100	(Babalola et al., 2011; Ghaffari et al., 2012)
Lamiaceae				Allergy	Leaf	Crushed	Rubbed	3	1	3	0.38	3	100	
<i>Ipomoea aquatica</i> Forssk.	Kalmi (B,H) Banialochi (R)	AUS2552	0.45	Jaundice	Leaf	Decoction	Drink	7	1	7	0.89	7	100	(Prasad et al., 2008)
Convolvulaceae				Cuts and Wounds	Leaf	Crushed	Rubbed	2	1	2	0.25	2	100	
<i>Jasminum sambac</i> (L.) Aiton	Beli (B)	AUS2553	0.60	Bone Fracture	Leaf	Crushed	Rubbed	2	2	5	0.63	3	60	(Sabharwal et al., 2011; Sengar et al., 2015)
Oleaceae				Arthritis	Leaf	Crushed	Rubbed	3						
				Diarrhoea	Leaf	Decoction	Drink	7	1	7	0.89	7	100	
<i>Jatropha curcas</i> L.	Dantigach (D), bherenda (B)	AUS2554	0.40	Eczema	Seed oil	Ointment	Rubbed	2	1	2	0.25	2	100	(Joshi and Joshi, 2000)
Euphorbiaceae				Cut and Wounds	Leaf	Crushed	Rubbed	6	1	6	0.76	6	100	
<i>Jatropha gossypifolia</i> L.	Lal bherenda (B)	AUS2555	0.40	Teeth Infection	Latex	Ointment	Rubbed	6	1	6	0.76	6	100	(Panda et al., 2009)
Euphorbiaceae				Snake Bite	Leaf	Crushed	Rubbed	2	1	2	0.25	2	100	
<i>Justicia gendarussa</i> Burmf.	Bishoma (C)	AUS2556	0.75	Waist Pain	Leaf	Crushed	Rubbed	7	2	15	1.90	8	53.3333	(Jothimanivannan et al., 2010; Paval et al., 2009)
Acanthaceae				Inflammation	Leaf	Crushed	Rubbed	8						
<i>Justicia adhatoda</i> L.	Basak (B)	AUS2557	0.90	Cough	Leaf	Decoction	Drink	4	2	18	2.29	14	77.7778	(Nath et al., 1992; Singh et al., 2000; Mahato and Chaudhary, 2005; Joshi and Joshi, 2000)
Acanthaceae				Fever	Leaf	Decoction	Drink	14						
<i>Lablab purpureus</i> (L.) Sweet.	Lalshim (B)	AUS2551	0.20	Eczema	Leaf	Crushed	Rubbed	4	1	4	0.50	4	100	(Islam et al., 2011)
Leguminosae				Teeth Infection	Root	Fresh	Chewed	9	1	9	1.14	9	100	(Vivek et al., 2013)
<i>Lagerstroemia speciosa</i> (L.) Pers.	Jarulata (B)	AUS2558	0.55	Cut and Wounds	Root	Crushed	Rubbed	2	1	2	0.25	2	100	
Lythraceae				Urinary Troubles	Leaf	Crushed	Drink	4	1	4	0.50	4	100	(Ali et al., 1996; Chaudhary et al., 2010)
<i>Lawsonia inermis</i> L.	Mehendi (B)	AUS2559	0.55	Catamenia	Leaf	Crushed	Pellet	3	1	3	0.38	3	100	
Lythraceae				Body Pain	Leaf	Decoction	Drink	4	1	4	0.50	4	100	
<i>Leucas aspera</i> (Willd.) Link	Dankalshi (B)	AUS2560	0.80	Piles	Leaf	Crushed	Rubbed	4	1	4	0.50	4	100	(Reddy et al., 1986; Srinivas et al., 2000)
Lamiaceae				Inflammation	Whole Plant	Crushed	Drink	7	2	12	1.52	7	58.3333	
				Dysentery	Leaf	Crushed	Drink	5						
<i>Litsea glutinosa</i> (Lour.) C.B. Rob.	Chengpislal (B) Musafaoma (R)	AUS2561	1.0	Gastric	Leaf	Crushed	Drink	5	2	14	1.78	9	64.2857	(Bhowmick et al., 2014; Chowdhury et al., 2008; Ghani, 2003; Bhuinya et al., 2010)
Lauraceae				Dysentery	Leaf	Crushed	Pellet	6	1	6	0.76	6	100	
				Semen Loss	Bark	Crushed	Drink	9						
<i>Litsea monopetala</i> (Roxb.) Pers. ^a	Kala deungra (B)	AUS2562	0.4	Diabetes	Seed	Decoction	Drink	6	2	8	1.01	6	75	

Table 1 (continued)

Plant name and family	Vernacular name	Voucher no.	RFC	Main disease	Parts used	Preparation	Application	Number of citations per preparation	No. of uses type	Ct	%Ct	Cm	%Cm	ISu	Other uses and mode of application	Relevant ethnobotanical/ pharmacological citation
Lauraceae				Fever	Leaf	Decoction	Drink	2								
<i>Litsea semecarpifolia</i> (Wall. Ex. Nees) Hook.f. ^a	Leichchoshiyang (C)	AUS2563	0.35	Diabetes	Seed	Crushed	Drink	2	1	2	0.25	2	100	0.25		
Lauraceae				Body Pain	Leaf	Crushed	Rubbed	5	1	5	0.63	5	100	0.63		
<i>Litsea salicifolia</i> (J. Roxb. Ex Nees) Hook.f. ^a	Shigonjalin odonmointto (C)	AUS2564	0.35	Arthritis	Leaf	Decoction	Drink	5	2	7	0.89	5	71.4286	0.63		
Lauraceae				Body Pain	Stem	Decoction	Drink	2								
<i>Melastoma malabathricum</i> L. ^a	Lutki (B)	AUS2565	0.35	Bone Fracture	Leaf	Crushed	Rubbed	2	1	2	0.25	2	100			
Melastomataceae				Asthma	Leaf	Crushed	Pellet	5	1	5	0.63	5	100	0.63		
<i>Mangifera indica</i> L.	Aam (B, H)	AUS2566	0.55	Ulcer	Bark	Crushed	Drink	3	2	5	0.63	3	60	0.38		(Carvalho et al., 2007; Severi et al., 2009)
Anacardiaceae				Gastric Diabetes	Leaf	Crushed	Drink	2								
<i>Mentha arvensis</i> L.	Pudina (B)	AUS2567	0.40	Indigestion	Leaf	Decoction	Drink	6	1	6	0.76	6	100	0.76		
Lamiaceae				Cattle Fever	Whole Plant	Fresh	Chewed	5	1	5	0.63	5	100	0.63		(Londonkar and Poddar, 2009)
<i>Micromelum integerrimum</i> (Buch.-Ham. ex DC.) Wight & Arn. Ex M. Roem. ^a	Banjamra (B)	AUS2568	0.50	Diabetes	Leaf	Decoction	Drink	3	1	3	0.38	3	100	0.38		
Rutaceae				Dysentery	Leaf	Crushed	Drink	5	2	8	1.01	5	62.5	0.63		
<i>Mikania cordata</i> (Burm.f.) B.L.Rob. ^a	Noreshbodug, pikash (D)	AUS2569	0.60	Arthritis	Leaf	Crushed	Rubbed	2	1	2	0.25	2	100	0.25		
Asteraceae				Diabetes	Leaf	Crushed	Drink	5	1	5	0.63	5	100	0.63		
<i>Mimosa pudica</i> L.	Lajjaboti (B, H, M)	AUS2570	0.70	Cut and Wounds	Leaf	Crushed	Rubbed	7	1	7	0.89	7	100	0.89		
Mimosaceae				Asthma	Leaf	Crushed	Rubbed	9	1	9	1.14	9	100	1.14		(Girish et al., 2004; Mahanta and Mukherjee, 2001; Yang et al., 2011)
				Stone	Whole Plant	Crushed	Drink	4	1	4	0.50	4	100	0.50		
				Dog/Snake Bite	Root	Crushed	Pellet	1	1	1	0.12	1	100	0.12		
<i>Momordica charantia</i> L.	Titkarala (B)	AUS2571	0.60	Indigestion	Leaf	Decoction	Drink	7	1	7	0.89	7	100	0.89		(Akhtar et al., 1981; Raman and Lau, 1996; Sarkar et al., 1996; Welihinda et al., 1986)
Cucurbitaceae				Diabetes	Leaf	Crushed	Drink	3	2	5	0.63	3	60	0.38		
<i>Morinda angustifolia</i> Roxb. ^a	Borshoilli (C)	AUS2572	0.45	Diabetes	Fruit	Crushed	Drink	2								
Rubiaceae				Dysentery	Stem	Decoction	Drink	3	1	3	0.38	3	100	0.38		
<i>Moringa oleifera</i> Lam.	Sajna (B)	AUS2573	0.45	Allergy	Leaf	Crushed	Rubbed	6	1	6	0.76	6	100	0.76		
Moringaceae				Jaundice	Bark	Crushed	Pellet	5	1	5	0.63	5	100			(Fakurazi et al., 2008; Pari and Kumar, 2002)
<i>Murraya koenigii</i> (L.) Spreng.	Baupata (M)	AUS2574	0.35	High B/P	Bark	Crushed	Drink	4	1	4	0.50	4	100	0.50		
Rutaceae				Stomach Ache	Leaf	Decoction	Drink	6	1	6	0.76	6	100	0.76		(Chakraborty et al., 1965; Shah et al., 2008)
<i>Myxopyrum smilacifolium</i> (Wall.) Blume ^a	Arcid (B)	AUS2575	0.45	Piles	Leaf	Crushed	Drink	1	1	1	0.12	1	100	0.12		
				Fever	Leaf	Decoction	Drink	3	1	3	0.38	3	100	0.38		




Fig. 2. Percentage distribution of informants category (a); informants gender (b); informants age (c); educational background of informants (d) and informants experience (e).

Table 2
Informant consensus factor for commonly used medicinal plants.

Disease Category	F _{IC}
Respiratory system disorder	0.85
Digestive system disorder	0.84
Endocrine system disorder	0.82
Infection	0.82
Poisoning	0.81
Injuries	0.80
Muscular skeletal disorder	0.78
Genitourinary system disorder	0.78
Mental disorder	0.76
Nervous System Disorders	0.75
Skin/Subcutaneous Cellular Tissue Disorders	0.75
Pain	0.69
Circulatory system disorder	0.69
Veterinary Diseases	0.50
Sensory System Disorders	0.50

et al., 2011). However, this argument might always not be true when traditional healer identified different role of plant parts with a particular disease, as observed in this field study. For instance, in *Alstonia scholaris*, use of bark for diarrhoea have the higher ISu as compared to use of leaves for bone fracture, while in *Jatropha gossypifolia* L., use of latex for teeth infection have more preference over use of leave for snake bites. Knowledge about difference in pharmacological action and concentration of bioactive compounds

between different plant parts might be the reason for selectivity of plant parts by the traditional healers (Elisabetsky and Etkin, 2005). Selectivity of the plant parts depend on the nature and type of the disease. When different plant parts have the same effectiveness for a disease, leaves usually became the initial choice. Apart from convenience, healers usually choose leaves as major medicinal source as uprooting usually lead to death of the plant and they had to maintain their continuous supply (Poffenberger et al., 1992; Abebe and Ayehu, 1993; Martínez et al., 2000; Zheng and Xing, 2009; Rehecho et al., 2011). Consciousness for conservation of medicinal plants are there among the traditional healers are there, that can be observed as cultivation of formerly wild-plants in their home gardens (Etkin and Ticktin, 2005). During the field survey, the researchers observed cultivation of many such plants and even collected some plants not available in the wild from their home-gardens.

Preparation of medicine from the raw plant materials was one of the most important processes in herbal therapies. Preference of raw materials over other types of preparation were also observed in other parts of the state (Majumdar and Datta, 2007; Majumdar et al., 2006) and in neighbouring communities of Hmar, Assamese etc., in the state of Assam, (Nath and Choudhury, 2010).

During our survey, some THPs took interest in concept of supernatural power and their role in disease treatment, similar to observation of earlier studies in the state (Majumdar and Datta, 2007). As our study mainly focussed on the use and efficacies of the medicinal plants we have not incorporated par se purely spiritual therapies in the analysis. However, maintenance of strict rules and regulations in collecting and preparation of traditional

recipes have relevancy in the efficacies of these therapies. Collection of the plant at a specific time, the order of mixing of different plants or adjuvant during the concoction, the time of day at which preparations are made or the time of taking the dose might be influenced by how traditional people are giving special treatment for harvesting the associated bioactive compounds (Elisabetsky and Etkin, 2005).

During the survey, it was observed the people in the district are using same vernacular name for *P. longifolia* and *O. indicum*. Fruits of these two plants are similar to each other. Their usage pattern in jaundice was similar but differs in dysentery. Though similar usage pattern with common vernacular name might represent replacement and substitution therapy (Khare, 2004), the healers also acknowledged difference in pharmacological properties by not including *P. longifolia* in the treatment of dysentery.

During the survey, informants without modern education were observed to retain more traditional knowledge about the medicinal plants as compared to those informants with modern education system. It could be attributed to negligence of literate people towards traditional remedy and more proclivities towards modern treatment (Majumdar et al., 2006). Lack of attention among younger generation is one of the reasons for loss of valuable heritage regarding medicinal plant usage in many parts of the world (Kadir et al., 2012). In the existing scenario, ongoing acculturation process remains the major impediment in finding reliable information about restorative properties of medicinal plants. Counselling of traditional health practitioners should be taken into consideration in order to smooth continuation and extension of traditional medical knowledge. At the same time, it is important to integrate and provide awareness about potential toxicological properties of the medicinal recipes for ensuring safe and effective therapy.

Out of the total 75 collected species 49 species have been pharmacologically tested, 7 species studied chemically for isolation of active principles and 19 plants (marked with asterisk in table -1) are yet to be studied either pharmacologically or chemically. Intensive biochemical screening of these plants may provide important leads for treating associated ailments.

Interestingly, it has been observed that none of the plants collected during the present work falls under IUCN threat category. Therefore, the use of enumerated species in herbal formulations creates no ecological problem.

5. Conclusion

The present study could highlight important medicinal plants with their specified disease. Medicinal plants such as *L. glutinosa*, *P. longifolia*, *C. roseus*, *J. adhatoda* etc. are observed to have priority ranking for further experimentation. Healers are observed to acknowledge the differential pharmacological properties of medicinal plants with regard to different diseases, which may be helpful in further bioprospective programmes. Documentation and nurturing of traditional knowledge through various means are required to inculcate the higher value to traditional knowledge among younger generation might help in conserving slowly eroding traditional knowledge and medicinal plants.

Acknowledgement

We are very grateful to the people of North Tripura district, Tripura for their kind cooperation during the survey. We are also indebted to DeLCON (DBT-Electronic Library Consortium), Govt. of India and Bioinformatics Centre of Assam University, Silchar for providing free access to the journals and articles.

References

- Abebe, D., Ayehu, A., 1993. Medicinal Plants and Enigmatic Health Practices of Northern Ethiopia.
- Akerele, O., 1984. WHO's traditional medicine programme: progress and perspectives. WHO Chron. 38, 76–81.
- Akhtar, M.S., Athar, M.A., Yaqub, M., 1981. Effect of *Momordica charantia* on blood glucose level of normal and alloxan-diabetic rabbits. Planta Med. 42, 205–212.
- Akinloye, O.A., Olaniyi, M.O., 2011. Hepatoprotective effect of *Cajanus cajan* on tissue defense system in D-galactosamine-induced hepatitis in rats. Turk. J. Biochem. 36, 237–241.
- Ali, B., Bashir, A., Tanira, M., 1996. Anti-inflammatory, antipyretic, and analgesic effects of *Lawsonia inermis* L. (henna) in rats. Pharmacology, 356–363.
- Andrade-Cetto, A., Heinrich, M., 2011. From the field into the lab: useful approaches to selecting species based on local knowledge. Front. Pharmacol. 2, 20.
- Angami, A., Gajurel, P., Rethy, P., Singh, B., Kalita, S., 2006. Status and potential of wild edible plants of Arunachal Pradesh. Indian J. Tradit. Knowl. 5, 541–550.
- Arul, V., Miyazaki, S., Dhananjayan, R., 2004. Mechanisms of the contractile effect of the alcoholic extract of *Aegle marmelos* Corr. on isolated guinea pig ileum and tracheal chain. Phytomedicine 11, 679–683.
- Arul, V., Miyazaki, S., Dhananjayan, R., 2005. Studies on the anti-inflammatory, antipyretic and analgesic properties of the leaves of *Aegle marmelos* Corr. J. Ethnopharmacol. 96, 159–163.
- Atta-ur-Rahman, A., Asif, M., Ghazala, M., Fatima, J., Alvi, K., 1985. Scholaricine, an alkaloid from *Alstonia inermis* L. (henna) in rats. Phytochemistry 24, 2771–2773.
- Babalola, O., Ojo, O., Oloyede, F., 2011. Hepatoprotective activity of aqueous extract of the leaves of *Hyptis suaveolens* (L.) Poit on acetaminophen induced hepatotoxicity in rabbits. Res. J. Chem. Sci. 1, 85–88.
- Ballabh, B., Chaurasia, O., Ahmed, Z., Singh, S.B., 2008. Traditional medicinal plants of cold desert Ladakh—used against kidney and urinary disorders. J. Ethnopharmacol. 118, 331–339.
- Banerjee, R.N., Basu, T.K., 1991. Hypoglycemic and antihyperglycemic effect of leaves of *Vinca rosea* Linn. Indian J. Physiol. Pharmacol. 35, 145–151.
- Bhandurge, P., Rajarajeshwari, N., Alagawadi, K., Agrawal, S., 2010. Antidiabetic and hyperlipaemic effects of *Citrus Maxima* Linn. fruits on alloxan-induced diabetic rats. Int. J. Drug Dev. Res. 2, 273–278.
- Bhowmick, R., Sarwar, M.S., Dewan, S.M.R., Das, A., Das, B., Uddin, M.M.N., Islam, M. S., Islam, M.S., 2014. In vivo analgesic, antipyretic, and anti-inflammatory potential in Swiss albino mice and in vitro thrombolytic activity of hydroalcoholic extract from *Litsea glutinosa* leaves. Biol. Res. 47, 56.
- Bhuiya, T., Singh, P., Mukherjee, S.K., 2010. Traditional uses of the Indian species of *Litsea* Lam. Phytotaxonomy 10, 45–46.
- Borah, P., Gogoi, P., Phukan, A., Mahanta, J., 2006. Traditional medicine in the treatment of gastrointestinal diseases in Upper Assam. Indian J. Tradit. Knowl. 5, 510–512.
- Brijesh, S., Daswani, P., Tetali, P., Antia, N., Birdi, T., 2009. Studies on the antidiarrhoeal activity of *Aegle marmelos* unripe fruit: validating its traditional usage. BMC Complement. Altern. Med. 9, 47.
- Bruni, A., Ballero, M., Poli, F., 1997. Quantitative ethnopharmacological study of the Campidano Valley and Urzulei district, Sardinia, Italy. J. Ethnopharmacol. 57, 97–124.
- Calixto, J.B., 2005. Twenty-five years of research on medicinal plants in Latin America: a personal view. J. Ethnopharmacol. 100, 131–134.
- Carvalho, A.C., Guedes, M.M., de Souza, A.L., Trevisan, M.T., Lima, A.F., Santos, F.A., Rao, V.S., 2007. Gastroprotective effect of mangiferin, a xanthonoid from *Mangifera indica*, against gastric injury induced by ethanol and indomethacin in rodents. Planta Med. 73, 1372–1376.
- Chakraborty, D., Barman, B., Bose, P., 1965. On the constitution of murrayanine, a carbazole derivative isolated from *Murraya koenigii* Spreng. Tetrahedron 21, 681–685.
- Chanda, S., Sarethy, I.P., De, B., Singh, K., 2013. *Paederia foetida* – a promising ethnomedicinal tribal plant of northeastern India. J. For. Res. 24, 801–808.
- Chatterjee, T., Chakraborty, A., Pathak, M., Sengupta, G., 1992. Effects of plant extract *Centella asiatica* (Linn.) on cold restraint stress ulcer in rats. Indian J. Exp. Biol. 30, 889–891.
- Chattopadhyay, R., Chattopadhyay, R., Nandy, A., Poddar, G., Maitra, S., 1987. Preliminary report on antihyperglycemic effect of a fraction of fresh leaves of *Azadirachta indica* (Beng. Neem). Bull. Calcutta Sch. Trop. Med. 35, 29–33.
- Chaudhary, G., Goyal, S., Poonia, P., 2010. *Lawsonia inermis* Linnaeus: a phytopharmacological review. Int. J. Pharm. Sci. Drug Res. 2, 91–98.
- Chavan, M.J., Wakte, P.S., Shinde, D.B., 2012. Analgesic and anti-inflammatory activities of the sesquiterpene fraction from *Annona reticulata* L. bark. Nat. Prod. Res. 26, 1515–1518.
- Cheng, C., Koo, M., 2000. Effects of *Centella asiatica* on ethanol induced gastric mucosal lesions in rats. Life Sci. 67, 2647–2653.
- Chowdhury, J.U., Bhuiyan, M.N.I., Nandi, N.C., 2008. Aromatic plants of Bangladesh: essential oils of leaves and fruits of *Litsea glutinosa* (Lour.) C.B. Rob. Bangladesh J. Bot. 37, 81–83.
- Dalziel, J.M., 1937. The Useful Plants of West Tropical Africa. Crown Agents for the Colonies, London, pp. 28–29.
- Das, H.B., Majumdar, K., Datta, B., Ray, D., 2009. Ethnobotanical uses of some plants by Tripuri and Reang tribes of Tripura. Nat. Prod. Radiance 8, 172–180.
- Das, S., Choudhury, M.D., 2010. Plants used against gastro-intestinal disorders and as anti-hemorrhagic by three tribes of North Tripura district, Tripura, India: a Report. Ethnobot. Leaflet. 14, 467–478.

- Das, S., Dash, S., Padhy, S., 2003. Ethno-medicinal information from Orissa State, India – a review. *J. Hum. Ecol.* 14, 165–227.
- Deb, D., Sarkar, A., Barma, B.D., Datta, B., Majumdar, K., 2013. Wild edible plants and their utilization in traditional recipes of Tripura, Northeast India. *Adv. Biol. Res.* 7, 203–211.
- Deb, D.B., 1968. Medicinal plants of Tripura state. *Indian Forester* 94, 753–765.
- Deb, D.B., 1975a. Economic plants of Tripura state timber yielding plants principal uses. *Indian For.* 101, 282–288.
- Deb, D.B., 1975b. Economic plants of Tripura state-edible plants: foodgrains, Vegetables and Fruits. *Indian For.* 101, 399–406.
- Deb, D.B., 1975c. Economic plants of Tripura state-timber yielding plants, miscellaneous uses. *Indian For.* 101, 328–336.
- Deb, D.B., 1976. Economic plants of Tripura state-IV. Edible plants: fats, oils, spices, seasoning herbs, beverages, narcotics and sugar. *Indian For.* 102, 273–277.
- Deb, D.B., 1978a. Economic plants of Tripura state V. Pasture and Fodder. *Indian For.* 104, 275–281.
- Deb, D.B., 1978b. Economic plants of Tripura state – VI Plants Yielding Fibre. *Indian For.* 104, 407–410.
- Deb, D.B., 1981a. The Flora of Tripura State, vol. I. Today and Tomorrow's Printers and Publishers, New Delhi.
- Deb, D.B., 1981b. Economic Plants of Tripura State IX. Plants yielding tannins. *Indian For.* 107, 426–431.
- Deb, D.B., 1983. The Flora of Tripura State, vol. II, Today and Tomorrow's Printers and Publishers, New Delhi.
- Deb, S., Arunachalam, A., Das, A., 2009. Indigenous knowledge of Nyishi tribes on traditional agroforestry systems. *Indian J. Tradit. Knowl.* 8, 41–46.
- Devbarma, M., 1976. 'Chauk' – a drink for tribals in Tripura. *Folklore* 17, 347–351.
- Elisabetsky, E., Etkin, N.L., 2005. *Ethnopharmacology: An Overview*, in *Ethnopharmacology*. EOLSS Publishers, Oxford.
- Etkin, N.L., Tickint, T., 2005. Integrating Ethnographic and Ecological Perspectives for Ethnopharmacology Field Research. EOLSS Publishers, Oxford.
- Fakurazi, S., Hairuszah, I., Nanthini, U., 2008. *Moringa oleifera* Lam. prevents acetaminophen induced liver injury through restoration of glutathione level. *Food Chem. Toxicol.* 46, 2611–2615.
- Friedman, J., Yaniv, Z., Dafni, A., Palewitch, D., 1986. A preliminary classification of the healing potential of medicinal plants, based on a rational analysis of an ethnopharmacological field survey among Bedouins in the Negev Desert, Israel. *J. Ethnopharmacol.* 16, 275–287.
- Gaire, B.P., Subedi, L., 2013. A review on the pharmacological and toxicological aspects of *Datura stramonium* L. *J. Integr. Med.* 11, 73–79.
- Ghaffari, H., Ghassam, B.J., Prakash, H., 2012. Hepatoprotective and cytoprotective properties of *Hyptis suaveolens* against oxidative stress-induced damage by CCl₄ and H₂O₂. *Asian Pac. J. Trop. Med.* 5, 868–874.
- Ghani, A., 2003. Medicinal Plants of Bangladesh: Chemical Constituents and Uses (Revised and Enlarged). Asiatic Society of Bangladesh, Old Nimtali, Dhaka, pp. 196–197.
- Ghosh, R.K., Gupta, I., 1980. Effect of *Vinca rosea* and *Ficus racemosus* on hyperglycemia in rats. *Indian J. Anim. Health* 19, 145–148.
- Giday, M., Asfaw, Z., Woldu, Z., 2009. Medicinal plants of the Meinit ethnic group of Ethiopia: an ethnobotanical study. *J. Ethnopharmacol.* 124, 513–521.
- Girish, K., Mohanakumari, H.P., Nagaraju, S., Vishwanath, B., Kemparaju, K., 2004. Hyaluronidase and protease activities from Indian snake venoms: neutralization by *Mimosa pudica* root extract. *Fitoterapia* 75, 378–380.
- Gogtay, N., Bhatt, H., Dalvi, S., Kshirsagar, N., 2002. The use and safety of non-allopathic Indian medicines. *Drug Saf.* 25, 1005–1019.
- Guo, J.S., Cheng, C.L., Koo, M.V., 2004. Inhibitory effects of *Centella asiatica* water extract and asiaticoside on inducible nitric oxide synthase during gastric ulcer healing in rats. *Planta Med.* 70, 1150–1154.
- Gupta, V., 2006. Plant used in folkloric medicine by Bangnis of East Kameng, Arunachal Pradesh. *Nat. Prod. Radiance* 5, 52–59.
- Heinrich, M., Ankli, A., Frei, B., Weimann, C., Sticher, O., 1998. Medicinal plants in Mexico: healers' consensus and cultural importance. *Soc. Sci. Med.* 47, 1859–1871.
- Hooker, J.D., 1872–1897. The Flora of British India. Vol. 1–7. L. Reeve & Co., London.
- Hossain, M.M., Ahmed, S.K., Dewan, S.M.R., Hassan, M.M., Istiaq, A., Islam, M.S., Moghal, M.M.R., 2014. In vivo antipyretic, antiemetic, in vitro membrane stabilization, antimicrobial, and cytotoxic activities of different extracts from *Spilanthes paniculata* leaves. *Biol. Res.* 47, 45.
- Hossain, M.S., Hanif, A., Khan, M., Bari, S., Jahan, R., Rahmatullah, M., 2009. Ethnobotanical survey of the Tripura tribe of Bangladesh. *Am. Eurasian J. Sustain. Agric.* 253–261.
- Hossain, S., Agarwala, B., Sarwar, S., Karim, M., Jahan, R., Rahmatullah, M., 2010. Traditional use of medicinal plants in Bangladesh to treat urinary tract infections and sexually transmitted diseases. *Ethnobot. Res. Appl.* 8, 61–74.
- Islam, F., Jahan, F.I., Seraj, S., Malek, I., Sadat, A., Bhuiyan, M.S.A., Swarna, A., Sanam, S., Rahmatullah, M., 2011. Variations in diseases and medicinal plant selection among folk medicinal practitioners: a case study in Jessore district, Bangladesh. *Am. Eurasian J. Sustain. Agric.* 5, 282–291.
- Jahan, F.I., Hasan, M.R.U., Jahan, R., Seraj, S., Chowdhury, A.R., Islam, M.T., Khatun, Z., Rahmatullah, M., 2011. A comparison of medicinal plant usage by folk medicinal practitioners of two adjoining villages in Lalmonirhat district, Bangladesh. *Am. Eurasian J. Sustain. Agric.* 5, 46–66.
- Joshi, A.R., Joshi, K., 2000. Indigenous knowledge and uses of medicinal plants by local communities of the Kali Gandaki Watershed Area, Nepal. *J. Ethnopharmacol.* 73, 175–183.
- Jothimanivannan, C., Kumar, R., Subramanian, N., 2010. Anti-inflammatory and analgesic activities of ethanol extract of aerial parts of *Justicia gendarussa* Burm. *Int. J. Pharmacol.* 6, 278–283.
- Kadir, M.F., Bin Sayeed, M.S., Mia, M., 2012. Ethnopharmacological survey of medicinal plants used by indigenous and tribal people in Rangamati, Bangladesh. *J. Ethnopharmacol.* 144, 627–637.
- Kamboj, A., Saluja, A.K., 2009. *Bryophyllum pinnatum* (Lam.) Kurz.: phytochemical and pharmacological profile: a review. *Pharmacogn. Rev.* 3, 364.
- Kanjilal, U.N., Kanjilal, P.C., Das, A., 1936. Flora of Assam, Vol. 2. Govt. of Assam, Shillong.
- Kanjilal, U.N., Kanjilal, P.C., Das, A., De, R.N., 1938. Flora of Assam, Vol. 3. Govt. of Assam, Shillong.
- Kanjilal, U.N., Kanjilal, P.C., Das, A., Purkayastha, C., 1934. Flora of Assam, Vol. 1. Govt. of Assam, Shillong.
- Kanjilal, U.N., Kanjilal, P.C., De, R.N., Das, A., 1940. Flora of Assam, Vol. 4. Govt. of Assam, Shillong.
- Kar, A., Choudhary, B., Bandyopadhyay, N., 2003. Comparative evaluation of hypoglycaemic activity of some Indian medicinal plants in alloxan diabetic rats. *J. Ethnopharmacol.* 84, 105–108.
- Khare, C.P., 2004. Indian Herbal Remedies: Rational Western Therapy, Ayurvedic, and Other Traditional Usage, Botany. Springer, New York, USA.
- Khare, C.P., 2007. Indian Medicinal Plants: An Illustrated Dictionary. Springer Science & Business Media, New York, USA.
- Khosla, P., Bhanwra, S., Singh, J., Seth, S., Srivastava, R., 2000. A study of hypoglycaemic effects of *Azadirachta indica* (Neem) in normal and alloxan diabetic rabbits. *Indian J. Physiol. Pharmacol.* 44, 69–74.
- Kim, M.-S., Kim, J.-K., Kim, H.-J., Moon, S.-R., Shin, B.-C., Park, K.-W., Yang, H.-O., Kim, S.-M., Park, R., 2003. *Hibiscus* extract inhibits the lipid droplet accumulation and adipogenic transcription factors expression of 3T3-L1 preadipocytes. *J. Altern. Complement. Med.* 9, 499–504.
- Kirtikar, K.R., Basu, B.D., 1933. 2nd ed. *Vinca rosea*. In *Indian Medicinal Plants*, Vol. III. Lalit Mohan Basu Publications, Allahabad, India.
- Kong, Y., Fu, Y.-J., Zu, Y.-G., Liu, W., Wang, W., Hua, X., Yang, M., 2009. Ethanol modified supercritical fluid extraction and antioxidant activity of cajanin stilbene acid and pinostrobin from pigeonpea [*Cajanus cajan* (L.) Millsp.] leaves. *Food Chem.* 117, 152–159.
- Kshirsagar, R., Upadhyay, S., 2009. Free radical scavenging activity screening of medicinal plants from Tripura, Northeast India. *Nat. Prod. Radiance* 8, 117–122.
- Kumar, M., Prasad, S.K., Hemalatha, S., 2014. A current update on the phytopharmacological aspects of *Houttuynia cordata* Thunb. *Pharmacogn. Rev.* 8, 22.
- Kundusen, S., Gupta, M., Mazumder, U.K., Haldar, P.K., Saha, P., Bhattacharya, S., Kar, B., Bala, A., 2011. Antihyperglycemic effect and antioxidant property of *Citrus maxima* leaf in streptozotocin-induced diabetic rats. *Diabetol. Croat.* 40, 113–120.
- Leong, C.N.A., Tako, M., Hanashiro, I., Tamaki, H., 2008. Antioxidant flavonoid glycosides from the leaves of *Ficus pumila* L. *Food Chem.* 109, 415–420.
- Liao, C.R., Kao, C.P., Peng, W.H., Chang, Y.S., Lai, S.C., Ho, Y.L., 2012. Analgesic and anti-inflammatory activities of methanol extract of *Ficus pumila* L. in mice. *Evid. Based Complement. Altern. Med.*, 2012.
- Londonkar, R.L., Poddar, P.V., 2009. Studies on activity of various extracts of *Mentha arvensis* Linn against drug induced gastric ulcer in mammals. *World J. Gastrointest. Oncol.* 1, 82.
- Mahanta, M., Mukherjee, A.K., 2001. Neutralisation of lethality, myotoxicity and toxic enzymes of *Naja kaouthia* venom by *Mimosa pudica* root extracts. *J. Ethnopharmacol.* 75, 55–60.
- Mahato, R., Chaudhary, R., 2005. Ethnomedicinal study and antibacterial activities of selected plants of Palpa district, Nepal. *Sci. World* 3, 26–31.
- Majumdar, K., Datta, B.K., 2007. A study on ethnomedicinal usage of plants among the folkloric herbalists and Tripuri medical practitioners: Part-II. *Nat. Prod. Radiance* 6, 66–73.
- Majumdar, K., Saha, R., Datta, B., Bhakta, T., 2006. Medicinal plants prescribed by different tribal and non-tribal medicine men of Tripura state. *Indian J. Tradit. Knowl.* 5, 559–562.
- Mao, A., Hynniewta, T., Sanjappa, M., 2009. Plant wealth of Northeast India with reference to ethnobotany. *Indian J. Tradit. Knowl.* 8, 96–103.
- Martínez, A., Bernal, Y., Cáceres, A., 2000. Fundamentos de agroecología de cultivo de plantas medicinales Iberoamericanas. Santafé de Bogotá. Convenio Andrés Bello/Ciencia y Tecnología para el Desarrollo, Colombia.
- Mia, M.M., Rahman, M.S., Begum, K., Begum, B., Rashid, M.A., 2007. Phytochemical and biological studies of *Averrhoa carambola*. *Dhaka Univ. J. Pharm. Sci.* 6, 125–128.
- Ming, L.C., 1999. *Ageratum conyzoides*: A tropical source of medicinal and agricultural products. In: Janick, J. (Ed.), Perspectives on New crops and New Uses. ASHS Press, Alexandria, VA, pp. 469–473.
- Misra, M.K., Mohanty, M.K., Das, P.K., 1993. Studies on the method-ethnobotany of *Calotropis gigantea* and *C. procera*. *Anc. Sci. Life* 13, 40.
- Morton, J., 1976. The pigeon pea (*Cajanus cajan* Millsp.) a high-protein, tropical bush legume. *Hortic. Sci.* 2, 11–19.
- Mukti, M., Ahmed, A., Chowdhury, S., Khatun, Z., Bhuiyan, P., Debnath, K., Rahmatullah, M., 2012. Medicinal plant formulations of Kavirajes in several areas of Faridpur and Rajbari districts, Bangladesh. *Am. Eurasian J. Sustain. Agric.* 6, 234–247.
- Muthu, C., Ayyanar, M., Raja, N., Ignacimuthu, S., 2006. Medicinal plants used by traditional healers in Kancheepuram District of Tamil Nadu, India. *J. Ethnobiol. Ethnomed.* 2, 43.

- Nadembega, P., Boussim, J.I., Nikiema, J.P., Poli, F., Antognoni, F., 2011. Medicinal plants in Baskoure, Kouritenga Province, Burkina Faso: an ethnobotanical study. *J. Ethnopharmacol.* 133, pp. 378–395.
- Nadkarni, K.M., Basu, B.D., 1996. *Alocasia indica*, Indian Materia Medica, 1st ed. Popular prakashan Ltd., Mumbai, India, pp. 72–73.
- Naim, M., Amjad, F.M., Sultana, S., Islam, S.N., Hossain, M.A., Begum, R., Rashid, M.A., Amran, M.S., 2012. Comparative study of antidiabetic activity of hexane-extract of lemon peel (*Limon citrus*) and glimepiride in alloxan-induced diabetic rats. *Bangladesh Pharm. J.* 15, 131–134.
- Nammi, S., Boini, M.K., Lodagala, S.D., Behara, R.B.S., 2003. The juice of fresh leaves of *Catharanthus roseus* Linn. reduces blood glucose in normal and alloxan diabetic rabbits. *BMC Complement. Altern. Med.* 3, 4.
- Nath, D., Sethi, N., Singh, R., Jain, A., 1992. Commonly used Indian abortifacient plants with special reference to their teratologic effects in rats. *J. Ethnopharmacol.* 36, 147–154.
- Nath, M., Choudhury, M.D., 2010. Ethno-medico-botanical aspects of Hmar tribe of Cachar district, Assam (Part-I). *Indian J. Tradit. Knowl.* 9, 760–764.
- Ningthoujam, S.S., Talukdar, A.D., Potsangbam, K.S., Choudhury, M.D., 2013. Traditional uses of herbal vapour therapy in Manipur, North East India: an ethnobotanical survey. *J. Ethnopharmacol.* 147, 136–147.
- Panda, B., Gaur, K., Kori, M., Tyagi, L., Nema, R., Sharma, C., Jain, A., 2009. Anti-inflammatory and analgesic activity of *Jatropha gossypifolia* in experimental animal models. *Glob. J. Pharmacol.* 3, 01–05.
- Pari, L., Kumar, N.A., 2002. Hepatoprotective activity of *Moringa oleifera* on antitubercular drug-induced liver damage in rats. *J. Med. Food* 5, 171–177.
- Patel, S., Sharma, V., Chauhan, N.S., Dixit, V.K., 2012. An updated review on the parasitic herb of *Cuscuta reflexa* Roxb. *J. Chin. Integr. Med.* 10, 249–255.
- Pattanayak, S.P., Mazumder, P.M., 2010. Phytochemical screening and safety evaluation of hydroalcoholic extract of *Dendrophthoe falcata* Ettingsh: summary of acute and subacute toxicological data. *Der Pharm. Lett.* 2, 127–138.
- Paval, J., Kaitheri, S.K., Potu, B.K., Govindan, S., Kumar, R.S., Narayanan, S.N., Moorkoth, S., 2009. Anti-arthritis potential of the plant *Justicia gendarussa* Burm.f. *Clinics* 64, 357–362.
- Poffenberger, M., McGean, B., Khare, A., Campbell, J., 1992. Field Methods Manual Volume II. Community Forest Economy and Use Patterns: Participatory Rural Appraisal (PRA) Methods in South Gujarat, India. Society for Promotion of Wetlands Development, New Delhi.
- Pokharen, N., Dahal, S., Anuradha, M., 2011. Phytochemical and antimicrobial studies of leaf extract of *Euphorbia nerifolia*. *J. Med. Plants Res.* 5, 5785–5788.
- Prasad, K.N., Shivamurthy, G., Aradhya, S., 2008. *Ipomoea aquatica*, an underutilized green leafy vegetable: a review. *Int. J. Bot.* 4, 123–129.
- Qureshi, R., Bhatti, G.R., 2008. Ethnobotany of plants used by the Thari people of Nara Desert, Pakistan. *Fitoterapia* 79, 468–473.
- Rahman, M.A., Solaiman, M., Haque, M.E., Das, A., 2011. Analgesic and anti-inflammatory activities of *Alocasia indica* (Roxb.) Schott. *Orient. Pharm. Exp. Med.* 11, 143–146.
- Rahmatullah, M., Ferdausi, D., Mollik, A., Jahan, R., Chowdhury, M.H., Haque, W.M., 2010a. A survey of medicinal plants used by Kavirajes of Chalna area, Khulna district, Bangladesh. *Afr. J. Tradit. Complement. Altern. Med.* 7, 91–97.
- Rahmatullah, M., Ferdausi, D., Mollik, M.A.H., Azam, M.N.K., Rahman, M., Jahan, R., 2009. Ethnomedicinal survey of Bheramara area in Kushtia district, Bangladesh. *Am. Eurasian J. Sustain. Agric.* 3, 534–541.
- Rahmatullah, M., Khatun, M.A., Morshed, N., Neogi, P.K., Khan, S.U.A., Hossain, M.S., Mahal, M.J., Jahan, R., 2010b. A randomized survey of medicinal plants used by folk medicinal healers of Sylhet Division, Bangladesh. *Adv. Nat. Appl. Sci.* 4, 52–62.
- Rai, L., Prasad, P., Sharma, E., 2000. Conservation threats to some important medicinal plants of the Sikkim Himalaya. *Biol. Conserv.* 93, 27–33.
- Rajendran, S., Agarwal, S., Sundaresan, V., 2004. Lesser known ethnomedicinal plants of the Ayyarkarkoil forest province of Southwestern Ghats, Tamil Nadu, India – Part I. *J. Herbs Spices Med. Plants* 10, 103–112.
- Raman, A., Lau, C., 1996. Anti-diabetic properties and phytochemistry of *Momordica charantia* L. (Cucurbitaceae). *Phytomedicine* 2, 349–362.
- Rasineni, K., Bellamkonda, R., Singareddy, S.R., Desireddy, S., 2010. Antihyperglycemic activity of *Catharanthus roseus* leaf powder in streptozotocin-induced diabetic rats. *Pharmacogn. Res.* 2, 195.
- Reddy, M.K., Viswanathan, S., Sambantham, P.T., Ramachandran, S., Kameswaran, L., 1986. Effect of *Leucas aspera* on experimental inflammation and mast cell degranulation. *Anc. Sci. Life* 5, 168.
- Rehecho, S., Uriarte-Pueyo, I., Calvo, J., Vivas, L.A., Calvo, M.I., 2011. Ethnopharmacological survey of medicinal plants in Nor-Yauyos, a part of the Landscape Reserve Nor-Yauyos-Cochas, Peru. *J. Ethnopharmacol.* 133, 75–85.
- Ripunjy, S., 2013. Indigenous knowledge on the utilisation of medicinal plants by the Sonowal Kachari tribe of Dibrugarh district in Assam, North-East India. *Int. Res. J. Biol. Sci.* 2, 44–50.
- Ross, I.A., 1999. *Hibiscus rosa sinensis*. In: Ross, I.A. (Ed.), *Medicinal Plants of the World. Chemical Constituents, Traditional and Modern Medicinal Uses*. Humana Press, Totowa, New Jersey, pp. 155–163.
- Roy, S., Hazra, B., Mandal, N., Chaudhuri, T.K., 2013. Assessment of the antioxidant and free radical scavenging activities of methanolic extract of *Diplazium esculentum*. *Int. J. Food Prop.* 16, 1351–1370.
- Sabharwal, S., Vats, M., Sardana, S., Aggarwal, S., 2011. Pharmacognostical, physico and phytochemical evaluation of the leaves of *Jasminum sambac* Linn. (Oleaceae). *Int. J. Pharm. Pharm. Sci.* 3, 237–241.
- Sairam, K., Rao, C.V., Goel, R., 2001. Effect of *Centella asiatica* Linn on physical and chemical factors induced gastric ulceration and secretion in rats. *Indian J. Exp. Biol.* 39, 137–142.
- Sambandan, K., Dhatchanamoorthy, N., 2012. Studies on the phytodiversity of a sacred grove and its traditional uses in Karaikal district, UT Puducherry. *J. Phytol.* 4, 16–21.
- Sannigrahi, S., Mazumder, U.K., Pal, D.K., Parida, S., Jain, S., 2010. Antioxidant potential of crude extract and different fractions of *Enhydra fluctuans* Lour. *Iranian J. Pharm. Res.* 9, 75.
- Sannigrahi, S., Mazumder, U.K., Pal, D., Mishra, M., Maity, S., 2011. Flavonoids of *Enhydra fluctuans* exhibits analgesic and anti-inflammatory activity in different animal models. *Pak. J. Pharm. Sci.* 24, 369–375.
- Sarkar, S., Pranava, M., MARITA, A.R., 1996. Demonstration of the hypoglycemic action of *Momordica charantia* in a validated animal model of diabetes. *Pharmacol. Res.* 33, 1–4.
- Sen, S., Chakraborty, R., De, B., Devanna, N., 2011. An ethnobotanical survey of medicinal plants used by ethnic people in West and South district of Tripura, India. *J. For. Res.* 22, 417–426.
- Sengar, N., Joshi, A., Prasad, S.K., Hemalatha, S., 2015. Anti-inflammatory, analgesic and anti-pyretic activities of standardized root extract of *Jasminum sambac*. *J. Ethnopharmacol.* 160, 140–148.
- Severi, J.A., Lima, Z.P., Kushima, H., Brito, Monteiro Souza, Campaner dos Santos, A. R., Vilegas, L., Hiruma-Lima, C.A., W., 2009. Polyphenols with antiulcerogenic action from aqueous decoction of mango leaves (*Mangifera indica* L.). *Molecules* 14, 1098–1110.
- Shah, A.J., Gowani, S.A., Zuberi, A.J., Ghayur, M.N., Gilani, A.H., 2010. Antidiarrhoeal and spasmolytic activities of the methanolic crude extract of *Alstonia scholaris* L. are mediated through calcium channel blockade. *Phytother. Res.* 24, 28–32.
- Shah, A.S., Wakade, A.S., Juvekar, A.R., 2008. Immunomodulatory activity of methanolic extract of *Murraya koenigii* (L.) Spreng. leaves. *Indian J. Exp. Biol.* 46, 505.
- Sharma, D.K., 2010. Bio-Diversity Resource Book of Tripura. International Book Distributor, Dehradun.
- Sharma, M., Sharma, C.L., Debbarma, J., 2013. Ethnobotanical uses of some tree species in Khowai district of Tripura, NE India. *Life Sci. Leaf.* 4, 60–80.
- Sharma, M., Sharma, C.L., Debbarma, J., 2014. Ethnobotanical studies of some plants used by tripuri of Tripura, NE India with special reference to Magico Religious Beliefs. *Int. J. Plant Anim. Environ. Sci.* 4, 518–528.
- Shibib, B.A., Khan, L.A., Rahman, R., 1993. Hypoglycaemic activity of *Coccinia indica* and *Momordica charantia* in diabetic rats: depression of the hepatic gluconeogenic enzymes glucose-6-phosphatase and fructose-1, 6-bisphosphatase and elevation of both liver and red-cell shunt enzyme glucose-6-phosphate dehydrogenase. *Biochem. J.* 292, 267–270.
- Shil, S., Choudhury, M.D., 2009. Ethnomedicinal importance of Pteridophytes used by Reang tribe of Tripura, North East India. *Ethnobot. Leaf.* 13, 634–643.
- Shil, S., Choudhury, M.D., Das, S., 2014. Indigenous knowledge of medicinal plants used by the Reang tribe of Tripura state of India. *J. Ethnopharmacol.* 152, 135–141.
- Shirwaikar, A., Khan, S., Malini, S., 2003. Antiosteoporotic effect of ethanol extract of *Cissus quadrangularis* Linn. on ovariectomized rat. *J. Ethnopharmacol.* 89, 245–250.
- Shoba, F.G., Thomas, M., 2001. Study of antidiarrhoeal activity of four medicinal plants in castor-oil induced diarrhoea. *J. Ethnopharmacol.* 76, 73–76.
- Shukla, A.N., Srivastava, S., Rawat, A., 2013. A survey of traditional medicinal plants of Uttar Pradesh (India) – used in treatment of infectious diseases. *Nat. Sci.* 11, 24–36.
- Singh, H., Hynniewta, T., Bora, P., 1997. Ethno-medico-botanical studies in Tripura, India. *Ethnobot. Res. Appl.* 9, 56–58.
- Singh, R.P., Padmavathi, B., Rao, A.R., 2000. Modulatory influence of *Adhatoda vesica* (justicia *adhatoda*) leaf extract on the enzymes of xenobiotic metabolism, antioxidant status and lipid peroxidation in mice. *Mol. Cell. Biochem.* 213, 99–109.
- Singh, S.N., Vats, P., Suri, S., Shyam, R., Kumria, M., Ranganathan, S., Sridharan, K., 2001. Effect of an antidiabetic extract of *Catharanthus roseus* on enzymic activities in streptozotocin induced diabetic rats. *J. Ethnopharmacol.* 76, 269–277.
- Siwakoti, M., Siwakoti, S., 2000. Ethnomedicinal uses of plants among the Satar tribe of Nepal. *J. Econ. Taxon. Bot.* 24, 323–333.
- Srinivas, K., Rao, M., Rao, S., 2000. Anti-inflammatory activity of *Heliotropium indicum* Linn. and *Leucas aspera* spreng. in albino rats. *Indian J. Pharmacol.* 32, 37–38.
- Srivastava, R., 2010a. Traditional knowledge of Nyishi (Daffla) tribe of Arunachal Pradesh. *Indian J. Tradit. Knowl.* 9, 26–37.
- Srivastava, R., Singh, R.K., Mukherjee, T., 2010b. Indigenous biodiversity of Apatani plateau: learning on biocultural knowledge of Apatani tribe of Arunachal Pradesh for sustainable livelihoods. *Indian J. Tradit. Knowl.* 9, 432–442.
- Surjushe, A., Vasani, R., Saple, D., 2008. Aloe vera: a short review. *Indian J. Dermatol.* 53, 163.
- Tardío, J., Pardo-de-Santayana, M., 2008. Cultural importance indices: a comparative analysis based on the useful wild plants of Southern Cantabria (Northern Spain) 1. *Econ. Bot.* 62, 24–39.
- Telefo, P., Lienou, L., Yemele, M., Lemfack, M., Mouokeu, C., Goka, C., Tagne, S., Moundipa, F., 2011. Ethnopharmacological survey of plants used for the treatment of female infertility in Baham, Cameroon. *J. Ethnopharmacol.* 136, 178–187.

- Trotter, R.T., Logan, M.H., 1986. Informant Consensus: A New Approach for Identifying Potentially Effective Medicinal Plants. Redgrave Publishing Company, Bedford Hills, NY.
- Umamaheswari, M., Chatterjee, T., 2008. In vitro antioxidant activities of the fractions of *Coccinia grandis* L. leaf extract. Afr. J. Tradit. Complement. Altern. Med. 5, 61–73.
- Valussi, M., Scirè, A.S., 2012. Quantitative ethnobotany and traditional functional foods. Nutrafoods 11, 85–93.
- Venkateswaran, S., Pari, L., 2003. Effect of *Coccinia indica* leaves on antioxidant status in streptozotocin-induced diabetic rats. J. Ethnopharmacol. 84, 163–168.
- Vivek, M., Sunil, S., Pramod, N., Prashith, K., Mukunda, S., Mallikarjun, N., 2013. Anticariogenic activity of *Lagerstroemia speciosa* (L.) Pers. Sci. Technol. Arts Res. J. 1, 53–56.
- Vogler, B., Ernst, E., 1999. *Aloe vera*: a systematic review of its clinical effectiveness. Br. J. Gen. Pract. 49, 823–828.
- Wagh, V.V., Jain, A.K., Kadel, C., 2011. Ethnomedicinal plants used for curing dysentery and diarrhea by tribals of Jhabua district (Madhya Pradesh). Indian J. Nat. Prod. Resour. 2, 256–260.
- Warrier, P.K., Nambiar, V.P., Ramankutty, C., Vasudevan, R., 1995. *Oroxylum indicum*, Indian Medicinal Plants: A Compendium of 500 Species, 1 ed. Orient Longman Private Ltd., Chennai, pp. 180–190.
- Welihinda, J., Karunanayake, E., Sheriff, M., Jayasinghe, K., 1986. Effect of *Momordica charantia* on the glucose tolerance in maturity onset diabetes. J. Ethnopharmacol. 17, 277–282.
- Yang, E.J., Lee, J.S., Yun, C.Y., Ryang, Y.S., Kim, J.B., Kim, I.S., 2011. Suppression of ovalbumin induced airway inflammatory responses in a mouse model of asthma by *Mimosa pudica* extract. Phytother. Res. 25, 59–66.
- Yogendra Kumar, S.H., 2008. Herbal remedies among the Khasi traditional healers and village folks in Meghalaya. Indian J. Tradit. Knowl. 7, 581–586.
- Yong, J.W., Ge, L., Ng, Y.F., Tan, S.N., 2009. The chemical composition and biological properties of coconut (*Cocos nucifera* L.) water. Molecules 14, 5144–5164.
- Zheng, X., Xing, F., 2009. Ethnobotanical study on medicinal plants around Mt. Yinggeling, Hainan Island, China. J. Ethnopharmacol. 124, 197–210.