



* A JEE Notebook

 $Part \ 1: \ on \ Mathematics$

Adyansh Mishra

March 8, 2024











Contents



i Preface

The Philosophy $\,i\,$

1 1 Sets and Manipulations

Naive Set Theory 1 • Equality and Subsets 2











Preface



§1 The Philosophy

First, before jumping to any content, I shall discuss the philosophy behind these notes. These notes are made because I did not like pen-paper notes, these have version control, can be easily rewritten are more accessible(on phone/pc and if needed, on print).

Another thing to note is that these notes follow a specific philosophy, in that they try to be expository. They are meant to explain things. I have noticed that books often talk circularly, create redundancies, and etc.

To combat this, I have tried to clearly lay out everything here. Also, because of how books are made, generally in various volumes, content often goes repeated. These notes will try to remove that

Lastly, I have often added some sort of additional non-JEE relevant content in many places, because 1) JEE is not the only thing I care for, 2) I am particularly interested in Mathematics/Physics so I study extra stuff, 3) They often help in exposition and better understanding. These are, however, mentioned in the text, and you can easily skip them.

Happy learning!













CHAPTER 1



Sets and Manipulations



"You don't have to be great to start, but you have to start to be great."

Zig Ziglar

§1.1 Naive Set Theory

Motivation: There is little Mathematics that does not use the language of sets.

Sets are perhaps the most important structures in all of Mathematics. Although an indepth discussion of sets is beyond these notes, and their axiomation is complicated, we shall not let this affect us and discuss a less rigorous formalisation of sets. Namely, *Naive Set Theory*.

Definition 1.1.1

A set is any collection of elements.

It is denoted as $\{a,b,c\}$ where a,b,c are elements of the set. This representation of sets is called *Roster form*.


Well, what do we mean by any collection? To give a few examples, the set of flowers would be $\{\text{roses, lilies, sunflowers, }\ldots\}$ The trailing ...are used to show that the elements continue on ad infinitum. Hence, any collection of objects is a set.

Often, instead of manually writing out {roses, lilies, sunflowers, ...}, we will use a condition that each element in the set follows. Here, the condition is that all elements all flowers. We denoted this by saying that S is the set of all x such that x is a flower. We represent this in what is called Set-Builder notation. Thus,


$$S = \{x : P(x)\}$$
 or $S = \{x \mid P(x)\}$

Here, P(x) represents the condition. We use the different symbols depending on the context, mostly for ensuring that the statements are clear.

These definitions are superficious
We have not and will not define what we mean by collection or elements. That is what makes this theory naive.







A STATE OF THE STA

Remark 1.1.2. Notice the equality symbol between S and $\{x: P(x)\}$. Generally, we denote sets as capital latin letters A, B, C, \ldots , and their elements by little latin letters a, b, c, \ldots

Example 1.1.3

The sets of all Natural numbers is denoted by \mathbb{N} . Generally, it is either used to represent $\{0,1,2,\ldots\}$ or $\{1,2,3,\ldots\}$. It matters little. We will let $\mathbb{N}=\{1,2,3,\ldots\}$ and $\mathbb{N}_0=\{0,1,2,\ldots\}$ in this text

Example 1.1.4

The other common sets are similarly denoted by blackboard bold letters as well.

- 1. \mathbb{Z} is the set of all integers.
- 2. \mathbb{Q} is the set of all rationals.
- 3. \mathbb{R} is the set of all reals.
- 4. \mathbb{C} is set of all complex numbers.

The symbol \in in $x \in S$ is used to show that x is an element of S. It may be read as 'belongs to'.

Example 1.1.5

The cartesian plane is also a set. It defined as

$$\{(x,y)\mid x,y\in\mathbb{R}\}$$

Extra

The set $\{\emptyset\}$ is not a null set! In-fact it is the

set that contains

the empty or null set.

§1.1.1 Why Naive?

One interesting formulation possible in naive set-theory is the "Russel's Paradox". It goes as follows. Let S be a set $\{x \mid x \notin S\}$. Basically, S is the set of all elements not in S. This leads to a paradox because for x to be in S, it has to follow the condition that it is not in S! And any element not in S must be in S.

There is no particular solution to this using naive set-theory. This is one of the reasons why this discussion of sets is so naive.

§1.1.2 Types of Sets

Although this division of sets is superfluous at best, we will still categorise them for a reference.

- Finite Set: A set having a finite number of elements is called a finite set.
- Infinite Set: A set which has infinite elements is called an infinite set.
- Null Set: A set which has no elements is called a null set. It is represented as {} or ∅.
- Singleton Set: A set containing a single element is called a singleton set.

§1.2 Equality and Subsets

§1.2.1 Sets of Sets

Remember that we referred to sets as a collection of *any* objects? That might lead to the question that can we also make sets of sets? In-fact, that is exactly right, we can! A set of sets maybe represented as $\{\{a,b,c\},\{d,e,f\},\{g,h,i\}\}$. All of the elements of this set are themselves sets. One thing to note here is that we cannot form a set of all sets, and we will see why in a moment.









§1.2.2 Subsets

Consider the sets \mathbb{N} and \mathbb{Z} . You might see that all the elements of \mathbb{N} are also present in \mathbb{Z} . If we were to describe it, we could say that \mathbb{N} is contained in \mathbb{Z} . Such a relation between sets is expressed by saying \mathbb{N} is a subset of \mathbb{Z} .

Definition 1.2.1

A set A is a subset of B if all elements of A are also elements of B. This is represented as

$$A \subseteq B$$

Example 1.2.2

 $\mathbb{N} \subseteq \mathbb{Z} \subseteq \mathbb{Q} \subseteq \mathbb{R} \subseteq \mathbb{C}$.

Example 1.2.3

Let $A = \{x \mid x \text{ is prime and } x \text{ is odd}\}$, and $B = \{x \mid x \text{ is prime}\}$, then $A \subseteq B$.

One thing to note here is that for A to be a subset of B, it does not need to have less elements than B. Infact, A is a subset of A itself because all elements of A are in A.

To show that each element of A is contained in B but all elements of B are not contained in A, we say that A is a proper subset of B.

Note that A is not a proper subset of A.

Definition 1.2.4

A is a proper subset of B if all elements of A are in B but all elements of B are not in A. We represent this as

$$A \subseteq B$$

§1.2.3 Superset

If A is a subset of B, then B is a superset of A. It is represented as

$$B \supseteq A$$

This is another way of saying the same thing. The definition of proper superset is also the same.

§1.2.4 Equality

We may say that two sets are equal if they have the same elements. However, there is one thing to consider. When discussing about sets, we do not care how many times an element appears. The sets $\{1, 1, 1, 1, \ldots\}$ is the same as $\{1\}$. Therefore, we always ignore the repeating terms. Thus, two sets are equal if they have the same number of elements, **regardless of the frequency they appear in.** The same thing applies for subsets and supersets, **the frequency of elements does not matter.**

We can phrase the equality of sets in another manner, in terms of subsets.

Definition 1.2.5 (Equality in sets)

Two sets A and B are equal if and only if $A \subseteq B$ and $B \subseteq A$. Therefore,

$$A = B \iff A \subseteq B \text{ and } B \subseteq A$$



