

# Discrete Mathematics: Lecture 26

Paths and Isomorphism, Counting Paths, Euler Paths and Circuits

Xuming He

Associate Professor

School of Information Science and Technology  
ShanghaiTech University

Spring Semester, 2022


Notes by Prof. Liangfeng Zhang

# Vertex Connectivity

**DEFINITION:** A connected undirected graph  $G = (V, E)$  is said to be **nonseparable**<sub>不可分的</sub> if  $G$  has no cut vertex.

**DEFINITION:** Let  $G = (V, E)$  be a connected simple graph.

- **vertex cut**<sub>点割集</sub>: A subset  $V' \subseteq V$  such that  $G - V'$  is disconnected
- **vertex connectivity**<sub>点连通度</sub>  $\kappa(G)$ : the minimum number of vertices whose removal disconnect  $G$  or results in  $K_1$ ; equivalently,
  - if  $G$  is disconnected,  $\kappa(G) = 0$ ; //additional definition
  - if  $G = K_n$ ,  $\kappa(G) = n - 1$  //  $K_n$  has no vertex cut
  - else,  $\kappa(G)$  is the minimum size of a vertex cut of  $G$



These graphs are all nonseparable


# Edge Connectivity

**DEFINITION:** Let  $G = (V, E)$  be a connected simple graph.  $E' \subseteq E$  is an **edge cut**<sub>边割集</sub> of  $G$  if  $G - E'$  is disconnected.

**DEFINITION:** Let  $G = (V, E)$  be a simple graph.


The **edge connectivity**<sub>边连通度</sub> ( $\lambda(G)$ ) of  $G$  is defined as below:

- $G$  disconnected:  $\lambda(G) = 0$
- $G$  connected:
  - $|V| = 1$ :  $\lambda(G) = 0$
  - $|V| > 1$ :  $\lambda(G)$  is the minimum size of edge cuts of  $G$ .



# Connectivity

**THEOREM:** Let  $G = (V, E)$  be a simple graph. Then  $\kappa(G) \leq \lambda(G) \leq \delta(G)$ , where  $\delta(G) = \min_{v \in V} \deg(v)$  is the least degree of  $G$ 's vertices.



- $\kappa(G) = 2$
- $\lambda(G) = 3$
- $\delta(G) = 4$


[https://cp-algorithms.com/graph/edge\\_vertex\\_connectivity.html](https://cp-algorithms.com/graph/edge_vertex_connectivity.html)

<http://www.math.caltech.edu/~2014-15/2term/ma006b/05%20connectivity%201.pdf>


# Paths and Isomorphism

## Theorem

*The existence of a simple circuit of length  $k$ ,  $k \geq 3$  is an isomorphism invariant for simple graphs.*



$G_1$



$G_2$

6 vertices, 8 edges

Degree sequence: 3, 3, 3, 3, 2, 2

# Paths and Isomorphism\*

## Theorem

*The existence of a simple circuit of length  $k$ ,  $k \geq 3$  is an isomorphism invariant for simple graphs.*

**Proof:** Let  $G_1 = (V_1, E_1)$  and  $G_2 = (V_2, E_2)$  be isomorphic graphs: there is a bijective function  $f : V_1 \rightarrow V_2$  respecting adjacency conditions.

Assume  $G_1$  has a simple circuit of length  $k$ :  $u_0, u_1, \dots, u_k = u_0$ , with  $u_i \in V_1$  for  $0 \leq i \leq k$ . Let's denote  $v_i = f(u_i)$ , for  $0 \leq i \leq k$ .


$(u_i, u_{i+1}) \in E_1 \Rightarrow (f(u_i), f(u_{i+1})) = (v_i, v_{i+1}) \in E_2$ , for  $0 \leq i \leq k - 1$ .

So  $v_0, \dots, v_k$  is a path of length  $k$  in  $G_2$ .


It is a circuit because  $v_k = f(u_k) = f(u_0) = v_0$ .

It is simple: if not, at least one edge is traversed more than once, so it would mean that there exist  $0 \leq i \neq j \leq k - 1$  such that

$(v_i, v_{i+1}) = (v_j, v_{j+1})$ . But this implies  $(u_i, u_{i+1}) = (u_j, u_{j+1})$  by bijectivity of  $f$ . This is impossible because  $u_0, u_1, \dots, u_k$  is simple.



G



H

5 vertices, 6 edges

Degree sequence: 3, 3, 2, 2, 2

1 simple circuit of length 3,

1 simple circuit of length 4,

1 simple circuit of length 5.

Isomorphic graphs ?

If there is an iso  $f : V_G \rightarrow V_H$ , the simple circuit of length 5

$u_1, u_4, u_3, u_2, u_5$  must be sent to the simple circuit of length 5 in H, respecting the degrees of vertices.

Check that  $f(u_1) = v_1, f(u_4) = v_2, f(u_3) = v_3, f(u_2) = v_4, f(u_5) = v_5$  is an isomorphism by writing adjacency matrices.

# Counting Paths Between Vertices

## Theorem

*Let  $G$  be a graph with adjacency matrix  $A$  with respect to the ordering of vertices  $v_1, \dots, v_n$ . The number of different paths of length  $r \geq 1$  from  $v_i$  to  $v_j$  equals the  $(i, j)$  entry of the matrix  $A^r$ .*

**Proof:** By induction

- $r = 1$ : the number of paths of length 1 from  $v_i$  to  $v_j$  is equal to the  $(i, j)$  entry of  $A$  by definition of  $A$ , as it corresponds to the number of edges from  $v_i$  to  $v_j$ .



- Assume the  $(i, j)$  entry of the matrix  $A^r$  is the number of different paths of length  $r$  from  $v_i$  to  $v_j$ .

We can write  $A^{r+1} = A^r A$

Let's denote  $A^r = (b_{ij})_{1 \leq i, j \leq n}$ , and  $A = (a_{ij})_{1 \leq i, j \leq n}$ . The  $(i, j)$  entry of  $A^{r+1}$  is given by:

$$\sum_{k=1}^n b_{ik} a_{kj} = b_{i1} a_{1j} + b_{i2} a_{2j} + \cdots + b_{in} a_{nj} \quad (1)$$


By hypothesis:  $b_{ik}$  equals the number of paths of length  $r$  from  $v_i$  to  $v_k$ .

"Path of length  $r + 1$  from  $v_i$  to  $v_j$  = path of length  $r$  from  $v_i$  to any vertex  $v_k$  + an edge from  $v_k$  to  $v_j$ ."

This is equal to the sum (1).

# Example

How many paths of length four are there from  $a$  to  $d$  in the simple graph  $G$



$G$

with ordering of vertices  $(a, b, c, d, e)$ :

$$A_G = \begin{pmatrix} 0 & 1 & 1 & 0 \\ 1 & 0 & 0 & 1 \\ 1 & 0 & 0 & 1 \\ 0 & 1 & 1 & 0 \end{pmatrix}$$

$$A_G^2 = \begin{pmatrix} 2 & 0 & 0 & 2 \\ 0 & 2 & 2 & 0 \\ 0 & 2 & 2 & 0 \\ 2 & 0 & 0 & 2 \end{pmatrix}$$







$$A_G^3 = \begin{pmatrix} 0 & 4 & 4 & 0 \\ 4 & 0 & 0 & 4 \\ 4 & 0 & 0 & 4 \\ 0 & 4 & 4 & 0 \end{pmatrix}$$

$$A_G^4 = \begin{pmatrix} 8 & 0 & 0 & 8 \\ 0 & 8 & 8 & 0 \\ 0 & 8 & 8 & 0 \\ 8 & 0 & 0 & 8 \end{pmatrix}$$

# Euler Paths and Circuits

**DEFINITION:** Let  $G = (V, E)$  be a graph.

- **Euler Path**<sub>欧拉路径</sub>: a simple path that traverses every edge of  $G$ .
- **Euler Circuit**<sub>欧拉回路</sub>: a simple circuit that traverses every edge of  $G$ .



**Remark:** When  $G$  has multiple edges, these edges will be given different names and considered as different. This is implicit in the textbook.

# Euler Circuits

**THEOREM:** Let  $G = (V, E)$  be a connected multigraph of order  $\geq 2$ .

Then  $G$  has an Euler circuit iff  $2 \mid \deg(x)$  for every  $x \in V$ .

- $\Rightarrow$ : Let  $P: \{x_0, x_1\}, \dots, \{x_{i-1}, x_i\}, \dots, \{x_{n-1}, x_n\}$  be an Euler circuit,  $x_0 = x_n$ 
  - Every occurrence of  $x_i$  in  $P$  contributes 2 to  $\deg(x_i)$ 
    - Every vertex  $x_i$  has an even degree
- $\Leftarrow$ : Let  $P: \{x_0, x_1\}, \dots, \{x_{n-1}, x_n\}$  be a longest simple path in  $G$ .
  - Let  $H = G[P]$ , the subgraph of  $G$  induced by all edges in  $P$ 
    - If  $x_n \neq x_0$ , then  $\deg_H(x_n)$  is odd and so  $P$  cannot be longest.
      - $x_n = x_0$ ,  $P$  is a simple circuit, and  $2 \mid \deg_H(x_i)$  for all  $i$ .
    - If  $\exists i \in \{0, 1, \dots, n-1\}$  such that  $\deg_H(x_i) < \deg_G(x_i)$ ,
      - then  $\exists y \in V$  such that  $\{x_i, y\} \notin P$ 
        - $y, x_i, x_{i+1}, \dots, x_n, x_1, \dots, x_{i-1}, x_i$  is longer than  $P$
    - Hence,  $\deg_H(x_i) = \deg_G(x_i)$  for all  $i \in \{0, 1, \dots, n-1\}$ .
      - $V = \{x_0, x_1, \dots, x_{n-1}\}$  and  $H = G$ .
        - $P$  is an Euler circuit.




**Remark:**  $H$  contains all vertices of  $G$ .  
Otherwise,  $P$  can be extended.

# Construction


## ALGORITHM (Hierholzer):

- **Input:**  $G = (V, E)$ , a connected multigraph,  $2 \mid \deg(x), \forall x \in V$
- **Output:** an Euler circuit
  - **circuit:** = a circuit in  $G$
  - $H := G - \text{circuit} - \text{isolated vertices}$
  - while  $H$  has edges do
    - **subcircuit:** = a circuit in  $H$  that intersects **circuit**
    - $H := H - \text{subcircuit} - \text{isolated vertices}$
    - **circuit:** = **circuit**  $\cup$  **subcircuit**
  - return **circuit**

# Example






**circuit** =  $a, b, e, a$




$H$

# Example







**circuit** =  $a, b, e, a$



$H$

**subcircuit** =  $a, c, f, a$

# Example







**circuit** =  $a, b, e, a$

**subcircuit** =  $a, c, f, a$

$H$






# Example




**circuit** =  $a, b, e, a, c, f, a$

$H$

# Example






**circuit** =  $a, b, e, a, c, f, a$



$H$

**subcircuit** =  $c, d, e, f, b, c$

# Example






**circuit** =  $a, b, e, a, c, f, a$

$H$

**subcircuit** =  $c, d, e, f, b, c$

# Example





**circuit** =  $a, b, e, a, c, d, e, f, b, c, f, a$

# Euler Paths

**THEOREM:** Let  $G = (V, E)$  be a connected multigraph of order  $\geq 2$ . Then  $G$  has an Euler path (not Euler circuit) iff  $G$  has exactly 2 vertices of odd degree.

## ALGORITHM:

- **Input:**  $G = (V, E)$ , a connected multigraph,  $x, y \in V$  have odd degrees
- **Output:** an Euler path
  - $H := G + \{x, y\}$
  - find an Euler circuit using Hierholzer's algorithm
  - remove the edge  $\{x, y\}$  from the circuit




$a, c, d, e, f, b, a, e, b, c, f, a$   
 $a, c, d, e, f, \textcolor{red}{b}, \textcolor{red}{a}, e, b, c, f, a$   
 $\textcolor{red}{a}, e, b, c, f, a, c, d, e, f, \textcolor{red}{b}$


# Hamilton Paths and Circuits

**DEFINITION:** Let  $G = (V, E)$  be a graph.


- **Hamilton Path:** A simple path that passes through every vertex exactly once.
- **Hamilton Circuit:** A simple circuit that passes through every vertex exactly once.




✓ Hamilton path  
✓ Hamilton circuit



✓ Hamilton path  
✓ Hamilton circuit



✓ Hamilton path  
× Hamilton circuit



× Hamilton path  
× Hamilton circuit

# Hamilton Circuits

## Determine if there is a Hamilton circuit in a given graph $G$ ?

- This problem is NP-Complete. //that means very difficult

## Necessary conditions on Hamilton circuit.



- If  $G$  has a vertex of degree 1, then  $G$  cannot have a Hamilton circuit.
- If  $G$  has a vertex of degree 2, then a Hamilton circuit of  $G$  traverses both edges.

## Sufficient conditions on Hamilton circuit.

- **Ore's Theorem:** Let  $G = (V, E)$  be a simple graph of order  $n \geq 3$ . If  $\deg(u) + \deg(v) \geq n$  for all  $\{u, v\} \notin E$ , then  $G$  has a Hamilton circuit.
- **Dirac's Theorem:** Let  $G = (V, E)$  be a simple graph of order  $n \geq 3$ . If  $\deg(u) \geq n/2$  for every  $u \in V$ , then  $G$  has a Hamilton circuit.
  - This is a corollary of Ore's Theorem
    - $\forall u \in V, \deg(u) \geq n/2 \Rightarrow \forall u, v \in V, \deg(u) + \deg(v) \geq n$





# Hamilton Circuits

- Examples (sufficient condition)





# Hamilton Circuits




**Remark:** Dirac's and Ore's Theorems do not give a necessary condition for the existence of a Hamilton circuit!

# Shortest Path Problem

## Definition

A **weighted graph** is a graph  $G = (V, E)$  such that each edge is assigned with a strictly positive number.

The **length** of a path in weighted graph is the sum of the weights of the edges of this path.




$a, b, c$  is a path of length 7 and  $b, e, d, c$  is a path of length 6


**Remark:** Observe that in a non-weighted graph the length of a path is the number of edges in the path!

# Shortest Path Problem

## Examples



distance




time

What is the shortest path in air distance between cities A and E?  
What combination of flights has the smallest total flight time?

# Shortest Path Problem

**Question:** Find the shortest path from  $a$  to  $d$ .




**Method:** Find the closest vertex to  $a$ , then the second closest, the third closest... until we reach  $d$ .

⇒ Dijkstra's algorithm

# Shortest Path Problem


## Remarks:

- Of course in the example above, we could have looked at all the paths between  $a$  and  $d$  and compute their length, but too complicated if the graph has a lot of edges.




- Advantage of Dijkstra's algorithm: we can compute the length of a shortest path from one vertex to all other vertices of the graph.

# Dijkstra's Algorithm



- 1** Find the closest vertex to  $a \rightsquigarrow$  analyse all the edges starting from  $a$ :  
 $a, b$  of length 4  
 $a, f$  of length 2  
 $\Rightarrow f$  is the closest vertex to  $a$ . The shortest path from  $a$  to  $f$  has length 2.
- 2** Find the second closest vertex to  $a \rightsquigarrow$  shortest paths from  $a$  to a vertex in  $\{a, f\}$  followed by an edge from a vertex in  $\{a, f\}$  to a vertex not in this set:  
 $a, b$  of length 4  
 $a, f, e$  of length 5  
 $\Rightarrow b$  is the second closest vertex to  $a$ . The shortest path from  $a$  to  $b$  has length 4.

# Dijkstra's Algorithm



- 3** Find the third closest vertex to  $a \rightsquigarrow$  shortest path from  $a$  to a vertex in  $\{a, f, b\}$  followed by an edge from a vertex in  $\{a, f, b\}$  to a vertex not in this set:  
 $a, b, c$  of length 7  
 $a, b, e$  of length 7  
 $a, f, e$  of length 5  
 $\Rightarrow e$  is the third closest vertex to  $a$ . The shortest path from  $a$  to  $e$  has length 5.
- 4** Find the fourth closest vertex to  $a \rightsquigarrow$  shortest path from  $a$  to a vertex in  $\{a, f, b, e\}$  followed by an edge from a vertex in  $\{a, f, b, e\}$  to a vertex not in this set:  
 $a, b, c$  of length 7  
 $a, f, e, d$  of length 6  
 $\Rightarrow d$  is the fourth closest vertex to  $a$ . The shortest path from  $a$  to  $d$  has length 6.

# Dijkstra's Algorithm

**Goal:** find the length of a shortest path from  $a$  to  $z$  with a series of iterations.

- A distinguished set of vertices is constructed by adding one vertex at each iteration.
- A labeling procedure is carried out at each iteration: a vertex  $w$  is labeled with the length of a shortest path from  $a$  to  $w$  that contains only vertices in the distinguished set.
- The vertex added to the distinguished set is one with minimal label among those vertices not already in the set.

**Notations:**  $S_k :=$  distinguished set after  $k$  iterations,  $L_k(v) :=$  length of a shortest path from  $a$  to  $v$  containing only vertices in  $S_k$  ("label" of  $v$ ).

**Initialization:**  $L_0(a) = 0,$   
 $L_0(v) = \infty$  for every vertex  $v \neq a,$   
 $S_0 = \emptyset.$


**$k$ th iteration:**

- $S_k$  is formed from  $S_{k-1}$  by adding a vertex  $u$  not in  $S_{k-1}$  with smallest label,
- Update the labels of all vertices not in  $S_k$  so that  $L_k(v)$  is the length of a shortest path from  $a$  to  $v$  containing only vertices in  $S_k$ , i.e.

$L_k(v) = \min\{L_{k-1}(v), L_{k-1}(u) + w(u, v)\}$  (with  $w(u, v)$  length of the edge  $(u, v)$ )




# Dijkstra's Algorithm



- **k=0 (initialization):**  $S_0 = \emptyset$ ,  
 $L_0(a) = 0$ ,  $L_0(b) = L_0(c) =$   
 $L_0(d) = L_0(e) = L_0(z) = \infty$


# Dijkstra's Algorithm



- **k=0 (initialization):**  $S_0 = \emptyset$ ,  
 $L_0(a) = 0$ ,  $L_0(b) = L_0(c) =$   
 $L_0(d) = L_0(e) = L_0(z) = \infty$

- **k=1:**  $u := a \rightsquigarrow S_1 = \{a\}$ ,  
 $L_0(a) + w(a, b) = 4 < L_0(b) \rightsquigarrow L_1(b) = 4$   
 $L_0(a) + w(a, c) = 2 < L_0(c) \rightsquigarrow L_1(c) = 2$
- **k=2:**  $u := c \rightsquigarrow S_1 = \{a, c\}$ ,  
 $L_1(c) + w(c, b) = 3 < L_1(b) \rightsquigarrow L_2(b) = 3$   
 $L_1(c) + w(c, d) = 10 < L_1(d) \rightsquigarrow L_2(d) = 10$   
 $L_1(c) + w(c, e) = 12 < L_1(e) \rightsquigarrow L_2(e) = 12$
- **k=3:**  $u := b \rightsquigarrow S_1 = \{a, c, b\}$ ,  
 $L_2(b) + w(b, d) = 8 < L_2(d) \rightsquigarrow L_3(d) = 8$
- **k=4:**  $u := d \rightsquigarrow S_1 = \{a, c, b, d\}$ ,  
 $L_3(d) + w(d, e) = 10 < L_3(e) \rightsquigarrow L_4(e) = 10$   
 $L_3(d) + w(d, z) = 14 < L_3(z) \rightsquigarrow L_4(z) = 14$
- **k=5:**  $u := e \rightsquigarrow S_1 = \{a, c, b, d, e\}$ ,  
 $L_4(e) + w(e, z) = 13 < L_4(z) \rightsquigarrow L_5(z) = 13$
- **k=6:**  $u := z \rightsquigarrow S_1 = \{a, c, b, d, z\}$ ,
- **return:**  $L(z) = 13$


# Traveling Salesperson Problem



**Traveling salesperson problem:** a traveling salesperson wants to visit each of the cities once and return to his starting point. In which order should he visit these cities to travel the minimum total distance?

⇒ **Hamiltonian circuit with minimum total weight in the complete graph.**

# Traveling Salesperson Problem



Route	Tot. dist.
A, B, C, D, E, A	610
A, B, C, E, D, A	516
A, B, E, D, C, A	588
A, B, E, C, D, A	458
A, B, D, E, C, A	540
A, B, D, C, E, A	504
A, D, B, C, E, A	598
A, D, B, E, C, A	576
A, D, E, B, C, A	682
A, D, C, B, E, A	646
A, C, D, B, E, A	670
A, C, B, D, E, A	728

**Traveling salesperson problem:** a traveling salesperson wants to visit each of the cities once and return to his starting point. In which order should he visit these cities to travel the minimum total distance?

⇒ **Hamiltonian circuit with minimum total weight in the complete graph.**