

Numerical analysis(SI211) Fall 2021-22 Homework 2

Prof. Jiahua Jiang

Name: xxx Student No.: xxx E-mail: xxx@shanghaitech.edu.cn

Acknowledgements:

1. Deadline: **2021-12-10 11:59:00**, no late submission is allowed.
2. No handwritten homework is accepted. You should submit your homework in [Blackboard](#) with [PDF](#) format, we recommend you use \LaTeX .
3. Giving your solution in English, solution in Chinese is not allowed.
4. Make sure that your codes can run and are consistent with your solutions, you can use any programming language.
5. Your PDF should be named as "your_student_id+HW2.pdf", package all your codes into "your_student_id+_Code2.zip" and upload. [Don't put your PDF in your code file](#)
6. [All the results from your code should be shown in pdf but please do not inset your code into \$\text{\LaTeX}\$.](#)
7. [Plagiarism is not allowed. Those plagiarized solutions and codes will get 0 point. If the results on the pdf are inconsistent with the results of code, your coding problem will get 0 point.](#)

1. **Numerical differentiation**(10 points.) Assume $f(x) \in C^3$, there are 3 points $f(x_0 - \alpha h), f(x_0), f(x_0 + h)$ with $\alpha > 0$.

- (a) use Lagrange Polynomials to construct an approximation for $f''(x_0)$,
- (b) evaluate the approximation error and find the approximation order.

Solution:

2. **Richardson extrapolation**(10 points.) The $f'(x_0)$ can be expressed as

$$f'(x_0) = \frac{1}{h}(f(x_0 + h) - f(x_0)) - \frac{h}{2}f''(x_0) - \frac{h^2}{6}f'''(x_0) + O(h^3). \quad (1)$$

Use idea of Richardson extrapolation to derive a 3 point formula for $f'(x_0)$ with $O(h^2)$ error. (Hint: Replace step size h with $2h$.)

Solution:

3. **Elements of Numerical Integration**(20 points.) For the integral $\int_2^6 \frac{1}{1+x} dx$, use numerical integration methods to approximate.

- (a) Given only the values of $f(x)$ at $x = 2, 3, 4, 5$ and 6 , use the Midpoint Rule, the Trapezoidal rule to approximate with the smallest step size possible.
- (b) Use Romberg integration to compute $R_{3,3}$.
- (c) Use Gaussian quadrature with $n = 2$ to approximate the integral.

Solutions:

4. **Coding of Simpson's Rule**(20 points.) For integral

$$\int_0^4 e^x dx, \quad (2)$$

we write it in the form:

$$\int_0^4 e^x dx = \sum_{i=0}^{N-1} \left\{ \int_{4i/N}^{4(i+1)/N} e^x dx \right\}, \quad (3)$$

then apply Simpson's rule on each part separately and sum up the results. You need to:

- Plot the actual error of this integral approximation versus N for $N \in \{1, 2, \dots, 100\}$.
- Derive a theoretical bound on the integral approximation in dependence on N and plot this upper bound, too.

Solutions:

5. **Bonus Coding**(Multiple Integrals) (20 points.) For the

$$\iint_{\mathcal{D}} e^{-xy} dx dy \tag{4}$$

with $\mathcal{D} =: \{0 \leq x \leq 1.5, 0 \leq y \leq 2\}$.

- Use Composite Simpson's rule with $n = 6$ and $m = 8$, i.e., $h_x = \frac{1.5}{6}, h_y = \frac{2}{8}$ to approximate (4). (**Note:** your code should input the box range \mathcal{D} and the integers n, m , you can use the Example.1 from the page-239 to debug).

Solution: