

ALDEHYDES AND KETONES

Introduction :


Organic Compounds having $>\text{C}=\text{O}$ group are called carbonyl compounds and $>\text{C}=\text{O}$ group is known as carbonyl or oxo group. Its general formula is $\text{C}_n\text{H}_{2n}\text{O}$ ($n = 1, 2, 3, \dots$) Carbonyl compounds are grouped into two categories.

- (a) **Aldehydes** : Aldehyde group is $\text{—}\overset{\text{O}}{\underset{\text{||}}{\text{C}}}\text{—H}$ (also known as formyl group). It is a monovalent group
- (b) **Ketones** : The carbonyl group ($>\text{C}=\text{O}$) is a Ketonic group when its both the valencies are satisfied by alkyl group. It is a bivalent group.

◆ Ketones are further classified as :


- (i) **Simple or Symmetrical ketones** : Having two similar alkyl groups. $\begin{matrix} \text{R} \\ \text{R} \end{matrix} > \text{C}=\text{O}$
- (ii) **Mixed or unsymmetrical ketones** : Having two different alkyl groups. $\begin{matrix} \text{R} \\ \text{R}' \end{matrix} > \text{C}=\text{O}$

Example : (Ketones) : Symmetrical




(Acetone or Dimethyl ketone)
2-Propanone

Unsymmetrical



(Ethyl methyl ketone)
2-Butanone




In all the compounds given above, lone pair of electrons and double bond are conjugate.

$\left(\begin{matrix} \text{O} \\ \text{||} \\ \text{—C—}\ddot{\text{Z}} \end{matrix} \right)$ so resonance occurs. These compounds have $\text{—}\overset{\text{O}}{\underset{\text{||}}{\text{C}}}\text{—}$ group still they are not carbonyl compounds because carbonyl group takes parts in resonance with the lone pair of electrons.

Structure :

In $>\text{C}=\text{O}$ compounds C-atom is sp^2 hybridised which forms two σ bonds with C and H-atom respectively and one σ bond with oxygen atom. The unhybridised atomic orbital of C-atom and the parallel 2p orbital of oxygen atom give the π bond in $>\text{C}=\text{O}$ group.



Due to electro-negativity difference in C & O atoms, the $>\text{C}=\text{O}$ group is polar.





❑ **General Methods of Preparation :**

(A) For both Aldehydes and Ketones :

(1) By Oxidation of Alcohols :

(a) By $K_2Cr_2O_7 / H_2SO_4$:

Oxidation of primary alcohols gives aldehyde and oxidation of secondary alcohols gives Ketones. Here, $(K_2Cr_2O_7 / H_2SO_4)$ is a strong oxidising agent.



Aldehydes are quite susceptible to further oxidation to acids -





Thus oxidation of primary alcohols is made at the temperature much above the boiling point of aldehyde and thus aldehydes are vapourised out and prevented from being oxidised.

◆ **Note :** Aldehydes can be prepared from 1° alcohol, secondary alcohols can be oxidized to ketones, by oxidation with pyridinium chlorochromate (PCC) in CH_2Cl_2 solvent, pyridinium dichromate (PDC) and with Jones reagent ($CrO_3 + H_2SO_4$) in acetone.

(b) **Oppenaur Oxidation :**


The oxidation of secondary alcohols to ketones by heating them with specific reagent : $[(CH_3)_3CO]_3Al$ (Aluminium-t-butoxide) in presence of acetone. Primary alcohols may also be oxidized to aldehydes if ketones is replaced by a better hydrogen acceptor, e.g. p-benzoquinone. The equilibrium can be controlled by the amount of acetone, an excess of which favours the oxidation of the alcohol.



◆ **Note :** The reaction is the reverse of Meerwein-Ponndorf -verley reduction.


(c) **Mild Oxidising Agent :**


1° alcohols will get oxidised with $CrO_3 / \text{Pyridine}$, **collin's reagent** Ag/O_2 at $250^\circ C$



By this reaction, **good yield of aldehyde** is possible.


(2) **Dehydrogenation of alcohols :**







(3) By Hydrolysis of gem dihalides :

Terminal gem-dihalides on hydrolysis give aldehydes while the non-terminal dihalides give ketone.





(4) By Oxidation of diols :

With periodic acid (HIO_4) & lead tetra acetate $(\text{CH}_3\text{COO})_4\text{Pb}$ vicinal diols gets oxidised to form carbonyl compounds



(5) By Ozonolysis of alkenes :


This reaction is used to determine the position of double bond in alkene.




Unbranched alkene	→	aldehyde
Branched alkene	→	ketone

(6) From Alkyne :





(a) **Hydration :** With dil H_2SO_4 & 1% HgSO_4 at $60-80^\circ\text{C}$.



Other alkynes give ketone :




(b) **Hydroboration** : Reaction with B_2H_6 , 2BH_3 or R_2BH give dialkyl borane.




(7) **By Nef's reaction** :


Nitro alkanes are used in this reaction. The α -H of nitro alkane shows acidic nature.







1 nitro alkane



2 nitro alkane





(8) **By hydrolysis of carbonyl derivatives** :




(9) By oxidation of alkyl halides:

Oxidation takes place by $(\text{CH}_3)_2\text{S}=\text{O}$ dimethyl sulphoxide (DMSO).





(10) From Grignard reagents :


(a) By Cyanides :



(b) By Esters : HCHO can't be prepared by this method.





(c) By acid chlorides :



(11) From β -keto acids :


The decarboxylation reaction takes place via formation of six membered ring transition state.




(B) For Aldehydes only :

(1) Reduction of acyl halides, esters and nitriles :


(a) Acyl chlorides can be reduced to aldehydes by treating them with lithium-tri-tert-butoxyaluminium hydride, $\text{LiAlH}[\text{OC}(\text{CH}_3)_3]$, at -78°C .






(2) From dialkyl Cadmium :


$RCdR'$ (dialkyl Cadmium) is an organometallic compound.





This reaction is superior than Grignard Reaction because the ketones formed, further reacts with Grignard reagent to form 3 alcohols.



(3) From R_2CuLi :






(4) By hydrolysis of Aceto Acetic Ester (AAE) :




❑ Other methods for aldehyde and ketone :


(1) By dry distillation of Ca-salts of carboxylic acid :



Calcium salts of acids other than formic acid on heating together gives ketone






To prepare ethyl methyl ketone Calcium acetate and Calcium propionate are used :




(2) **By Thermal decomposition of carboxylic acids :**

Vapour of carboxylic acids when passed over $\text{MnO}/300^\circ\text{C}$ give carbonyl compounds




(3) **Wacker process :**

In this reaction double bond is not cleaved so same C-atom aldehyde and ketones are formed.



All other alkenes gives ketone.



□ **Physical Properties :**

◆ **State :**


Only formaldehyde is gas, all other carbonyl compounds upto C_{11} are liquids and C_{12} & onwards solid.

◆ **Odour :**

Lower aldehydes give unpleasant smell, higher aldehydes and all ketones have pleasant smell.

◆ **Solubility :**

C_1 to C_3 (formaldehyde, acetaldehyde and propionaldehyde) and acetone are freely soluble in water due to polarity of $\begin{array}{c} \delta+ \quad \delta- \\ \text{>C=O} \end{array}$ bond and can form H-bond with water molecule. C_5 onwards are insoluble in water.



◆ **Boiling point :** Boiling point \propto Molecular weight

Boiling point order is - **Alcohol** > **Carbonyl compounds** > **Alkane**

This is because in alcohols intermolecular H-bonding is present but in carbonyl compounds H-bonding doesn't exist, instead dipole-dipole vander waal force of attraction is present. Alkanes are non polar.




◆ **Density** : Density of carbonyl compounds is lower than water.

□ **Chemical Properties** :

Reactions of both aldehydes and ketones :

Due to strong electronegativity of oxygen, the mobile π electrons pulled strongly towards oxygen, leaving the carbon atom deficient of electrons.




Carbon is thus readily attacked by $\ddot{\text{N}}\text{u}$. The negatively charged oxygen is attacked by electron deficient (electrophile) E^+ .

$>\text{C}=\text{O}$ bond in carbonyl group is stronger than $\text{C}=\text{C}$ bond in alkanes.




Reactivity of carbonyl group \propto **Magnitude of +ve charge** \propto **- I group** $\propto \frac{1}{+\text{I group}}$

Ex : Why carbonyl compound gives nucleophilic addition reaction (NAR) ?




Ex. Arrange the following for reactivity in decreasing order



Sol. (A) $\text{I} > \text{II} > \text{III}$ (B) $\text{II} > \text{I} > \text{III} > \text{IV}$ (C) $\text{IV} > \text{III} > \text{II} > \text{I}$ (D) $\text{IV} > \text{I} > \text{II} > \text{III}$

[Hint : CH_3- is +I group, decreases the intensity of +ve charge on C-atom of $>\text{C}=\text{O}$ group.

$\text{Cl}-$ is -I group increases the intensity of +ve charge on C-atom of $>\text{C}=\text{O}$ group.]






Sol. I is more reactive than II.

❑ Chemical Reactions :

Carbonyl compounds in general under goes nucleophilic addition reaction :


(A) Nucleophilic addition reactions :

(1) Addition of HCN :





(2) Addition of NaHSO_3 :

This reaction is utilized for the separation of carbonyl compounds from non - carbonyl compounds.




Sodium bi sulphite

Bisulphite compound
(Crystalline)




(3) Reaction with ammonia derivatives :

These are condensation or addition elimination reaction. These proceeds well in weakly acidic medium.





Addition - elimination (Condensation)



Mechanism :







◆ Ammonia derivatives (NH_2Z) :





2, 4-Dinitro phenyl hydrazine (DNP) Brady's reagent.










(4) With alcohol and thioalcohol :




Tri ethoxy methane $[\text{HC}(\text{OC}_2\text{H}_5)_3]$ remove the water formed during the reaction and so the reaction proceeds in forward direction.




All sulphones compounds are hypnotic compounds.




(5) Reaction with glycol (group protection) :




(6) Reaction with sodium alkynide :




(7) Reaction with Grignard reagent :



Ethanol (1 alcohol)




2-Propanol (2 alcohol)



2-Methyl-2-propanol (3 alcohol)

(8) Reaction with H_2O : It is a reversible reaction.



Ex : Which compound form more stable hydrate with H_2O ?

(A) HCHO

(B) CH_3CHO

(C) CH_3COCH_3


(D) $\text{CH}_3\text{COC}_2\text{H}_5$

[Hint : HCHO since it is more reactive towards this reaction.]

Ans. (A)

Ex : Which carbonyl compound not gives reversible reaction with water ?

Sol. Chloral hydrate.




(Chloral)

(Chloral hydrate)

Stable by intra molecular hydrogen bonding.



(B) Other Reactions :

(c) Proton exchange





When molecules are same	→	Simple cannizaro reaction
Two different molecules	→	Mixed cannizaro reaction.

In mixed or crossed cannizaro reaction more reactive aldehyde is oxidised and less reactive aldehyde is reduced.



(3) Tischenko reaction :





It is a modified cannizaro reaction. All aldehydes undergo this reaction in presence of $(\text{C}_2\text{H}_5\text{O})_3\text{Al}$, to form ester.




(4) Reaction With Halogen :


(a) Replacement of α-H atoms :

This reaction is not shown by formaldehyde (HCHO), since α -H atoms are absent, as enolisation does not takes place in HCHO.






(b) **Replacement of O-atom of $>\text{C}=\text{O}$ group :** It takes place by PCl_5 or SOCl_2 .




Phosphorus penta chloride



Thionyl chloride


(c) **Haloform reactions :**

Chlorine or bromine replaces one or more α -hydrogen atoms in aldehydes and ketones, e.g., acetone may be brominated in glacial acetic acid to give monobromoacetone :



The halogenation of carbonyl compounds is catalysed by acids and bases. Let us consider the case of acetone. In alkaline solution, tribromoacetone and bromoform are isolated. Thus, the introduction of a second and a third bromine atom is more rapid than the first. In aqueous sodium hydroxide, the rate has been shown to be independent of the bromine concentration, but first order with respect to both acetone and base i.e.,

$$\text{Rate} = k [\text{acetone}] [\text{OH}^\ominus]$$



(5) **Aldol Condensation :**

Two molecules of an aldehyde or a ketone undergo condensation in the presence of a base to yield a β -hydroxyaldehyde or a β -hydroxyketone. This reaction is called the aldol condensation. In general Carbonyl compounds which contain α -H atoms undergo aldol condensation with dil. NaOH. Aldol contains both alcoholic and carbonyl group.


□ **Mechanism of aldol condensation :** It takes place in the following two stages :

(a) Formation of Carbanion


(b) Combination of carbanion with other aldehyde molecule.

(a) **Formation of Carbanion :**


α -H atom of $>\text{C}=\text{O}$ group are quite acidic which can be removed easily as proton, by a base.



Carbanion thus formed is stable because of resonance -



(b) Combination of carbanion with other aldehyde molecule :




Aldol condensation is possible between :


1. Two aldehyde (Same or different)
2. Two ketones (Same or different)
3. One aldehyde and one ketone


Identical carbonyl compounds	→ Simple or self aldol condensation.
Different carbonyl compounds	→ Mixed or crossed aldol condensation.

Simple or Self condensation :







Mixed or Crossed aldol Condensation :






Ex : $\text{CH}_3\text{CHO} + \text{CH}_3\text{CH}_2\text{CHO} \xrightarrow[\text{(WB)}]{\text{OH}^\ominus}$ Total 4 products. Write structure of products ?



□ **Intramolecular aldol condensation :**




(α,β - Unsaturated Ketone)

(Aldol)


Here 5 membered ring is more stable than 3 membered ring so above product is formed as a major product.

Note :

- ◆ If in crossed aldol condensation reaction , only one carbonyl compound have α -H than total two product formed.




(Crotonaldehyde)




(6) Claisen condensation :

When two molecules of ester undergo a condensation reaction, the reaction is called Claisen condensation. The product of the claisen condensation is a β -keto ester.





After nucleophilic attack, the aldol addition and the Claisen condensation differ. In the claisen condensation, the negatively charged oxygen reforms the carbon oxygen π -bond and eliminates the OR group.

Mixed claisen condensation :



(7) Intramolecular claisen condensation :




Dieckmann condensation : The addition of base to a 1,6-diester causes the diester to undergo intramolecular claisen condensation, thereby forming a five membered ring β -keto ester. An intramolecular claisen condensation is called a Dieckmann condensation.



(8) Perkin reaction :


In perkin reaction, condensation has been effected between aromatic aldehydes and aliphatic acid anhydride in the presence of sodium or potassium salt of the acid corresponding to the anhydride, to yield α , β -unsaturated aromatic acids.

The acid anhydride should have at least two α -H.




(9) Knoevenagel Reaction :

Condensation of aldehydes and ketones with compounds having active methylene group in the presence of basic catalyst to form α, β -unsaturated compounds is called Knoevenagel Reaction. The basic catalyst may be ammonia or its derivative. Thus 1, 2, 3 amines i.e., aniline, di-or tri - alkyl amines, pyridine or piperidine are used.





(10) Reformatsky Reaction :

A similar reaction like the addition of organometallic compounds on carbonyl compounds that involves the addition of an organozinc reagent to the carbonyl group of an aldehyde or ketone. This reaction, called Reformatsky reaction, extends the carbon skeleton of an aldehyde or ketone and yields β -hydroxy esters. It involves treating an aldehyde or ketone with an α -bromo ester in the presence of zinc metal; the solvent most often used is benzene. The initial product is a zinc alkoxide, which must be hydrolysed to yield the β -hydroxy ester.



(11) Schmidt Reaction :


This is the reaction between a carbonyl compound and hydrazoic acid in the presence of a strong acid concentrated sulphuric acid. Aldehydes give a mixture of cyanide and formyl derivatives of primary amines, whereas ketones give amides :




Schiff's Base

(12) Benzoin condensation :

The benzoin condensation is essentially a dimerisation of two aromatic aldehydes under the catalytic influence of cyanide ions to give benzoin (I).




The hydrogen atom attached to the carbonyl group of aldehyde is not active enough to be removed easily but the addition of the cyanide ion to the carbonyl carbon places this hydrogen in the alpha position of the nitrile thus rendering it relatively acidic. The carbanion, thus generated, attacks the carbonyl carbon of the second aldehyde molecule in a rate-determining step forming an unstable cyanohydrin of benzoin which immediately breaks down into benzoin and hydrogen cyanide.




(13) Benzilic acid rearrangement :

The addition of a strong base to a carbonyl group results in the formation of an anion. The reversal of the anionic charge may cause expulsion of the attached group X, e.g.




However, in a 1, 2-diketone the group X may migrate to the adjacent electron-deficient carbonyl carbon forming α -hydroxy acid.

Thus, benzil on treatment with a strong base forms benzilic acid (salt), hence the name benzilic acid rearrangement.



(14) The Beckmann rearrangement :

The acid catalysed transformation of a ketoxime to an N-substituted amide is known as the Beckmann rearrangement.





The rearrangement is catalysed by a variety of acidic reagents such as H_3PO_2 , H_2SO_4 , $SOCl_2$, PCl_5 , etc.

(C) Oxidation Reactions :

(a) By $K_2Cr_2O_7/H_2SO_4$:

On oxidation with $K_2Cr_2O_7/H_2SO_4$ 1^o alc. gives aldehyde, which on further oxidation gives acid with same C-atom. While, 2^o alcohol on oxidation gives ketone which on further oxidation gives acid with less C-atom.





(i) 3^o alcohol is not oxidised within 2 or 3 minutes.

(ii) 1^o and 2^o alcohol converts orange colour of $K_2Cr_2O_7$ to green in 2-3 minutes.

(b) SeO_2 (Selenium Oxide) :


Ketones or aldehydes on oxidation with SeO_2 gives dicarbonyl compounds. This reaction is possible only in compounds containing α -carbon.

HCHO doesn't show this reaction.




(c) **Baeyer's Villiger oxidation :**





Both aldehyde and ketones are oxidized by peroxy acids. This reaction, called the Baeyer-villiger oxidation, is especially useful with Ketones, because it converts them to carboxylic esters. For example, treating acetophenone with a peroxy acid converts it to the ester phenyl acetate.



Mechanism :




The product of this reaction shows that a phenyl group has a greater tendency to migrate than a methyl group. Had this not been the case, the product would have been C₆H₅COOCH₃ and not CH₃COOC₆H₅. This tendency of a group to migrate is called migratory aptitude. Studies of the Baeyer-villiger oxidation and other reactions have shown that the migratory aptitude of groups is H > phenyl > 3° alkyl > 2° alkyl > 1° alkyl > methyl. In all cases, this order is for groups migrating with their electron pairs, that is, as anions.




(D) Reduction :**(a) The wolf kishner reduction :**

When a ketone or an aldehyde is heated in a basic solution of hydrazine, the carbonyl group is converted to a methylene group this process is called Deoxygenation because an oxygen is removed from the reactant. The reaction is known as the Wolf-kishner Reduction.

**(b) Clemmensen Reduction :**

The reduction of carbonyl groups of aldehydes and ketones to methylene groups with amalgamated zinc and concentrated hydrochloric acid is known as Clemmensen reduction.




The nature of product depends upon the reducing agent used. It can be summarized as.



Reducing agents are



- ◆ Red P/HI at 150 C
- ◆ Zn-Hg/HCl [Clemmensen's reduction]
- ◆ $NH_2-NH_2/C_2H_5OH, OH^\ominus$ [Wolff Kischner's reduction]



Reducing agents are

- ◆ $LiAlH_4$ (Nitetron brown)
- ◆ Na/C_2H_5OH (Bouval blank)
- ◆ $NaH/Benzene$ (Darzen reaction)
- ◆ $[(CH_3)_2CHO]_3Al$ (Aluminium isopropoxide)
- ◆ $(CH_3)_2CHOH$ (Isopropyl alcohol)

Reduction with aluminium isopropoxide in excess of isopropanol is called MPV (**Meerwein Ponndroff Verley**) reduction. Other reducible groups are not attacked like $-NO_2$, $-CH=CH_2$, $-C\equiv C-$.

**(E) Reactions given by only aldehydes :****(1) Polymerisation :** It is a reversible process.**Formaldehyde :**

Formalin


(40% HCHO)

Paraformaldehyde

$n = 6-50$


Hydrated white crystal

Paraformaldehyde is a linear polymer which show reducing character with Tollen's reagent, Fehling solution etc.



Poly oxy methylene



$n > 100$



Meta formaldehyde (Trioxane)

Cyclic polymer (Trioxymethylene)



Cyclic polymer doesn't show reducing character with Tollen's reagent etc.



Formose sugar

A linear polymer (α -acrose)

Acetaldehyde :





Para acetaldehyde

Paraldehyde (cyclic polymer)

Pleasant smelling liquid

Hypnotic compound



Meta aldehyde


White crystalline solid.

Cyclic polymer

Used as solid fuel or killing snails

(4) Reaction with ammonia :

Except formaldehyde, all other aldehydes give addition reactions (HCHO give addition elimination i.e. condensation reaction)







Urotropine (Hexamine)

White crystalline solid

Used in preparation of explosive

Used in treatment of urine infection diseases





Trimethyl hexahydro Triazine trihydrate

(5) Reducing character :


Aldehydes are easily oxidised so they are strong reducing agents.

(a) Tollen's reagent :

It oxidises aldehydes. Tollen's reagent is ammoniacal silver nitrate solution



Silver mirror




(b) Fehling's solution :

It is a mixture of CuSO_4 , NaOH and sodium potassium tartrate.


Fehling solution A- (aq.) solution of CuSO_4

Fehling solution B- Roschelle salt (Sodium potassium tartrate + NaOH)

Fehling solution A + Fehling's solution B (Dark blue colour of cupric tartrate)



(Cuprous oxide-Red ppt.)




(Cupric - Blue)

(Cuprous - Red ppt.)

(c) Benedict's solution :



It is a mixture of CuSO_4 + sodium citrate + Na_2CO_3 . It provides Cu^{+2} . It is reduced by aldehyde to give red ppt of cuprous oxide.



(Cuprous oxide-Red ppt.)

(d) **Mercuric chloride :**

HgCl_2 is a corrosive sublimate. It is reduced by aldehyde to give white ppt of mercurous chloride (Calonal) which further react with aldehyde to give black ppt of Hg.




(e) **Reaction with schiff's reagent :**

Schiff's reagent is dil solution of p-roseniline hydrochloride or magenta dye.


Its pink colour is discharged by passing SO_2 gas and the colourless solution is called schiff's reagent, Aldehyde reacts with this reagent to restore the pink colour.

(F) **Reaction of only ketones :**


(1) **Reduction :** Acetone is reduced by magnesium amalgam and water to give pinacol.




(2) **Reaction with chloroform :**



(3) **Reaction with HNO_2 :**





(4) **Oxidation reaction :** According to popoff's rule >C=O group stays with smaller alkyl group.




(5) **Condensation reaction :**


(a) In presence of dry HCl - aldol condensation takes place




(b) In presence of conc. H_2SO_4



(6) Reaction with ammonia :




(7) Pyrolysis :








TEST FOR HCHO , CH_3CHO , CH_3COCH_3

S.No.	Test	HCHO	CH_3CHO	CH_3COCH_3
1.	Legel's test : $\text{Na} [\text{Fe}(\text{NO})(\text{CN})_3]$ sodium nitroprusite (alk.) Only methyl $>\text{C}=\text{O}$ compound gives this test	-	Red	Red
2.	Iodoform test $(\text{I}_2 + \text{NaOH})$	-	- yellow ppt	- yellow ppt
3.	Pyragallol	white ppt.	-	-
4.	Orthonitro benzaldehyde	-	-	Blue
5.	Tollen's reagent - Fehling's reagent - Mercuric chloride - Schiff's reagent -	Silver mirror Red ppt Black ppt Pink colour	Silver mirror Red ppt Black ppt Pink colour	- - - -
6.	DNP	Orange colour	Orange colour	Orange colour







SOLVED EXAMPLES


Ex 1. What is A in the following reaction ?




Ex 2. is the final product obtained when one of the following is reacted with base :







Sol.




Ex 3.









The product (A) in the given reaction would be :










Ex 4. End product of the following sequence of reactions is :




∴ (B)

Ex 5. In which of the following substrates, rate of Benzoin condensation will be maximum ?



Benzoin condensation is due to stability of intermediate (III) when negative charge on C extensively delocalised in benzene ring, nitro and C≡N group. In all other cases, such dispersal is not extensively possible. On the other hand, NO₂⁻ is also creating a positive charge center on carbonyl carbon, making it more susceptible to nucleophilic attack of CN⁻.

Ex 6. This intermediate is converted into product in the wittig reaction :




Out of following which statements are correct ?

- (A) C-O bond is weaker as compared to P-O bond
- (B) Lone pair of oxygen atom participate in $p\pi-d\pi$ bonding with phosphorous atom
- (C) C-P bond is weaker as compare to C-C bond
- (D) C-C bond is weaker as compare to C-O bond

Sol. (A), (B), (C)

Ex 7. Consider the following sequence :



Which of following statements are correct for above reaction sequence ?

- (A) Step I is acid-base reaction
- (B) Step II is nucleophilic addition reaction
- (C) Step III is acid base reaction
- (D) Step IV is elimination reaction

Sol. (A), (B), (C), (D)

Ex 8. Which of the following oxidation reaction can be carried out with chromic acid in aqueous acetone at 5-10 C.

- (A) $\text{CH}_3(\text{CH}_2)_3\text{C}\equiv\text{C}-\underset{\text{OH}}{\text{CH}}-\text{CH}_3 \longrightarrow \text{CH}_3(\text{CH}_2)_3\text{C}\equiv\text{C}-\overset{\text{O}}{\underset{\text{||}}{\text{C}}}-\text{CH}_3$
- (B) $\text{CH}_3(\text{CH}_2)_3\text{CH}=\text{CH}-\text{CH}_2\text{OH} \longrightarrow \text{CH}_3(\text{CH}_2)_3\text{CH}=\text{CH}-\text{CHO}$
- (C) $\text{C}_6\text{H}_5\text{CH}_3 \longrightarrow \text{C}_6\text{H}_5\text{COOH}$
- (D) $\text{CH}_3(\text{CH}_2)_3\text{CH}_2\text{OH} \longrightarrow \text{CH}_3(\text{CH}_2)_3\text{CHO}$

Sol. (A), (B), (C), (D)

Ex 9. $\text{CH}_3-\overset{\text{O}}{\underset{\text{||}}{\text{C}}}-\text{CH}_3 \xrightarrow{\text{SeO}_2} \text{A}$; A will :

- (A) Reduce Tollen's reagent
- (B) Give iodoform test
- (C) Form oxime
- (D) Give Cannizaro reaction

Sol. SeO_2 oxidises - α - CH_2 - a w.r.t. keto group

\therefore (A), (B), (C) and (D)

Ex 10. $3\text{HCHO} + \text{CH}_3\text{CHO} \xrightarrow{\text{NaOH}} \text{A}$. A found can


(A) Reduce Tollen's reagent

(B) Give Cannizaro reaction

(C) React with Na




(D) Give green colour with $\text{Cr}_2\text{O}_7^{2-}/\text{H}^+$

Sol. A is by aldol condensation



\therefore (A), (B), (C) and (D)



Ex 11. $2\text{CH}_3 - \overset{\text{O}}{\underset{\text{O}}{\text{C}}} - \text{CH}_3 \xrightarrow[\text{H}^+]{\text{Mg/Hg}}$ Product, product in the reaction is :



(D) None of these



Sol. (A) $2\text{CH}_3 - \text{CO} - \text{CH}_3 \xrightarrow[\text{H}_2\text{O}]{\text{Mg/Hg}} \begin{array}{c} \text{CH}_3 \quad \text{CH}_3 \\ | \quad | \\ \text{H}_3\text{C} - \text{C} - \text{C} - \text{CH}_3 \\ | \quad | \\ \text{OH} \quad \text{OH} \end{array}$
(Pinacol)

Ex 12. Benzaldehyde on reaction with acetophenone in the presence of sodium hydroxide solution gives :



Sol. (A) $\text{C}_6\text{H}_5\text{CHO} + \text{CH}_3\text{COC}_6\text{H}_5 \xrightarrow[-\text{H}_2\text{O}]{\text{NaOH}} \text{C}_6\text{H}_5 - \text{CH} = \text{CH} - \overset{\text{O}}{\underset{\text{O}}{\text{C}}} - \text{C}_6\text{H}_5$

Ex 13. Product in following reaction is :



Sol. (C) $\text{H} - \text{CHO} + \text{CH}_3\text{MgI} \rightarrow \text{CH}_3 - \text{CH}_2 - \text{OH} + \text{Mg} \begin{array}{c} \text{OH} \\ \diagup \\ \text{I} \end{array}$
