

Quantum-Classical Equivalence for AND Functions

CCC 2026

Sreejata Kishor Bhattacharya, Farzan Byramji, Arkadev Chattopadhyay, Yogesh Dahiya, Shachar Lovett

Tata Institute of Fundamental Research
University of California San Diego

Quantum-Equivalence Conjecture: For a total function
 $f : \{0, 1\}^n \rightarrow \{0, 1\}$,

$$R(f) \leq \text{poly}(C(f))$$

where

- $R(f)$ = classical randomized communication complexity
- $Q(f)$ = quantum communication complexity (with prior entanglement)

Major open problem in communication complexity. First formally proposed by Shi and Zhu

Our result: For AND functions, $f = g \circ \text{AND}_2$,

$$D(f) \leq \text{poly}(C(f), \log(n))$$

where $D(f)$ = classical deterministic communication complexity.
Thus, Quantum Equivalence Conjecture is true for AND function up to $\text{polylog}(n)$ factors.

This answers an open problem of Razborov [2002].

Why study AND functions?

- Lifted functions form a large class of interesting communication problems. They form a testbed for all major conjectures in communication complexity (e.g, Log Rank Conjecture, Log Approximate Rank Conjecture [now known to be false], Quantum Equivalence Conjecture)

Why study AND functions?

- Lifted functions form a large class of interesting communication problems. They form a testbed for all major conjectures in communication complexity (e.g, Log Rank Conjecture, Log Approximate Rank Conjecture [now known to be false], Quantum Equivalence Conjecture)
- Of particular interest is the case when the lifting gadget is as simple as possible, i.e., it has only two bits. In this case, the gadget does not have too much ability to obfuscate the inputs between Alice and Bob.

Why study AND functions?

- Lifted functions form a large class of interesting communication problems. They form a testbed for all major conjectures in communication complexity (e.g, Log Rank Conjecture, Log Approximate Rank Conjecture [now known to be false], Quantum Equivalence Conjecture)
- Of particular interest is the case when the lifting gadget is as simple as possible, i.e., it has only two bits. In this case, the gadget does not have too much ability to obfuscate the inputs between Alice and Bob.
- There are only two nontrivial gadgets on 2 bits: XOR and AND. Both XOR and AND functions have been extensively studied in the literature.

AND functions

- There is a base function $g : \{0, 1\}^n \rightarrow \{0, 1\}$. Alice has an input $x \in \{0, 1\}^n$, Bob has an input $y \in \{0, 1\}^n$. Their goal is to compute

$$g(x_1 \wedge y_1, \dots, x_n \wedge y_n)$$

- **Razborov (2002)** proved the Equivalence result for functions of the form $g \circ \text{AND}_2$, where g is a symmetric function.

- **Razborov (2002)** proved the Equivalence result for functions of the form $g \circ \text{AND}_2$, where g is a symmetric function.
- An easy adaptation of the same argument proves the results for functions of the form $g \circ h$, where h is any gadget inside whose truth table both AND and OR can be embedded.

- **Razborov (2002)** proved the Equivalence result for functions of the form $g \circ \text{AND}_2$, where g is a symmetric function.
- An easy adaptation of the same argument proves the results for functions of the form $g \circ h$, where h is any gadget inside whose truth table both AND and OR can be embedded.
- **Burhman and deWolf (2003)** noticed a relation between $Q(g \circ \text{AND}_2)$ and various complexity measures of g , and conjectured a particular relation between them (to be described in detail soon).

- **Razborov (2002)** proved the Equivalence result for functions of the form $g \circ \text{AND}_2$, where g is a symmetric function.
- An easy adaptation of the same argument proves the results for functions of the form $g \circ h$, where h is any gadget inside whose truth table both AND and OR can be embedded.
- **Burhman and deWolf (2003)** noticed a relation between $Q(g \circ \text{AND}_2)$ and various complexity measures of g , and conjectured a particular relation between them (to be described in detail soon).
- **Kremer (1996)** proved log-approximate rank lower bounds quantum communication complexity.

- **Razborov (2002)** proved the Equivalence result for functions of the form $g \circ \text{AND}_2$, where g is a symmetric function.
- An easy adaptation of the same argument proves the results for functions of the form $g \circ h$, where h is any gadget inside whose truth table both AND and OR can be embedded.
- **Burhman and deWolf (2003)** noticed a relation between $Q(g \circ \text{AND}_2)$ and various complexity measures of g , and conjectured a particular relation between them (to be described in detail soon).
- **Kremer (1996)** proved log-approximate rank lower bounds quantum communication complexity.
- **Lovett, Knop, McGuire, Yuan (2021)** proved the log rank conjecture for AND functions. By an intermediate result in their proof, it became clear that proving the relation conjectured by Burhman and deWolf would also imply the Quantum Equivalence Conjecture for AND functions.

- **Chattopadhyay, Dahiya, Lovett (2026)** proved that de-Morgan sparsity and approximate de-Morgan sparsity are polynomially related.

- **Chattopadhyay, Dahiya, Lovett (2026)** proved that de-Morgan sparsity and approximate de-Morgan sparsity are polynomially related.
- They also show the Quantum Equivalence Conjecture for functions of the form $g \circ \text{EQ}_4$.

- **Chattopadhyay, Dahiya, Lovett (2026)** proved that de-Morgan sparsity and approximate de-Morgan sparsity are polynomially related.
- They also show the Quantum Equivalence Conjecture for functions of the form $g \circ \text{EQ}_4$.
- We extend their work to show the relation between quantities conjectured by Burhman and deWolf. This implies the Quantum-Equivalence Conjecture for AND functions.

- A result of Kremer ¹:

$$Q^{cc}(F) \geq \text{poly}(\log(\text{rank}^{1/3}(F)))$$

where

$$\text{rank}^{1/3}(F) = \inf\{\text{rank}(G) \mid \|G - F\|_{\infty} \leq 1/3\}$$

¹The result of Kremer only lower bounds clean-quantum communication complexity.[BdW01] observed that it readily implies general quantum communication lower bounds too. Another proof was given by Lee and Shraibman [LS07]

- A result of Kremer ¹:

$$Q^{cc}(F) \geq \text{poly}(\log(\text{rank}^{1/3}(F)))$$

where

$$\text{rank}^{1/3}(F) = \inf\{\text{rank}(G) \mid \|G - F\|_{\infty} \leq 1/3\}$$

- Thus, log-approximate rank conjecture for a class of functions also implies quantum-equivalence conjecture for that class of functions.

¹The result of Kremer only lower bounds clean-quantum communication complexity.[BdW01] observed that it readily implies general quantum communication lower bounds too. Another proof was given by Lee and Shraibman [LS07]

- A result of Kremer ¹:

$$Q^{cc}(F) \geq \text{poly}(\log(\text{rank}^{1/3}(F)))$$

where

$$\text{rank}^{1/3}(F) = \inf\{\text{rank}(G) \mid \|G - F\|_{\infty} \leq 1/3\}$$

- Thus, log-approximate rank conjecture for a class of functions also implies quantum-equivalence conjecture for that class of functions.
- It is known that LARC is false in general (**Mande, Suhail, Chattopadhyay, 2019**). Nevertheless, in this work we show LARC is true for AND functions (upto $\text{polylog}(n)$ factors); thereby also establishing the QEC for AND functions.

¹The result of Kremer only lower bounds clean-quantum communication complexity.[BdW01] observed that it readily implies general quantum communication lower bounds too. Another proof was given by Lee and Shraibman [LS07]

In case Yogesh has defined this properly in his previous talk I shall skip the next few slides.

- Any function $g : \{0, 1\}^n \rightarrow \mathbb{R}$ has a unique representation as a multilinear polynomial:

$$g(x) = \sum_{S \subseteq [n]} \alpha_S x_S$$

where $x_S = \prod_{i \in S} x_i$.

Note: This is not the Fourier decomposition where the domain is $\{\pm 1\}$. Also note that we are working over $\mathbb{R}$, not $\mathbb{F}_2$.

- Any function $g : \{0, 1\}^n \rightarrow \mathbb{R}$ has a unique representation as a multilinear polynomial:

$$g(x) = \sum_{S \subseteq [n]} \alpha_S x_S$$

where $x_S = \prod_{i \in S} x_i$.

Define sparsity of g , $\text{spar}(g)$, to be the number of nonzero monomials in this representation.

- The conjecture in [BdW01]:

$$\text{rank}^{1/3}[g \circ \text{AND}_2] \approx \text{spar}(g)$$

²Other direction is trivial

- The conjecture in [BdW01]:

$$\text{rank}^{1/3}[g \circ \text{AND}_2] \approx \text{spar}(g)$$

- Our main result:

$$\log \text{spar}(g) \leq \text{poly} \left(\log \left(\text{rank}^{1/3}(g \circ \text{AND}_2), \log(n) \right) \right)$$

2

²Other direction is trivial

How this implies the Quantum Equivalence Conjecture

- Recall we want to upper bound $D(f)$ in terms of $\log \left(\text{rank}^{1/3}(f \circ \text{AND}_2) \right)$

How this implies the Quantum Equivalence Conjecture

- Recall we want to upper bound $D(f)$ in terms of $\log \left(\text{rank}^{1/3}(f \circ \text{AND}_2) \right)$
- **Knop, Lovett, McGuire, Yuan** showed that

$$\log D(g \circ \text{AND}_2) \leq \text{poly}(\log \text{spar}(g), \log(n))$$

How this implies the Quantum Equivalence Conjecture

- Recall we want to upper bound $D(f)$ in terms of $\log \left(\text{rank}^{1/3}(f \circ \text{AND}_2) \right)$
- **Knop, Lovett, McGuire, Yuan** showed that

$$\log D(g \circ \text{AND}_2) \leq \text{poly}(\log \text{spar}(g), \log(n))$$

- Since our result upper bounds $\text{spar}(g)$ in terms of $\text{rank}^{1/3}(g \circ \text{AND}_2)$, the result immediately follows.

Goal: Upper bound $\text{spar}(g)$ in terms of $\text{rank}^{1/3}(g \circ \text{AND}_2)$

- Assume, for the sake of contradiction, $\text{spar}(g)$ is large and $\text{rank}^{1/3}(g \circ \text{AND}_2)$ is small.

Goal: Upper bound $\text{spar}(g)$ in terms of $\text{rank}^{1/3}(g \circ \text{AND}_2)$

- Assume, for the sake of contradiction, $\text{spar}(g)$ is large and $\text{rank}^{1/3}(g \circ \text{AND}_2)$ is small.
- From the second assumption, we have functions

$$a_i, b_i : \{0, 1\}^n \rightarrow [0, 1], i = 1, 2, \dots, m$$

and constants $\alpha_i, i = 1, 2, \dots, m$ such that

$$\left| g(x \wedge y) - \sum_{i=1}^m \alpha_i a_i(x) \wedge b_i(y) \right| \leq 1/3$$

- By standard techniques (eg John's lemma), we can assume $\sum |\alpha_i| \leq L$ (where $\log L \leq \text{polylog}(m)$)

Goal: Upper bound $\text{spar}(g)$ in terms of $\text{rank}^{1/3}(g \circ \text{AND}_2)$

- We shall construct a function f (to be obtained from g) such that
 - Approximate degree of f is small
 - Degree of f is high

This contradicts known relations between degree and approximate degree (first proven by Nisan-Szegedy; constants were optimized later).

Goal: Upper bound $\text{spar}(g)$ in terms of $\text{rank}^{1/3}(g \circ \text{AND}_2)$

- We shall construct a function f (to be obtained from g) such that
 - Approximate degree of f is small
 - Degree of f is high

This contradicts known relations between degree and approximate degree (first proven by Nisan-Szegedy; constants were optimized later).

- We shall get a low-degree approximating polynomial from the inequality

$$\left| g(x \wedge y) - \sum_{i=1}^m \alpha_i a_i(x) \wedge b_i(y) \right| \leq 1/3$$

whereas the high sparsity of g will guarantee the original polynomial has high degree.

Goal: Upper bound $\text{spar}(g)$ in terms of $\text{rank}^{1/3}(g \circ \text{AND}_2)$

- Note that every function $f : \{0, 1\}^n \rightarrow \mathbb{R}$ can also be represented as a function $\tilde{f} : \{\pm 1\}^n \rightarrow \mathbb{R}$. It is easy to see that their degrees are the same.

Convention: For a function $f : \{0, 1\}^n \rightarrow \mathbb{R}$, we denote its equivalent function with the domain $\{\pm 1\}^n$ by $\tilde{f} : \{\pm 1\}^n \rightarrow \mathbb{R}$.

Goal: Upper bound $\text{spar}(g)$ in terms of $\text{rank}^{1/3}(g \circ \text{AND}_2)$

- Note that every function $f : \{0, 1\}^n \rightarrow \mathbb{R}$ can also be represented as a function $\tilde{f} : \{\pm 1\}^n \rightarrow \mathbb{R}$. It is easy to see that their degrees are the same.

Convention: For a function $f : \{0, 1\}^n \rightarrow \mathbb{R}$, we denote its equivalent function with the domain $\{\pm 1\}^n$ by $\tilde{f} : \{\pm 1\}^n \rightarrow \mathbb{R}$.

- For upper-bounding the approximate degree, we shall look at functions in the $\{\pm 1\}^n$ domain. For lower bounding the degree, we shall look at functions in the $\{0, 1\}^n$ domain. Since the degree is invariant under change of domain, this is ok.

- **Observation:** Given the expression

$$g(x \wedge y) \approx \sum_{i=1}^m \alpha_i a_i(x) \wedge b_i(y),$$

we can hit each a_i by a restriction+averaging operator (ρ, δ) , replace each b_i by a constant, and get an approximation for g_ρ

Restriction+averaging operators:

A **restriction+averaging operator** is denoted by a string $R \in \{0, 1, *, \Delta\}^n$. It converts a function $f : \{0, 1\}^n \rightarrow \mathbb{R}$ to a function $f : \{0, 1\}^{\text{free}(R)} \rightarrow \mathbb{R}$ where $\text{free}(R) = \{i \mid R_i = *\}$. The transformation is as follows:

- If $R_i = b \in \{0, 1\}$, fix the i -th coordinate to b .
- If $R_i = \Delta$, average over the i -th coordinate.
- If $R_i = *$, keep the i -th coordinate free.

We shall associate a partial assignment $\rho \in \{0, 1, *\}^n$ to each restriction+averaging operator R : ρ is obtained from R by changing all the Δ 's to 0's.

Goal: Upper bound $\text{spar}(g)$ in terms of $\text{rank}^{1/3}(g \circ \text{AND}_2)$

- We shall define a distribution $\mathcal{D}$ over *restriction+averaging operators* R such that for any $\tilde{a} : \{\pm 1\}^n \rightarrow [0, 1]$,

$$\mathbb{E}_{R \sim \mathcal{D}}[|\tilde{a}_R|^{\geq d}] \leq \varepsilon$$

where $|\tilde{f}|^{\geq d}$ denotes ℓ^1 Fourier weight above level d

Goal: Upper bound $\text{spar}(g)$ in terms of $\text{rank}^{1/3}(g \circ \text{AND}_2)$

- We shall define a distribution $\mathcal{D}$ over *restriction+averaging operators* R such that for any $\tilde{a} : \{\pm 1\}^n \rightarrow [0, 1]$,

$$\mathbb{E}_{R \sim \mathcal{D}}[|\tilde{a}_R|^{\geq d}] \leq \varepsilon$$

where $|\tilde{f}|^{\geq d}$ denotes ℓ^1 Fourier weight above level d

- ... and such that $\mathbb{E}_{R \sim \mathcal{D}}[f_\rho] \geq D$, where ρ is the partial assignment associated with R .

- The first property, along with the expression for approximate rank, will give that $\text{approx-deg}(g_\rho) \leq d$ for a typical $R \sim \mathcal{D}$

- The first property, along with the expression for approximate rank, will give that $\text{approx-deg}(g_\rho) \leq d$ for a typical $R \sim \mathcal{D}$
- The second property will give that $\text{deg}(g_\rho) \geq D$ for a typical $R \sim \mathcal{D}$.

- The first property, along with the expression for approximate rank, will give that $\text{approx-deg}(g_\rho) \leq d$ for a typical $R \sim \mathcal{D}$
- The second property will give that $\text{deg}(g_\rho) \geq D$ for a typical $R \sim \mathcal{D}$.
- If the gap between d and D is large enough, we are done.

Obtaining an approximation for f_ρ

- Recall the expression we got from the low-approximate rank assumption:

$$g(x \wedge y) \approx \sum_{i=1}^m \alpha_i a_i(x) b_i(y)$$

$$a_i, b_i : \{0, 1\}^n \rightarrow [0, 1]$$

$$\sum_i |\alpha_i| \leq L$$

Obtaining an approximation for f_ρ

- Recall the expression we got from the low-approximate rank assumption:

$$g(x \wedge y) \approx \sum_{i=1}^m \alpha_i a_i(x) b_i(y)$$

$$a_i, b_i : \{0, 1\}^n \rightarrow [0, 1]$$

$$\sum_i |\alpha_i| \leq L$$

- Claim:** Given a restriction+averaging operator R , with ρ its associated partial assignment, there exist constants $c_1, c_2, \dots, c_m \in [0, 1]$ such that

$$g_\rho(z) \approx \sum_{i=1}^m \alpha_i c_i(a_i)_R(z)$$

Obtaining an approximation for f_ρ

- **Claim:** Given a restriction+averaging operator R , with ρ its associated partial assignment, there exist constants $c_1, c_2, \dots, c_m \in [0, 1]$ such that

$$g_\rho(z) \approx \sum_{i=1}^m \alpha_i c_i(a_i)_R(z)$$

Obtaining an approximation for f_ρ

- **Claim:** Given a restriction+averaging operator R , with ρ its associated partial assignment, there exist constants $c_1, c_2, \dots, c_m \in [0, 1]$ such that

$$g_\rho(z) \approx \sum_{i=1}^m \alpha_i c_i(a_i)_R(z)$$

- I will show this only for the special case $R = \{\Delta, *, *, *, \dots, *\}$. General case should be clear from this.

Obtaining an approximation for f_ρ

- **Claim:** Given a restriction+averaging operator R , with ρ its associated partial assignment, there exist constants $c_1, c_2, \dots, c_m \in [0, 1]$ such that

$$g_\rho(z) \approx \sum_{i=1}^m \alpha_i c_i(a_i)_R(z)$$

- I will show this only for the special case $R = \{\Delta, *, *, *, \dots, *\}$. General case should be clear from this.
- So, $\rho = \{0, *, *, *, \dots, *\}$.

Obtaining an approximation for f_ρ

- I will show this only for the special case $R = \{\Delta, *, *, *, \dots, *\}$.
General case should be clear from this. So, $\rho = \{0, *, *, *, \dots, *\}$.

Obtaining an approximation for f_ρ

- I will show this only for the special case $R = \{\Delta, *, *, *, \dots, *\}$.
General case should be clear from this. So, $\rho = \{0, *, *, *, \dots, *\}$.
- Plugging $y_0 = (0, 1, 1, \dots, 1), x = (1, z_2, z_3, \dots, z_n)$, we get

$$g(0, z_2, z_3, \dots, z_n) \approx \sum_{i=1}^m \alpha_i a_i(1, z_2, \dots, z_n) b_i(y_0)$$

Plugging $y_0 = (0, 1, 1, \dots, 1), x = (0, z_2, z_3, \dots, z_n)$,

$$g(0, z_2, c_3, \dots, z_n) \approx \sum_{i=1}^m \alpha_i a_i(0, z_2, z_3, \dots, z_n) b_i(y_0)$$

Obtaining an approximation for f_ρ

- I will show this only for the special case $R = \{\Delta, *, *, *, \dots, *\}$.
General case should be clear from this. So, $\rho = \{0, *, *, *, \dots, *\}$.
- Plugging $y_0 = (0, 1, 1, \dots, 1), x = (1, z_2, z_3, \dots, z_n)$, we get

$$g(0, z_2, z_3, \dots, z_n) \approx \sum_{i=1}^m \alpha_i a_i(1, z_2, \dots, z_n) b_i(y_0)$$

Plugging $y_0 = (0, 1, 1, \dots, 1), x = (0, z_2, z_3, \dots, z_n)$,

$$g(0, z_2, c_3, \dots, z_n) \approx \sum_{i=1}^m \alpha_i a_i(0, z_2, z_3, \dots, z_n) b_i(y_0)$$

- Average these two expressions:

$$g(0, z_2, \dots, z_n) \approx \sum_{i=1}^m \alpha_i b_i(y_0) (a_i)_R(z)$$

Exhibiting a gap between degree and approximate degree

- **Conclusion:** If R is a restriction+averaging operator and ρ is the associated partial assignment, we get an approximation for g_ρ in terms of the $(a_i)_R$'s.

Exhibiting a gap between degree and approximate degree

- **Conclusion:** If R is a restriction+averaging operator and ρ is the associated partial assignment, we get an approximation for g_ρ in terms of the $(a_i)_R$'s.
- **Our next goal:** Using the fact that sparsity of g is high, we define a distribution $\mathcal{D}$ over restriction+averaging operators R such that:
 - For all $a : \{0, 1\}^n \rightarrow [0, 1]$, $E_{R \sim \mathcal{D}} \tilde{\deg}(a_R) \leq d$
 - $E_{(R \sim \mathcal{D})} [\deg(g_\rho)] \geq D$ where ρ is the partial assignment corresponding to R .