1.

$$abla f(x) =
abla (rac{1}{2}x^TAx + b^Tx) = Ax + b$$

(b)

$$\frac{\partial g(h(x))}{\partial x_i} = \frac{\partial g(h(x))}{\partial h(x)} \frac{\partial h(x)}{\partial x_i} = g'(h(x)) \frac{\partial h(x)}{\partial x_i}$$
$$\nabla f(x) = \nabla g(h(x)) = g'(h(x)) \nabla h(x)$$

(c)

$$egin{aligned}
abla^2 f(x) &= \left[rac{\partial
abla f(x)}{\partial x_1} & rac{\partial
abla f(x)}{\partial x_2} & \ldots & rac{\partial
abla f(x)}{\partial x_n}
ight] \ &= \left[rac{\partial
abla (Ax+b)}{\partial x_1} & rac{\partial
abla (Ax+b)}{\partial x_2} & \ldots & rac{\partial
abla f(x)}{\partial x_n}
ight] \ &= \left[egin{aligned} A_{11} & A_{12} & \ldots & A_{1n} \ A_{21} & A_{22} & \ldots & A_{2n} \ dots & dots & \ddots & dots \ A_{n1} & A_{n2} & \ldots & A_{nn} \end{array}
ight] = A \end{aligned}$$

(d)

$$\nabla f(x) = \nabla g(a^T x) = g'(a^T x)\nabla(a^T x) = g'(a^T x)a$$

$$\frac{\partial^2 g(h(x))}{\partial x_i \partial x_j} = \frac{\partial^2 g(h(x))}{\partial (h(x))^2} \frac{\partial h(x)}{\partial x_i} \frac{\partial h(x)}{\partial x_j} = g''(h(x)) \frac{\partial h(x)}{\partial x_i} \frac{\partial h(x)}{\partial x_j}$$

$$\frac{\partial^2 g(a^T x)}{\partial x_i \partial x_j} = g''(a^T x) \frac{\partial (a^T x)}{\partial x_i} \frac{\partial (a^T x)}{\partial x_j} = g''(a^T x)a_i a_j$$

$$\nabla^2 f(x) = \nabla^2 g(a^T x) = g''(a^T x) \begin{bmatrix} a_1 a_1 & a_1 a_2 & \dots & a_1 a_n \\ a_2 a_1 & a_2 a_2 & \dots & a_2 a_n \\ \vdots & \vdots & \ddots & \vdots \\ a_n a_1 & a_n a_2 & \dots & a_n a_n \end{bmatrix} = g''(a^T x)aa^T$$

2.

$$A^T=(zz^T)^T=zz^T=A$$
 $x^TAx=x^Tzz^Tx=x^Tz(x^Tz)^T=(x^Tz)^2\geq 0$

(b)

$$Ax=zz^Tx=z(z^Tx)=0$$
 $N(A)=\{x\in\mathbb{R}^n:z^Tx=0\}$ $R(A)=R(zz^T)\leq \min\{R(z),R(z^T)\}=1$

$$R(A) = 1$$

(c)

$$(BAB^T)^T = BA^TB^T = BAB^T$$
 $x^TBAB^Tx = (x^TB)A(x^TB)^T \geq 0$

3.

(a)

$$AT = T\Lambda$$

$$A\left[t^{(1)} \quad t^{(2)} \quad \dots \quad t^{(n)}\right] = \left[t^{(1)} \quad t^{(2)} \quad \dots \quad t^{(n)}\right] \begin{bmatrix} \lambda_1 & 0 & \dots & 0 \\ 0 & \lambda_2 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & \lambda_n \end{bmatrix}$$

$$\left[At^{(1)} \quad At^{(2)} \quad \dots \quad At^{(n)}\right] = \left[\lambda_1 t^{(1)} \quad \lambda_2 t^{(2)} \quad \dots \quad \lambda_n t^{(n)}\right]$$

$$At^{(i)} = \lambda_i t^{(i)}$$

 $A = T\Lambda T^{-1}$

(b)

$$A = U\Lambda U^T$$
 $AU = U\Lambda U^T U = U\Lambda$

(c)

$$At^{(i)} = \lambda_i t^{(i)} \ (t^{(i)})^T At^{(i)} = \lambda_i \|t^{(i)}\|_2 = \lambda_i \geq 0$$