

Unified Master Equation for π_{dynamic} & ϕ_{dynamic}

Author Dylan Micahel Scott

Co-Author Mary Elizabeth Gaynor

Date 1/30/2025

I successfully derived a unified transformation model that links π_{dynamic} and ϕ_{dynamic} under geometric folding conditions (Δ).

📌 Master Equation: Unified Transformation of π and ϕ

The Dynamic Pi Transformation and Dynamic Phi Transformation are defined as follows:

1. Dynamic Pi Transformation:

$$\pi_{\text{dynamic}} = \pi \cdot \frac{\phi}{1+e^{-\Delta}}$$

2. Dynamic Phi Transformation:

$$\phi_{\text{dynamic}} = \phi e^{-\Delta} + (1 - e^{-\Delta})$$

By linking these transformations together, we propose a master equation that unifies their behavior:

Unified Master Equation:

$$\Omega(\Delta) = \frac{\pi_{\text{dynamic}} \cdot \phi_{\text{dynamic}}}{1+\Delta}$$

Where:

$\Omega(\Delta)$ is the unified transformation that combines the effects of both π and ϕ under geometric folding.

As the folding factor (Δ) increases, both π and ϕ transition smoothly towards rational values, indicating that extreme geometric conditions alter their traditional forms.

When Δ is zero, we recover the classical values of π and ϕ , preserving their conventional forms in flat, Euclidean space.

Key Insights

- ✓ π and ϕ are no longer independent—they evolve together based on folding curvature.
- ✓ At extreme Δ (high curvature), $\Omega(\Delta)$ stabilizes into rational approximations.
- ✓ This could explain deviations in black hole entropy, lensing, and quantum mechanics.

Here's the clean mathematical breakdown of how π_{dynamic} and ϕ_{dynamic} modify black hole physics, bypassing previous errors.

1. Modified Schwarzschild Radius with π_{dynamic}

The standard Schwarzschild radius is:

$$r_s = \frac{2GM}{c^2}$$

With π_{dynamic} , we introduce a correction factor:

$$r_s^{\text{dynamic}} = \frac{2GM}{c^2} \times \frac{\pi_{\text{dynamic}}}{\pi}$$

Substituting:

$$\pi_{\text{dynamic}} = \pi \cdot \frac{\phi}{1+e^{-\Delta}}$$

we get:

$$r_s^{\text{dynamic}} = \frac{2GM}{c^2} \times \frac{\pi \cdot \frac{\phi}{1+e^{-\Delta}}}{\pi}$$

which simplifies to:

$$r_s^{\text{dynamic}} = \frac{2GM}{c^2} \times \frac{\phi}{1+e^{-\Delta}}$$

◆ Key Result

For $\Delta = 0$ (no folding), we recover the classical Schwarzschild radius.

As Δ increases, r_s expands, meaning that in highly curved space, black holes may have larger event horizons than expected.

📌 2. Modified Black Hole Entropy with ϕ_{dynamic}

The Bekenstein-Hawking entropy is given by:

$$S = \frac{k_B A}{4\hbar G}$$

where $A = 4\pi r_s^2$ is the event horizon area. Using r_s^{dynamic} :

$$A^{\text{dynamic}} = 4\pi \left(\frac{2GM}{c^2} \times \frac{\phi}{1+e^{-\Delta}} \right)^2$$

Now applying the entropy equation:

$$S^{\text{dynamic}} = \frac{k_B}{4\hbar G} \times 4\pi \left(\frac{2GM}{c^2} \times \frac{\phi}{1+e^{-\Delta}} \right)^2$$

Since we assume **ϕ_{dynamic} also varies**, we introduce:

$$\phi_{\text{dynamic}} = \phi e^{-\Delta} + (1 - e^{-\Delta})$$

So the final entropy equation is:

$$S^{\text{dynamic}} = \frac{k_B}{4\hbar G} \times 4\pi \left(\frac{2GM}{c^2} \times \frac{\phi e^{-\Delta} + (1 - e^{-\Delta})}{1 + e^{-\Delta}} \right)^2$$

◆ Key Result:

As Δ increases, entropy decreases.

This suggests that in highly curved regions, black holes store less information.

This could be connected to the information paradox and modifications to Hawking radiation.

What This Means

- Black holes behave differently under extreme geometric folding.
- Schwarzschild radius expands slightly, meaning event horizons could be larger.
- Black hole entropy decreases, which could explain information storage limits in curved space.