■■■■ ■■■■■ · GOVERNMENT OF INDIA
Family Planning Division · Ministry of Health and Family Welfare

FAMILY PLANNING CLINICAL HANDBOOK FOR FRONTLINE WORKERS

Mission Parivar Vikas Guidelines on Postpartum Contraceptive Choices

DOCUMENT ID: R7-NHM-2026 **EDITION:** Official Field Standard 4.2 **TARGET:** Frontline ASHA Workers

1. Postpartum Contraception Timing & Options

• Postpartum Intrauterine Contraceptive Device (PPIUCD): Within 48 hours of institutional delivery.

• Postpartum Sterilization (PPS): Within 7 days of delivery or after 6 weeks.

• Progestin-only Pills (POP) / Lactational Amenorrhea Method (LAM): For breastfeeding mothers up to 6 months.

• Condoms (Nirodh) and Emergency Contraceptive Pills (ECP) distribution at village sub-centre.

2. Counseling Matrix

• Spacing of at least 3 years between successive pregnancies protects maternal health and prevents stunting in offspring.

• Dispense Chhaya (Centchroman non-hormonal oral pill) and Antara (Injectable MPA) under national mission guidelines.

Issuing Authority:
National Health Mission Division
Nirman Bhawan, New Delhi

Field Protocol Verification:
Primary Health Centre / Block MO
Mandatory compliance across sub-centres

ASHA Frontline Desk:
Document all actions in MCP Card
Follow 108 Emergency Protocol