[redacted] · GOVERNMENT OF INDIA
Immunization Division · Ministry of Health & Family Welfare · Government of India

UNIVERSAL IMMUNIZATION PROGRAMME (UIP) HANDBOOK

National Immunization Schedule & Cold Chain Protocol for Healthcare Workers

DOCUMENT ID: R3-NHM-2026 **EDITION:** Official Field Standard 4.2 **TARGET:** Frontline ASHA Workers

1. Birth & Infancy Vaccination Schedule

• At Birth: BCG (0.1 ml ID), Hepatitis B Birth dose (0.5 ml IM within 24 hrs), OPV Zero dose (2 drops oral).

• 6 Weeks: Pentavalent-1 (DPT+HepB+Hib), OPV-1, Fractional IPV-1, Rotavirus-1, PCV-1.

• 10 Weeks: Pentavalent-2, OPV-2, Rotavirus-2.

• 14 Weeks: Pentavalent-3, OPV-3, Fractional IPV-2, Rotavirus-3, PCV-2.

• 9-12 Months: MR-1 (Measles-Rubella), PCV Booster, Japanese Encephalitis-1 (endemic districts), Vitamin A (1 lakh IU).

2. Toddler & Pregnancy Immunization

• 16-24 Months: MR-2, DPT Booster-1, OPV Booster, JE-2, Vitamin A 2nd dose.

• 5-6 Years: DPT Booster-2.

• 10 & 16 Years: Td (Tetanus & adult Diphtheria).

• Pregnant Women: Td-1 early in pregnancy + Td-2 after 4 weeks.

3. Adverse Events Following Immunization (AEFI) Management

• Maintain child observation for 30 minutes post-injection at the Village Health & Nutrition Day (VHND) site.

• Paracetamol syrup (10-15 mg/kg) for mild fever and fussiness; reassure family.

• Immediate escalation to MO for severe allergic reaction / anaphylaxis (keep Adrenaline 1:1000 accessible).

Issuing Authority:
National Health Mission Division
Nirman Bhawan, New Delhi

Field Protocol Verification:
Primary Health Centre / Block MO
Mandatory compliance across sub-centres

ASHA Frontline Desk:
Document all actions in MCP Card
Follow 108 Emergency Protocol