

antidepressant 역시 neuroplasticity를 증진시켜 우울증을 치료한다고 하는데, 이때의 plasticity가 비특이적으로 가소성 potential을 올리는 것인지, 아니면 specific하게 abnormal한 부분만 rewiring하는 것인지에 대해 각각을 지지하는 연구들을 찾아서 비교 설명해줘 최대한 상세히 설명해줘

Aristotle Instant

Synthesized evidence comparing general plasticity enhancement and specific circuit normalization by antidepressants

Neuroplasticity and depression: Rewiring ...

Neuroplasticity As A Common...

Frontiers | From Serotonin to Neuroplasti...

[PDF] How Antidepressants and Cognitiv...

[PDF] NEUROPLASTICITY IN DEPRESSI...

Neuroplasticity and the next wave of anti...

Spandidos Publications: Experimental an...

The Role of BDNF on Neural Plasticity in ...

Frontiers | Neuroplasticity and the next w...

The Role of Neural Plasticity in Depressio...

Adult neuroplasticity: A new "cure" for ma...

Rapid neuroplasticity changes and respo...

Neuroplasticity and Mechanisms of Actio...

5. Monoamine Theory Of...

Neuroplasticity and Mechanisms of Actio...

The user is asking whether antidepressant-induced neuroplasticity works through:

1. Non-specific enhancement of overall plasticity potential, or
2. Specific rewiring of abnormal circuits

I need to synthesize the information I found to compare these two views, gathering supporting evidence for each.

Here's a breakdown of the evidence:

Evidence for Non-specific/General Plasticity Enhancement:

- BDNF is upregulated broadly across many brain regions.
- General neurogenesis occurs in the hippocampus.
- The cAMP-CREB pathway is activated, which is a general mechanism.
- Learning and memory-like processes are involved.
- LTP-like processes are observed.
- The timing of these changes aligns with how antidepressants typically work.

Evidence for Specific/Targeted Rewiring:

- Effects are region-specific; for instance, the hippocampus and amygdala show opposite changes.
- BDNF effects themselves are region-specific.
- Specific circuits, like the default mode network (DMN) and limbic-prefrontal circuits, appear to be normalized.
- Different antidepressants seem to have different regional impacts.
- Specific receptor mechanisms, such as 5-HT1A, have cell-type specific effects.
- There's evidence for targeting specific biotypes, like anhedonia, which involves striatal circuits.

I'll structure this comprehensively in Korean as requested.

Antidepressant-Induced Neuroplasticity: Non-Specific Enhancement vs. Specific Rewiring

The question of whether antidepressants enhance neuroplasticity in a generalized, non-specific manner or whether they selectively target and rewire abnormal circuits is a central debate in depression neurobiology. The evidence suggests a nuanced picture where both mechanisms may operate, though the relative contribution of each remains under investigation.

The Non-Specific Plasticity Enhancement Hypothesis

This view holds that antidepressants broadly increase the brain's capacity for plastic change, essentially "reopening" a window of heightened neuroplasticity similar to critical periods in development. The therapeutic effect then depends on environmental inputs and learning experiences to guide adaptive rewiring.

Supporting Evidence

1. Generalized Upregulation of Neurotrophic Factors

Antidepressants consistently increase brain-derived neurotrophic factor (BDNF) levels across multiple brain regions, not just those implicated in depression. The neuroplasticity theory of depression is supported by evidence that antidepressants elevate neurotrophic factor concentrations broadly and improve neuroplasticity in both the hippocampus and prefrontal cortex ^{[1][2][7][12][14]}. This upregulation follows a common intracellular cascade: monoamine receptor stimulation activates adenylate cyclase, which catalyzes ATP to cAMP, subsequently activating CREB through protein kinase A ^{[7][16]}. This pathway operates as a general mechanism rather than targeting specific dysfunctional circuits.

2. Hippocampal Neurogenesis as a General Phenomenon

Virtually all effective antidepressant treatments, including SSRIs, tricyclics, vagal stimulation, and deep brain stimulation, positively influence hippocampal neurogenesis ^[5]. This consistent finding across mechanistically diverse treatments suggests that enhancing general neuroplastic capacity may be a final common pathway. ECT, the most potent inducer of neuroplasticity, triggers multiple mechanisms including synaptogenesis, neurogenesis, dendrogenesis, angiogenesis, and gliogenesis simultaneously ^{[1][15]}.

3. The "Permissive" Model

Research indicates that antidepressant-induced neuroplasticity may function permissively, allowing learning and environmental inputs to reshape neural circuits. One review proposed that CBT may help "imprint healthy cognitions and behaviors on the neuroplasticity changes induced by medication" ^[2]. This suggests antidepressants create a general state of enhanced plasticity rather than directing specific circuit changes.

4. Critical Period Reopening

Some research frames antidepressant action as reopening juvenile-like critical periods of plasticity. SSRIs have been shown to reactivate visual cortex plasticity in adult animals, demonstrating effects extending well beyond mood-related circuits ^[5]. This broad enhancement of plasticity potential supports the non-specific hypothesis.

5. Potential for Maladaptive Plasticity

A critical piece of evidence supporting non-specificity is that enhanced plasticity can sometimes be harmful. BDNF promotes neuroplasticity not only in the hippocampus and PFC (where it is beneficial) but also in the amygdala, ventral tegmental area (VTA), and nucleus accumbens (NAc), where it may provoke depressive-like behaviors or exacerbate symptoms ^{[5][7][16][12][9]}. Stress increases BDNF in the basolateral amygdala while decreasing it in the hippocampus, and antidepressants do not always selectively reverse this pattern ^[5]. This bidirectional effect suggests the plasticity enhancement is not inherently targeted to pathological circuits.

The Specific Rewiring Hypothesis

This view proposes that antidepressants selectively reverse maladaptive plasticity in specific circuits while potentially sparing or differentially affecting healthy neural networks.

Supporting Evidence

1. Region-Specific Effects of SSRIs vs. SNRIs

A comparative study found that SSRIs (citalopram, escitalopram) produced decreased limbic activity and normalization of frontoparietal network activity during cognitive tasks, while serotonin-norepinephrine reuptake inhibitors showed limited impact on the same limbic regions ^[1]. This differential effect suggests mechanistic specificity rather than global plasticity enhancement.

2. Default Mode Network Targeting

The default mode network (DMN), which is hyperactive in depression and associated with negative rumination, appears to be specifically targeted by certain antidepressants. Ketamine treatment specifically reduced DMN metabolic activity and diminished DMN connectivity within the cingulate and prefrontal cortices ^{[3][5]}. This targeted effect could help "give depressed patients the needed 'push' to move them out of a current negative ruminative state" ^[5].

3. BDNF Region-Specificity

While BDNF is generally upregulated, its effects are demonstrably region-specific. One review noted that "BDNF's mechanism of action is controversial and it was observed that its neuroplasticity effects are region-specific and interact with other genetic and environmental factors" ^[1]. The downstream consequences of BDNF signaling depend heavily on local circuit properties.

4. Reversal of Stress-Induced Changes

Antidepressants appear to specifically reverse the neuroanatomical changes caused by chronic stress and depression. Pathological stress causes neuronal atrophy in the hippocampus and prefrontal cortex while simultaneously increasing dendritic arborization in the amygdala ^{[2][5]}. Antidepressant treatment reverses these changes in a pattern-specific manner: restoring hippocampal and prefrontal volume while normalizing (not further increasing) amygdala structure ^{[1][2]}.

5. Circuit-Specific Functional Connectivity Changes

Neuroimaging studies demonstrate that antidepressant therapy produces specific, not generalized, changes in functional connectivity. After treatment, patients show increased functional connectivity in neocortical regions (prefrontal cortex) and subcortical regions (hippocampus, parahippocampus, insula) that was correlated with clinical improvement ^[1]. Specifically, there is decreased activation in limbic areas in response to negative stimuli and increased activity in dorsolateral prefrontal cortex in response to positive stimuli, indicating selective reversal of the depression-associated pattern ^{[1][18]}.

6. Amygdala-Specific Normalization

After antidepressant therapy, there is decreased activation of the amygdala during emotional processing, specifically reducing negative emotional processing while normalizing activity in regions involved in resting state and emotional response ^{[1][15]}. This targeted effect on emotion-processing circuits supports specific rather than global rewiring.

7. Biotype-Specific Treatment Responses

Research suggests that different depression biotypes may benefit from medications targeting neuroplasticity in specific circuits. For example, anhedonia is mediated through hypoactivation of striatal pathways, and patients exhibiting this symptom may specifically benefit from medication targeting neuroplasticity in striatal dopaminergic circuits (such as pramipexole) ^[1]. This circuit-specific approach implies that plasticity effects are not uniform.

Comparative Analysis

Aspect	Non-Specific Enhancement	Specific Rewiring
BDNF mechanism	Global upregulation across brain regions ^[2] ^[5]	Region-specific downstream effects depending on local circuit properties ^[1] ^[16]
Hippocampal neurogenesis	Universal feature of all effective antidepressants ^[5]	May be necessary but not sufficient; functional integration matters ^[13] ^[19]
DMN effects	Part of broader cortical plasticity	Specifically targeted by ketamine, possibly SSRIs ^[3] ^[5]
Amygdala changes	Can be enhanced or suppressed ^[5]	Specifically normalized after treatment ^[1] ^[15] ^[4] ^[10]
Time course	Gradual, parallels neuroplasticity development ^[2]	Some effects rapid (ketamine), suggesting targeted mechanisms ^[3] ^[6]
Treatment implications	Environmental enrichment and therapy enhance outcomes ^[2]	Circuit-specific interventions may be more effective ^[1]

Integrative Framework: The "Priming and Guiding" Model

The most current understanding suggests both mechanisms operate in concert. Several sources propose an integrative model where antidepressants enhance overall plasticity potential (non-specific), but the direction and consolidation of plastic changes is guided by several factors:

1. Concurrent Psychotherapy

The observation that antidepressant-CBT combinations are generally more effective than either alone ^[2] suggests that pharmacologically enhanced plasticity requires guided learning to produce optimal circuit-specific changes. Antidepressants may "prime" the brain for change while therapy provides the directed input.

2. Activity-Dependent Consolidation

Neuroplasticity induced by antidepressants may be activity-dependent, meaning that circuits that are actively engaged during treatment are preferentially modified. This explains why enhanced plasticity does not randomly rewire the brain but instead strengthens relevant circuits.

3. Homeostatic Regulation

The brain possesses homeostatic mechanisms that may help channel non-specific plasticity toward restoring normal function. Depressed circuits may be more "responsive" to plasticity-promoting signals because they are further from homeostatic set points.

Critical Unresolved Questions

1. The Amygdala Paradox

Both stress and some antidepressant mechanisms can enhance amygdala plasticity, yet the clinical outcome differs ^[5]^[13]. Understanding what determines whether enhanced amygdala plasticity is adaptive or maladaptive remains unclear.

2. Individual Variation

The neuroplasticity effects of antidepressants show substantial individual variation that may depend on genetic factors, stress history, and baseline circuit states ^{[1][16]}. This variability complicates the specific vs. non-specific distinction.

3. Acute vs. Chronic Treatment

Acute and chronic antidepressant treatment may have different specificity profiles. Acute treatment with fast-acting antidepressants like ketamine activates signaling pathways modulating glutamate release through GABAergic neurons and astrocytes, while chronic treatment stimulates cAMP-CREB transcription factors regulating genes involved in neuroplasticity, cell survival, and cognition ^[8]. The temporal dynamics of specificity require further investigation.

4. The Role of Inflammation

Immune and stressor-induced changes in neuroplasticity, involving adult neurogenesis, synaptogenesis, dendritic remodeling, and trophic signaling, are posited to underlie the "faulty wiring" in depression ^{[3][5][14][10]}. Whether anti-inflammatory mechanisms of some antidepressants contribute to specific or non-specific plasticity effects is unclear.

Conclusions

The evidence does not clearly favor either pure non-specific enhancement or purely specific rewiring. Instead, the data suggest:

- 1 . Antidepressants engage broad neuroplasticity mechanisms (BDNF upregulation, neurogenesis, synaptogenesis) that could theoretically affect any neural circuit ^{[1][2][5][8][17][12][14]}.
- 1 . However, the functional consequences of this enhanced plasticity are shaped by circuit-specific factors, including baseline dysfunction severity, local receptor distributions, activity patterns, and concurrent environmental inputs ^{[1][7][16][11]}.
- 1 . The therapeutic benefit likely requires both: sufficient plasticity potential (non-specific enhancement) AND appropriate guidance of plastic changes toward adaptive circuit configurations (which may be partially intrinsic through homeostatic mechanisms and partially dependent on therapy and environment).
- 1 . Future research directions include identifying biomarkers that predict which patients will benefit from plasticity-enhancing treatments and developing strategies to more precisely direct enhanced plasticity toward therapeutic circuit changes ^[19].

This integrative view has important clinical implications: maximizing antidepressant efficacy may require not only pharmacological enhancement of plasticity but also concurrent interventions (psychotherapy, environmental enrichment, behavioral activation) that help guide plastic changes in therapeutically beneficial directions ^{[2][5][12][9]}.

Citations

- 1 . [Neuroplasticity and depression: Rewiring the brain's networks ...](https://pmc.ncbi.nlm.nih.gov/articles/PMC8383338/)
<https://pmc.ncbi.nlm.nih.gov/articles/PMC8383338/>
- 2 . [Neuroplasticity As A Common...](https://pmc.ncbi.nlm.nih.gov/articles/PMC3025168/)
<https://pmc.ncbi.nlm.nih.gov/articles/PMC3025168/>
- 3 . [Frontiers | Neuroplasticity and the next wave of antidepressant strategies](https://www.frontiersin.org/journals/cellular-neuroscience/articles/10.3389/fncel.2013.00218/full)
<https://www.frontiersin.org/journals/cellular-neuroscience/articles/10.3389/fncel.2013.00218/full>
- 4 . [Adult neuroplasticity: A new “cure” for major depression? - PMC](https://pmc.ncbi.nlm.nih.gov/articles/PMC6488487/)
<https://pmc.ncbi.nlm.nih.gov/articles/PMC6488487/>
- 5 . [Neuroplasticity and the next wave of antidepressant strategies](https://pmc.ncbi.nlm.nih.gov/articles/PMC3834236/)
<https://pmc.ncbi.nlm.nih.gov/articles/PMC3834236/>
- 6 . [Rapid neuroplasticity changes and response to intravenous ketamine](#)

- <https://www.nature.com/articles/s41398-023-02451-0>
7. [Frontiers | From Serotonin to Neuroplasticity: Evolvement of Theories for Major Depressive Disorder](https://www.frontiersin.org/journals/cellular-neuroscience/articles/10.3389/fncel.2017.00305/full)
<https://www.frontiersin.org/journals/cellular-neuroscience/articles/10.3389/fncel.2017.00305/full>
 8. [Neuroplasticity and Mechanisms of Action of Acute and ...](https://pmc.ncbi.nlm.nih.gov/articles/PMC11727250/)
<https://pmc.ncbi.nlm.nih.gov/articles/PMC11727250/>
 9. [Neuroplasticity and depression: Rewiring the brain's networks](https://www.spandidos-publications.com/10.3892/etm.2021.10565)
<https://www.spandidos-publications.com/10.3892/etm.2021.10565>
 0. [\[PDF\] NEUROPLASTICITY IN DEPRESSION: A NARRATIVE REVIEW ...](https://www.psychiatria-danubina.com/UserDocsImages/pdf/dnb_vol34_no3/dnb_vol34_no3_390.pdf)
https://www.psychiatria-danubina.com/UserDocsImages/pdf/dnb_vol34_no3/dnb_vol34_no3_390.pdf
 1. [The Role of Neural Plasticity in Depression: From Hippocampus to Prefrontal Cortex](https://onlinelibrary.wiley.com/doi/10.1155/2017/6871089)
<https://onlinelibrary.wiley.com/doi/10.1155/2017/6871089>
 2. [Source 12](https://www.nature.com/articles/s41398-023-02451-0)
<https://www.nature.com/articles/s41398-023-02451-0>
 3. [Source 13](https://pmc.ncbi.nlm.nih.gov/articles/PMC11727250/)
<https://pmc.ncbi.nlm.nih.gov/articles/PMC11727250/>
 4. [Source 14](https://pmc.ncbi.nlm.nih.gov/articles/PMC3649690/)
<https://pmc.ncbi.nlm.nih.gov/articles/PMC3649690/>
 5. [Source 15](https://www.mdpi.com/2227-9059/12/12/2744)
<https://www.mdpi.com/2227-9059/12/12/2744>
 6. Source 16
 7. Source 17
 8. Source 18
 9. Source 19



pmc.ncbi.nlm.nih.gov
pmc.ncbi.nlm.nih.gov

Neuroplasticity and depression:
Rewiring the brain's networks ...



pmc.ncbi.nlm.nih.gov
pmc.ncbi.nlm.nih.gov

Neuroplasticity As A Common...



www.frontiersin.org
www.frontiersin.org

Frontiers | From Serotonin to
Neuroplasticity: Evolvement of
Theories for Major Depressive
Disorder



digitalcommons.pcom.edu
digitalcommons.pcom.edu

[PDF] How Antidepressants and
Cognitive Behavior Therapy (CBT)
can ...



www.psychiatria-danubina.com
www.psychiatria-danubina.com

[PDF] NEUROPLASTICITY IN
DEPRESSION: A NARRATIVE
REVIEW ...



pmc.ncbi.nlm.nih.gov
pmc.ncbi.nlm.nih.gov

Neuroplasticity and the next wave of
antidepressant strategies



www.spandidos-publications.com
www.spandidos-publications.com

Spandidos Publications:
Experimental and Therapeutic
Medicine



www.frontiersin.org
www.frontiersin.org

The Role of BDNF on Neural
Plasticity in Depression



www.frontiersin.org
www.frontiersin.org

Frontiers | Neuroplasticity and the
next wave of antidepressant
strategies



onlinelibrary.wiley.com
onlinelibrary.wiley.com

The Role of Neural Plasticity in
Depression: From Hippocampus to
Prefrontal Cortex



pmc.ncbi.nlm.nih.gov
pmc.ncbi.nlm.nih.gov

Adult neuroplasticity: A new "cure"
for major depression? - PubMed
Central



www.nature.com
www.nature.com

Rapid neuroplasticity changes and
response to intravenous ketamine