A randomized trial of entropy-guided therapy

Robin Researcher, MD; Kim Analyst, PhD; Lee Author, MSc

ABSTRACT

Brain entropy describes the diversity of neural states visited over time. We measured signal complexity with established estimators across sessions. Each participant completed two scanning visits separated by one week. The analysis controlled for head motion and physiological noise. Higher entropy was associated with broader subjective experience.

INTRODUCTION

We measured signal complexity with established estimators across sessions. Each participant completed two scanning visits separated by one week. The analysis controlled for head motion and physiological noise. Higher entropy was associated with broader subjective experience. These observations agree with prior reports from independent groups. Alternative explanations based on arousal alone were considered. The effect remained after correction for multiple comparisons. Brain entropy describes the diversity of neural states visited over time. We measured signal complexity with established estimators across sessions. Each participant completed two scanning visits separated by one week.

MATERIALS AND METHODS

Participants

Each participant completed two scanning visits separated by one week. The analysis controlled for head motion and physiological noise. Higher entropy was associated with broader subjective experience. These observations agree with prior reports from independent groups. Alternative explanations based on arousal alone were considered. The effect remained after correction for multiple comparisons. Brain entropy describes the diversity of neural states visited over time. We measured signal complexity with established estimators across sessions.

Statistical analysis

The analysis controlled for head motion and physiological noise. Higher entropy was associated with broader subjective experience. These observations agree with prior reports from independent groups. Alternative explanations based on arousal alone were considered. The effect remained after correction for multiple comparisons. Brain entropy describes the diversity of neural states visited over time. We measured signal complexity with established estimators across sessions. Each participant completed two scanning visits separated by one week.

RESULTS

Higher entropy was associated with broader subjective experience. These observations agree with prior reports from independent groups. Alternative explanations based on arousal alone were considered. The effect remained after correction for multiple comparisons. Brain entropy describes the diversity of neural states visited over time. We measured signal complexity with established estimators across sessions. Each participant completed two scanning visits separated by one week. The analysis controlled for head motion and physiological noise. Higher entropy was associated with broader subjective experience. These observations agree with prior reports from independent groups. Alternative explanations based on arousal alone were considered. The effect remained after correction for multiple comparisons.

These observations agree with prior reports from independent groups. Alternative explanations based on arousal alone were considered. The effect remained after correction for multiple comparisons. Brain entropy describes the diversity of neural states visited over time. We measured signal complexity with established estimators across sessions. Each participant completed two scanning visits separated by one week. The analysis controlled for head motion and physiological noise. Higher entropy was associated with broader subjective experience. These observations agree with prior reports from independent groups.

DISCUSSION

Alternative explanations based on arousal alone were considered. The effect remained after correction for multiple comparisons. Brain entropy describes the diversity of neural states visited over time. We measured signal complexity with established estimators across sessions. Each participant completed two scanning visits separated by one week. The analysis controlled for head motion and physiological noise. Higher entropy was associated with broader subjective experience. These observations agree with prior reports from independent groups. Alternative explanations based on arousal alone were considered. The effect remained after correction for multiple comparisons. Brain entropy describes the diversity of neural states visited over time. We measured signal complexity with established estimators across sessions.

Limitations

The effect remained after correction for multiple comparisons. Brain entropy describes the diversity of neural states visited over time. We measured signal complexity with established estimators across sessions. Each participant completed two scanning visits separated by one week. The analysis controlled for head motion and physiological noise.

CONCLUSIONS

Brain entropy describes the diversity of neural states visited over time. We measured signal complexity with established estimators across sessions. Each participant completed two scanning visits separated by one week. The analysis controlled for head motion and physiological noise.

CONFLICTS OF INTEREST

None declared.

REFERENCES

1. Author A, Writer B. Entropy in clinical populations. Lancet. 1991;12:1-9.
2. Author A, Writer B. Entropy in clinical populations. Lancet. 1992;12:1-9.
3. Author A, Writer B. Entropy in clinical populations. Lancet. 1993;12:1-9.
4. Author A, Writer B. Entropy in clinical populations. Lancet. 1994;12:1-9.
5. Author A, Writer B. Entropy in clinical populations. Lancet. 1995;12:1-9.
6. Author A, Writer B. Entropy in clinical populations. Lancet. 1996;12:1-9.
7. Author A, Writer B. Entropy in clinical populations. Lancet. 1997;12:1-9.
8. Author A, Writer B. Entropy in clinical populations. Lancet. 1998;12:1-9.
9. Author A, Writer B. Entropy in clinical populations. Lancet. 1999;12:1-9.
10. Author A, Writer B. Entropy in clinical populations. Lancet. 2000;12:1-9.
11. Author A, Writer B. Entropy in clinical populations. Lancet. 2001;12:1-9.
12. Author A, Writer B. Entropy in clinical populations. Lancet. 2002;12:1-9.
13. Author A, Writer B. Entropy in clinical populations. Lancet. 2003;12:1-9.
14. Author A, Writer B. Entropy in clinical populations. Lancet. 2004;12:1-9.
15. Author A, Writer B. Entropy in clinical populations. Lancet. 2005;12:1-9.