




# Talha SEECS

## IceWatch FYP Report.docx

-  My Files
-  My Files
-  University

---





### Document Details

**Submission ID****trn:oid::17268:93659653****Submission Date****Apr 30, 2025, 2:31 PM GMT+5:30****Download Date****Apr 30, 2025, 2:34 PM GMT+5:30****File Name****IceWatch FYP Report.docx****File Size****5.3 MB****71 Pages****10,522 Words****64,884 Characters**




# 11% Overall Similarity

The combined total of all matches, including overlapping sources, for each database.

## Match Groups

-  **90** Not Cited or Quoted 10%  
Matches with neither in-text citation nor quotation marks
-  **5** Missing Quotations 0%  
Matches that are still very similar to source material
-  **6** Missing Citation 1%  
Matches that have quotation marks, but no in-text citation
-  **0** Cited and Quoted 0%  
Matches with in-text citation present, but no quotation marks

## Top Sources

- 7%  Internet sources
- 6%  Publications
- 8%  Submitted works (Student Papers)

## Integrity Flags

### 0 Integrity Flags for Review

No suspicious text manipulations found.

Our system's algorithms look deeply at a document for any inconsistencies that would set it apart from a normal submission. If we notice something strange, we flag it for you to review.

A Flag is not necessarily an indicator of a problem. However, we'd recommend you focus your attention there for further review.

## Match Groups

- 90 Not Cited or Quoted 10%**  
Matches with neither in-text citation nor quotation marks
- 5 Missing Quotations 0%**  
Matches that are still very similar to source material
- 6 Missing Citation 1%**  
Matches that have quotation marks, but no in-text citation
- 0 Cited and Quoted 0%**  
Matches with in-text citation present, but no quotation marks

## Top Sources

- 7% Internet sources
- 6% Publications
- 8% Submitted works (Student Papers)

## Top Sources

The sources with the highest number of matches within the submission. Overlapping sources will not be displayed.

1	Internet	esdd.copernicus.org	1%
2	Submitted works	Swinburne University of Technology on 2022-06-01	<1%
3	Internet	www.coursehero.com	<1%
4	Internet	www.mdpi.com	<1%
5	Publication	Mehdi Ghayoumi. "Generative Adversarial Networks in Practice", CRC Press, 2023	<1%
6	Submitted works	Monash University on 2022-05-19	<1%
7	Internet	assets.researchsquare.com	<1%
8	Internet	core.ac.uk	<1%
9	Internet	link.springer.com	<1%
10	Submitted works	HAS Den Bosch on 2025-03-14	<1%

11	Submitted works	JIS University on 2025-04-17	<1%
12	Internet	www.frontiersin.org	<1%
13	Internet	www.hindawi.com	<1%
14	Submitted works	University of Greenwich on 2022-01-17	<1%
15	Submitted works	University of Witwatersrand on 2023-09-14	<1%
16	Internet	research.utwente.nl	<1%
17	Internet	www.earth-science.net	<1%
18	Submitted works	CSU, Fullerton on 2024-12-13	<1%
19	Submitted works	Chung-Ang University on 2023-06-14	<1%
20	Submitted works	German Jordanian University on 2025-02-12	<1%
21	Submitted works	University of South Australia on 2024-06-09	<1%
22	Publication	Codruț-Andrei Diaconu, Konrad Heidler, Jonathan L. Bamber, Harry Zekollari. "Mu...	<1%
23	Submitted works	University of Dundee on 2025-03-27	<1%
24	Internet	oulurepo.oulu.fi	<1%

25	Internet	www.researchsquare.com	<1%
26	Submitted works	Colorado Technical University Online on 2025-04-19	<1%
27	Publication	Fang, Ruoming. "Application of Deep Learning to Land Cover Classification: Practi...	<1%
28	Internet	studenttheses.uu.nl	<1%
29	Submitted works	University of Canberra on 2024-03-31	<1%
30	Submitted works	University of Essex on 2025-04-25	<1%
31	Submitted works	University of Maryland, Global Campus on 2023-12-12	<1%
32	Internet	climatefactchecks.org	<1%
33	Submitted works	Johns Hopkins University on 2024-09-21	<1%
34	Submitted works	University of East London on 2025-03-24	<1%
35	Publication	Dorota Jelonek, Narendra Kumar, Mamta Chahar, Rusudan Kinkladze, Lilla Knop. ...	<1%
36	Submitted works	University of Northampton on 2025-01-05	<1%
37	Internet	www.freepatentsonline.com	<1%
38	Submitted works	Harrisburg University of Science and Technology on 2019-12-02	<1%

39	Submitted works	RMIT University on 2024-11-11	<1%
40	Submitted works	University of Leicester LTI on 2025-04-27	<1%
41	Submitted works	University of Stirling on 2024-09-10	<1%
42	Internet	www.groundwatergovernance.org	<1%
43	Publication	"Precision agriculture '19", Walter de Gruyter GmbH, 2019	<1%
44	Submitted works	Ajou University Graduate School on 2024-03-20	<1%
45	Submitted works	Al Akhawayn University in Ifrane on 2025-04-26	<1%
46	Submitted works	National University of Ireland, Maynooth on 2023-08-07	<1%
47	Submitted works	Robert Kennedy College on 2025-04-13	<1%
48	Publication	Shashi Kant Dargar, Shilpi Birla, Abha Dargar, Avtar Singh, D. Ganeshaperumal. "...	<1%
49	Submitted works	University of Sydney on 2024-11-01	<1%
50	Internet	uspa.memberclicks.net	<1%
51	Internet	www.forbes.com	<1%
52	Submitted works	Curtin University of Technology on 2013-06-03	<1%

53	Publication	Iam Palatnik de Sousa, Marley Maria Bernardes Rebuzzi Vellasco, Eduardo Costa ...	<1%
54	Submitted works	Institución Tecnológica Metropolitana de Medellín on 2025-03-27	<1%
55	Publication	Konstantin A. Maslov, Claudio Persello, Thomas Schellenberger, Alfred Stein. "Glo...	<1%
56	Submitted works	Notre Dame of Marbel University on 2025-04-27	<1%
57	Submitted works	University of Huddersfield on 2020-04-30	<1%
58	Submitted works	University of Plymouth on 2014-01-28	<1%
59	Submitted works	University of Sheffield on 2020-01-22	<1%
60	Internet	arxiv.org	<1%
61	Internet	eartharxiv.org	<1%
62	Internet	pdf.usaid.gov	<1%
63	Internet	www.researchgate.net	<1%
64	Internet	www.tandfonline.com	<1%
65	Submitted works	Baker University on 2024-07-30	<1%
66	Publication	Shalli Rani, Ashu Taneja. "WSN and IoT - An Integrated Approach for Smart Applic...	<1%

67	Submitted works	
VIT University on 2023-02-14		<1%
68	Submitted works	
University of Sheffield on 2020-09-14		<1%
69	Submitted works	
VIT University on 2025-04-16		<1%
70	Publication	
Yingisani Chabalala, Elhadi Adam, Khalid Adem Ali. "Machine Learning Classificati...		<1%
71	Publication	
Zunxun Liang, Fangxiong Wang, Jianfeng Zhu, Peng Li, Fuding Xie, Yifei Zhao. "Au...		<1%





**NUST**

**SCHOOL OF ELECTRICAL ENGINEERING  
& COMPUTER SCIENCE**

# **IceWatch**

## **GLOF Prediction Using Deep Learning**

**Final Year Project Report**

**By**

**Zuha Fatima**

**Muhammad Anser Sohaib**

**Muhammad Talha**

**In Partial Fulfillment**

**Of the Requirements for the degree**

**Bachelor of Electrical Engineering (BEE)**

**School of Electrical Engineering and Computer Science**

**National University of Sciences and Technology**

**Islamabad, Pakistan**

**(2025)**

2

# DECLARATION

We hereby declare that this project report entitled **IceWatch** submitted to the **School of Electrical Engineering & Computer Science (SEECs)**, is a record of an original work done by us under the guidance of Advisor Ms. Ayesha Kanwal and Co-Advisor Ms. Nazia Pervaiz and that no part has been plagiarized without citations. This project work is submitted in the partial fulfillment of the requirements for the degree of Bachelor of Electrical Engineering.

## Team Members

## Signature

Zuha Fatima

Muhammad Anser Sohaib

Anser

Muhammad Talha

Talha

## Supervisor

## Signature

Ms. Ayesha Kanwal

Ms. Nazia Pervaiz

Nazia Pervaiz

**Date:**

25<sup>th</sup> April, 2025

## DEDICATION

44 We dedicate this work to our families and mentors, whose unwavering support and encouragement made this journey possible.

## 2 ACKNOWLEDGEMENTS

We would like to express our sincere gratitude to our Advisor Ms. Ayesha Kanwal and Co-Advisor Ms. Nazia Pervaiz for their invaluable guidance and support throughout the course of this project.

# TABLE OF CONTENT

## Contents

IceWatch.....	0
DECLARATION.....	1
DEDICATION .....	2
ACKNOWLEDGEMENTS.....	2
TABLE OF CONTENT .....	3
LIST OF FIGURES .....	9
LIST OF TABLES .....	11
ABSTRACT.....	12
Chapter 1 .....	13
INTRODUCTION .....	13
Background and Motivation .....	13
Project Aim .....	13
System Overview .....	14
TempFlow: Temperature Prediction Using LSTM .....	14
TerraFlow: Glacier Velocity Prediction Using Transformer .....	14
RiskFlow: CNN Classifier for Velocity Mosaics.....	14
Parameter Selection .....	14
Contributions.....	15
Organization of the Paper .....	15
Chapter 2.....	16
LITERATURE REVIEW .....	16

Chapter 3 .....	18
PROBLEM DEFINITION .....	18
GLOF Trends and Growing Risks .....	18
1. Increasing Frequency of GLOFs Globally.....	18
2. Global Distribution of GLOF Occurrences.....	19
3. Expansion of Glacial Lakes in the Himalayas .....	20
4. Populations at Risk from GLOFs.....	20
Causes of GLOFs.....	21
1. Climate Change and Glacier Retreat.....	21
2. Dam Instability (Moraine and Ice Dams) .....	21
3. Triggering Mechanisms .....	21
4. Changes in Lake Geometry.....	21
Impacts of GLOFs.....	23
• Human Displacement:.....	23
• Infrastructure Damage: .....	23
• Economic Losses: .....	23
• Ecological Effects: .....	23
Importance of Early Warning and Monitoring.....	23
Limitations of Existing GLOF Warning Systems: .....	24
1. Heavy Reliance on Manual Monitoring and Field Sensors .....	24
2. Lack of Multivariate Data Integration .....	24
3. Threshold-Based Logic Is Rigid and Outdated.....	24
4. Poor Temporal Prediction Capabilities .....	24
5. Limited Geographic Coverage.....	25
Addressing Limitations through IceWatch: .....	25

1. Automated Monitoring Using Satellite Data.....	26
2. Multimodal Deep Learning Integration .....	26
3. Learned Representations, Not Static Thresholds .....	26
4. Early Forecasting with Lead Time .....	26
5. Scalable and Deployable System .....	26
Chapter 4.....	27
METHODOLOGY .....	27
Framework: .....	27
1. Riskflow .....	27
2. TerraFlow .....	27
3. TempFlow .....	27
Data Sources and Characteristics:.....	27
Sentinel-2 Satellite Imagery (for RiskFlow CNN Model).....	28
NASA ITS LIVE Glacier Velocity Data (for TerraFlow Model).....	31
MODIS Land Surface Temperature (for TempFlow Model): .....	32
Supplementary Labels and Ground Truth .....	35
TerraFlow Methodology .....	36
Data Collection: .....	36
Preprocessing: .....	36
TempFlow Methodology.....	38
Data Collection: .....	38
Preprocessing: .....	38
Data Structuring:.....	39
Model Processing:.....	39
Output Generation:.....	39

RiskFlow Methodology .....	40
Data Collection: .....	40
Preprocessing: .....	40
Dataset Splitting: .....	40
Model Processing: .....	40
Output Generation and Evaluation: .....	41
Chapter 5 .....	42
DETAILED DESIGN AND ARCHITECTURE .....	42
TerraFlow Design and Architecture: .....	42
Input Layer: .....	42
Embedding Layer: .....	42
Positional Encoding: .....	42
Transformer Encoder: .....	42
Dropout and Output: .....	43
Model Parameters: .....	43
Optimizer and Loss: .....	43
TempFlow Design and Architecture .....	45
Reason for Using LSTM: .....	47
• Long-Term Dependency Capture: .....	47
• Non-Linear Modeling: .....	47
• Robustness to Noise: .....	47
• Sequential Processing: .....	48
Architecture: .....	48
Input Layer: .....	48
Bidirectional LSTM (128 units): .....	48

Dropout (0.4): .....	48
LSTM (64 units): .....	48
Dropout (0.4): .....	49
LSTM (32 units): .....	49
Dropout (0.3): .....	49
Dense Layer: .....	49
Optimizer and Loss: .....	49
RiskFlow Design and Architecture .....	50
Input Layer: .....	51
Conv2d Layer (128 filters): .....	51
Maxpooling (2×2): .....	51
Conv2d Layer (32 filters): .....	51
Maxpooling (2×2): .....	51
Flattening: .....	51
Fully Connected Layer (128 neurons): .....	51
Output Layer: .....	52
Optimizer and Loss: .....	52
Integration in IceWatch System .....	53
Data Flow: .....	53
System Architecture: .....	53
Implementation: .....	53
Chapter 6 .....	54
IMPLEMENTATION AND TESTING .....	54
Implementation .....	54
Training Environment .....	54



Model Development.....	55
General Training Challenges.....	62
Testing.....	62
Testing Process.....	63
Chapter 7.....	64
RESULTS AND DISCUSSION.....	64
Web Application Overview.....	64
• Interface Description:.....	64
• Parallel Model Execution:.....	65
• Deployment:.....	65
• User Features: .....	65
Chapter 8.....	67
CONCLUSION AND FUTURE WORK.....	67
Conclusion .....	67
Future Work .....	67
Chapter 9.....	69
REFERENCES .....	69

# LIST OF FIGURES

Figure 1: Number of documented GLOF events globally from 1900 to 2020, illustrating an upward trend in occurrence frequency (Lützow et al., 2023).	18
Figure 2 Spatial distribution of recorded GLOF events globally(Lützow et al.,2023[6]).	19
Figure 3 Regional distribution of reported (a) GLOF source locations, (b) all GLOFs, and (c) ratio of the number of GLOFs and source locations.	20
Figure 4 Glacial Lake Formation	22
Figure 5 Glacial Lake	22
Figure 6 Glacial Lake Expansion	22
Figure 7 GLOF occurs after Glacial Lake Breaches	23
Figure 8 Region of Interest: Shisper Glacier	28
Figure 9 Sentinel-2, rendering of Bands B8, B11 and B12 on QGIS	29
Figure 10 Time series of LST from 2000 to 2022	<b>Error! Bookmark not defined.</b>
Figure 11 A kernel density plot showing LST distributions across high, medium, and low-quality	35
Figure 12 TerraFlow Architecture (a)	43
Figure 13 TerraFlow Architecture (b)	44
Figure 14 Terraflow Model	45
Figure 15 Tempflow Model	46
Figure 16 Tempflow model (a)	46
Figure 17 Tempflow model (b)	47

Figure 18 Code and Architechture of Tempflow .....	49
Figure 19 RiskFlow Model Architecture .....	50
Figure 20 RiskFlow Architecture.....	52
Figure 21 Nvidia GH200 overview using nvidia-smi.....	55
Figure 22 Accuracy of RiskFlow .....	60
Figure 23 RiskFlow Confusion Matrix .....	61
Figure 24 IceWatch Web App (a).....	66
Figure 25 IceWatch Web App (b).....	66

# LIST OF TABLES

Table 1 Traditional GLOF Detection Systems vs IceWatch .....	25
Table 2 Sentinel-2 Bands used in RiskFlow ( wavelength, resolution, Band Number)29	
Table 3 Reasons For Selection of Sentinel-2 bands for GLOF Detection .....	30
Table 4 Features Used for TerraFlow.....	31
Table 5 Key Variables for TempFlow .....	33
Table 6 Model Used, Respective Datasets and their Sources, Purpose .....	36
Table 7 RiskFlow Training Details .....	58
Table 8 RiskFlow Evaluation and Results .....	59

# ABSTRACT

Glacial Lake Outburst Floods (GLOFs) are sudden and devastating events that occur when the natural dams containing glacial lakes fail, leading to massive downstream flooding. With the ongoing impacts of climate change, the size and number of glacial lakes are rapidly increasing, making GLOFs more frequent and unpredictable. These floods pose a significant threat to communities, infrastructure, and ecosystems in high-altitude regions, where early detection and prevention measures are limited. Traditional methods of monitoring and predicting GLOFs often rely on manual observations and static risk maps, which are insufficient for capturing the dynamic changes in glacial environments.

To address this growing challenge, we developed **IceWatch**, an end-to-end system designed to predict the probability of GLOF occurrence using machine learning. The system analyzes key features such as glacial lake area, elevation, distance to glacier fronts, geographic coordinates, and changes in lake size and position over time. The IceWatch system delivers increased glacial lake risk assessments through its probability-based prediction method which surpasses conventional binary classification methods. Users benefit from a visual dashboard which provides them with straightforward access to lake monitoring as well as probability score displays and high-risk geo-area identification capabilities. Leaders who use predictive modeling combined with graphical tools obtain decisions-ready information for taking purposeful actions.

The IceWatch system works upon enhancing disaster warning capabilities along with emergency readiness efforts in regions where populations actively face disaster risks. The system provides prompt data-based predictions that help researchers together with policymakers and emergency management agencies to establish intervention requirements while minimizing the destructive impact of GLOFs. This project illustrates how machine learning and intelligent visualization technologies enhance climate protection to protect vulnerable communities at risk.

# Chapter 1

## INTRODUCTION

### Background and Motivation

58  
9  
64  
Glacial Lake Outburst Floods (GLOFs) represent the most dangerous and unpredictable geological disasters which occur within mountainous areas. A natural dam created from ice or moraines, or glacial debris will fail during GLOFs to release an extensive amount of water rapidly from glacial lakes. Glacial Lake Outburst Floods result in destructive flooding which jeopardizes human settlements and infrastructure and agricultural land and environment. Current climate change effects have accelerated glacial retreat worldwide which created the expansion of glacial lakes and the development of new ones. The expansion of glacial outburst flooding events has increased notably while becoming more intense in mountainous vulnerable areas including the Karakoram and Hindu Kush along with the greater Himalayas.

The growing threat to society is not properly addressed through existing traditional GLOF monitoring systems and warning mechanisms. Current hazard assessment techniques include static hazard mapping with manual observation techniques and threshold criteria, but these fail to reflect the natural dynamic modifications of glacial lakes together with climatic patterns and glacier movement. The present situation requires adaptive data-driven systems which monitor risk using various environmental variables in continuously operating modes.

### Project Aim

The research introduces IceWatch as a full-scale machine learning prediction tool that evaluates the possibility of GLOF occurrences. Satellite imagery provides environmental data to this system which includes land surface temperature along with glacier velocity measurements and glacial lake characteristics and velocity distribution maps. Implementing Transformers, LSTMs and Convolutional Neural Networks (CNNs) within IceWatch builds a superior system to assess real-time potential GLOF events in sensitive areas.

## System Overview

The IceWatch framework contains three essential components that power its operation:

### **TempFlow: Temperature Prediction Using LSTM**

The TempFlow model uses LSTM-based modules to carry out training while processing NASA MODIS Land Surface Temperature (LST) data during twenty-two consecutive years. The system detects large climate pattern shifts together with sudden temperature changes. The evolution of glacial patterns functions as an essential signal to determine upcoming glacial lake formation periods or fast growth situations from ice melt. Quality-controlled data combined with cyclic time features in the model allows the production of reliable seasonal temperature predictions.

### **TerraFlow: Glacier Velocity Prediction Using Transformer**

**TerraFlow** utilizes a Transformer-based sequence model that predicts upcoming glacier ice velocity by processing spatial-temporal features containing latitudinal and longitudinal positions in addition to day-of-year data encoding. TerraFlow operates with millions of NASA ITS\_LIVE observations to detect glacier dynamics modifications which suggest rising glacial lake pressure or instability through flow acceleration or surging signatures.

### **RiskFlow: CNN Classifier for Velocity Mosaics**

The IceWatch system combines a CNN-based image classifier with velocity mosaics to process satellite imagery showing glacier motion fields as a part of its design. This model detects anomalous spatial patterns in glacier flow, offering a visual cross-validation layer for TerraFlow predictions and improving the overall interpretability of the system.

## Parameter Selection

The choice of parameters in IceWatch is informed by glaciological science and prior GLOF case studies. Rising land surface temperature is a well-established driver of glacier melt and lake formation. Ice velocity changes reflect glacier surges or dam instability, while lake area and distance from the glacier terminus are commonly used indicators of hazard potential. Elevation and geolocation provide spatial context for generalizing across regions. Integrating these variables ensures that IceWatch captures both physical and temporal precursors to GLOF events.

## Contributions

The key contributions of this research are:

- A **multi-modal GLOF prediction framework** combining sequential (LSTM), attention-based (Transformer), and image-based (CNN) deep learning models.
- A **probability-based GLOF risk scoring system** that moves beyond binary classification to provide nuanced risk assessments.
- The development of a **visual dashboard interface** that enables researchers and disaster response authorities to monitor, interpret, and act upon real-time GLOF risk.
- A scalable pipeline that can be adapted to various high-risk glacial regions beyond the initial study area (Shisper Glacier, Pakistan).

## Organization of the Paper

The remainder of this paper is structured as follows:

Section 2 reviews relevant literature review in GLOF detection and glacier modeling.

Section 3 describes the Problem Definition

Section 4 outlines the Methodology

Section 5 presents the Detailed Architecture and Design

Section 6 Implementation and Testing

Section 7 Results and Discussion

Section 8 Conclusion and Future Work



## Chapter 2

# LITERATURE REVIEW

Recent advancements in remote sensing and machine learning have significantly contributed to the monitoring and prediction of Glacial Lake Outburst Floods (GLOFs).

Early research predominantly focused on mapping glacial lakes and assessing their hazard potential using satellite imagery. Fujita et al. [1] quantified potential flood volumes from Himalayan glacial lakes using topographical and morphological parameters, providing foundational knowledge for hazard estimation. Worni et al. [2] emphasized the importance of dynamic lake volume monitoring and introduced process-based flood simulations, although these methods lacked real-time predictive capability.

The application of machine learning has improved GLOF susceptibility mapping. Zhou et al. [3] proposed a hybrid machine learning framework combining environmental indicators and glacier evolution data to predict debris flows associated with GLOFs. Similarly, Abbas et al. [4] compared machine learning, deep learning, and hybrid models for GLOF susceptibility mapping in the Central Karakoram National Park, demonstrating that hybrid approaches outperform classical models. However, these models primarily relied on static terrain features without incorporating temporal glacier movement data.

Deep learning techniques have been employed for glacial lake segmentation. Pandey and Malakar [5] utilized CNNs for water body detection from remote sensing imagery, achieving high segmentation accuracy. Dey et al. [6] improved upon this by applying U-Net architectures for glacial lake extraction, although their models were sensitive to cloud interference and seasonal variations in the imagery.

Recent work has introduced Transformer architectures for glacier monitoring. Maslov et al. [7] developed Glacier-VisionTransformer-U-Net (GlaViTU), a hybrid CNN-Transformer model that outperformed traditional CNNs in glacier mapping tasks by capturing long-range spatial dependencies. However, limited labeled datasets and complex training processes remain challenges for Transformer-based glacier analysis.

Efforts to integrate multi-modal datasets have shown promise. **Mohanty et al.** [8] reviewed the use of **multi-sensor approaches** (combining optical, SAR, and DEM data) for glacier velocity and facies characterization, highlighting the advantages of fusing different observation sources. **Peng et al.** [9] explored the use of deep learning combined with open Earth observation datasets for **large-scale glacier mapping** but noted persistent challenges in real-time processing and data quality variability.

Despite these advancements, existing systems typically focus on static mapping or susceptibility scoring rather than real-time GLOF prediction integrating thermal, spatial, and dynamic glacier movement data. The gaps identified in previous works motivate the development of a system that combines temporal temperature forecasting, dynamic velocity prediction, and spatial anomaly detection for a comprehensive, real-time GLOF early warning solution.

# Chapter 3

## PROBLEM DEFINITION

### GLOF Trends and Growing Risks

Glacial Lake Outburst Floods (GLOFs) have emerged as an escalating hazard in high-mountain regions, driven by accelerating glacier retreat under global warming. Recent studies and observational datasets highlight the alarming rise in GLOF events globally. Satellite observations show a substantial increase in both the number and size of glacial lakes globally, particularly in vulnerable regions such as the Himalayas, the Andes, and parts of North America.

#### 1. Increasing Frequency of GLOFs Globally

Recent data show a steady increase in the frequency of documented GLOF events since the early 20th century, with a pronounced rise from the 1990s onward. This trend correlates with intensified glacier retreat due to anthropogenic climate change.




Figure 1: Number of documented GLOF events globally from 1900 to 2020, illustrating an upward trend in occurrence frequency (Lützow et al., 2023).

## 2. Global Distribution of GLOF Occurrences

GLOFs are not geographically uniform. High-risk zones are predominantly located in the Himalayas, Andes, and certain parts of North America (e.g., Alaska and Western Canada). The map below illustrates global hotspots for recorded GLOF events.




Figure 2 Spatial distribution of recorded GLOF events globally (Lützow et al., 2023 [6]).

This figure gives an overview map of the source locations of GLOFs and selected regional database contents. Colors show the lake type of the GLOF locations (bubbles) and proportions of GLOF locations in each region (pie charts). Boxes are regional statistics on the total number of outbursts (and the number of outbursts of GLOFs since the availability of satellite images); the percentage of cases with reported dam type and drainage mechanisms; and parameters on GLOF magnitude such as peak discharge ( $Q_p$ ), flood volume ( $V_0$ ), and the area before ( $A_b$ ) and after ( $A_a$ ) the GLOF.




Figure 3 Regional distribution of reported (a) GLOF source locations, (b) all GLOFs, and (c) ratio of the number of GLOFs and source locations.

### 3. Expansion of Glacial Lakes in the Himalayas

Glacial lakes are increasing both in number and area, particularly across the Himalayas, thereby heightening the potential for GLOFs. Between 2011 and 2023, a significant surge, over **10% increase in glacial lake formation** has been observed. [7]

### 4. Populations at Risk from GLOFs

An estimated **15 million people** globally live within vulnerable zones that could be impacted by GLOFs. South Asian countries, particularly India, Pakistan, Nepal, and Bhutan host the largest at-risk populations.

# Causes of GLOFs

The formation and sudden failure of glacial lakes are influenced by several interrelated natural and climatic factors:

## 1. Climate Change and Glacier Retreat

10

- Rising global temperatures are accelerating glacier melt, leading to the rapid formation and expansion of proglacial lakes.
- As glaciers retreat, the ice acts as a dam, behind which meltwater accumulates.
- Increased melt rates lead to higher lake volumes, increasing hydrostatic pressure on natural dams, making them more prone to failure.

## 2. Dam Instability (Moraine and Ice Dams)

- Many glacial lakes are dammed by loose accumulations of debris known as **moraines** or by remnant **glacier ice**.
- Ice dams are susceptible to melting and mechanical failure due to warming temperatures and internal stress.

## 3. Triggering Mechanisms

Several immediate triggers can precipitate a dam failure:


- **Ice or Rock Avalanches:** Landslides, icefalls, or rockfalls from adjacent mountain slopes into the lake can cause a sudden surge in water level (known as a displacement wave), overtopping and breaching the dam.
- **Earthquakes:** Seismic activity can destabilize the dam material, leading to a rapid collapse.
- **Heavy Rainfall or Rapid Snowmelt:** Excess water input into already full glacial lakes can stress dam integrity.
- **Subglacial Drainage:** Some GLOFs occur when water accumulates beneath a glacier and then bursts through the ice.

## 4. Changes in Lake Geometry


- Expansion of lake surface area and depth over time increases the hydraulic pressure acting on the dam.
- Thinning glaciers adjacent to lakes can reduce support against the moraine walls, making breaches more likely.



*Figure 4 Glacial Lake Formation*




*Figure 5 Glacial Lake*



*Figure 6 Glacial Lake Expansion*





*Figure 7 GLOF occurs after Glacial Lake Breaches*

## Impacts of GLOFs

- **Human Displacement:** Villages and towns located in vulnerable river valleys can be inundated within hours of a GLOF event.
- **Infrastructure Damage:** Roads, bridges, hydroelectric plants, and irrigation systems are often severely impacted.
- **Economic Losses:** GLOFs can wipe out agricultural land and disrupt water supplies, impacting livelihoods.
- **Ecological Effects:** GLOFs can reshape river channels, erode forests, and deposit massive amounts of sediment, affecting ecosystems for years.

## Importance of Early Warning and Monitoring

Given their sudden nature and devastating impacts, developing effective GLOF monitoring, prediction, and early warning systems is crucial for safeguarding human life and minimizing economic and environmental damages. This underscores the urgency for innovative, scalable, and automated solutions, such as the IceWatch project proposed in this research, that integrate satellite observations, deep learning models, and environmental forecasting to provide timely GLOF risk assessments.



# Limitations of Existing GLOF Warning Systems:

## 1. Heavy Reliance on Manual Monitoring and Field Sensors

Most operational GLOF monitoring systems depend on in-situ sensors such as water level gauges, weather stations, or manual glacier surveys. These instruments:

- Are often expensive to install and maintain in remote glacial environments.
- Provide localized data that may not capture regional trends.
- Are vulnerable to harsh weather conditions or mechanical failure.

## 2. Lack of Multivariate Data Integration

Traditional EWS usually monitor one or two variables (e.g., lake area or rainfall) without considering dynamic factors like glacier motion or thermal stress. This narrow focus:

- Limits the ability to detect complex, multicausal events.
- Fails to account for precursors like surface velocity spikes or anomalous warming.

## 3. Threshold-Based Logic Is Rigid and Outdated

Current systems often operate on static rules or thresholds (e.g., lake size > X = high risk), which:

- Cannot adapt to diverse lake-glacier morphologies.
- Are prone to false positives and false negatives.
- Do not learn from past events or trends.

## 4. Poor Temporal Prediction Capabilities

Most existing models offer situational awareness (e.g., current lake level), but not true forecasting. As a result:

- Warnings are issued late, often after triggering mechanisms have already started.
- Authorities have little lead time for evacuation or mitigation.

## 5. Limited Geographic Coverage

Due to cost and access constraints, only a small subset of glacial lakes are currently monitored in high-risk regions like the Himalayas, Andes, and Alaska.

*Table 1 Traditional GLOF Detection Systems vs IceWatch*

Feature	Traditional GLOF Systems	IceWatch
Data Source	Ground sensors, static maps	Dynamic satellite + temporal data
Prediction Type	Rule-based alerts	AI-driven forecasting
Multivariable Analysis	No	Yes (image + velocity + temperature)
Scalability	Limited	Global (remote sensing)
Temporal Forecasting	Reactive	Proactive
Cost	High (maintenance, logistics)	Low (cloud-based, automated)

## Addressing Limitations through IceWatch:

*IceWatch* introduces a next-generation, AI-powered early warning system that addresses the core limitations of conventional approaches through a multi-model, data-driven framework.

## 1. Automated Monitoring Using Satellite Data

Unlike sensor-based systems, *IceWatch* operates on open-access satellite data (Sentinel-2, MODIS, ITS LIVE), enabling:

- Wide-scale, repeatable, and remote monitoring of hundreds of glaciers.
- Cost-effective surveillance even in inaccessible or politically sensitive regions.

## 2. Multimodal Deep Learning Integration

The system integrates diverse environmental signals via:

- A **CNN** model for visual GLOF detection from imagery.
- A **Transformer** model (*TerraFlow*) for glacier motion prediction.
- An **LSTM** model (*TempFlow*) for temperature trend forecasting.

This fusion captures both immediate visual cues and slow-onset precursors, improving detection accuracy.

## 3. Learned Representations, Not Static Thresholds

By training on historical event data and natural variability, *IceWatch*:

- Learns complex spatiotemporal patterns leading to GLOFs.
- Can generalize across different glaciers and climate regimes.
- Offers probabilistic predictions rather than binary alerts.

## 4. Early Forecasting with Lead Time

With its time-series forecasting capabilities, especially via *TempFlow*, *IceWatch*:

- Anticipates thermal and mechanical stress on glaciers days to weeks in advance.
- Provides lead time for disaster response planning.

## 5. Scalable and Deployable System

*IceWatch* is:

- Cloud-deployable (via platforms like Google Colab, AWS, or GEE).
- Lightweight enough for integration into local disaster management dashboards.
- Easily extensible with additional data sources (e.g., radar, precipitation).

## Chapter 4

# METHODOLOGY

Project IceWatch is a real-time Glacial Lake Outburst Flood (GLOF) prediction system that integrates three deep learning frameworks: **TerraFlow** (transformer-based for ice velocity prediction), **TempFlow** (LSTM-based for temperature forecasting), and **RiskFlow** (CNN-based for GLOF classification). These models analyze geospatial, temporal, and satellite imagery data to monitor and predict GLOF events in the **Shispare Glacier** region (Lat: 36.32°–36.47°, Lon: 74.60°–74.90°). Below is a detailed methodology, with each model's details organized under its respective heading, including data sources, preprocessing, architecture, training, evaluation, and integration into the IceWatch system.

### Framework:

The three specialized models make up the IceWatch framework structure:

1. **Riskflow** – A CNN Model Detects visual features indicative of GLOF events from Sentinel-2 satellite imagery.
2. **TerraFlow** – TerraFlow utilizes Transformer technology for predicting glacier surface velocity through the analysis of spatiotemporal table data.
3. **TempFlow** – These three models provide outputs that enable assessment of GLOF risk levels for delivering meaningful insights.

The IceWatch system establishes multi-modal GLOF detection through its integrated models to monitor the Shispare Glacier area which demonstrates high GLOF vulnerability due to its sensitive glacial activities and climate changes.

### Data Sources and Characteristics:

IceWatch requires various datasets which capture how the Shispare Glacier region changes in space and time and spectrum. The models implement diverse datasets because each predictive task has unique requirements.

Internal to IceWatch each of the main three operation models receives unique domain-specific datasets for visual detection through CNN and glacier velocity using TerraFlow and temperature prediction by TempFlow.

## Sentinel-2 Satellite Imagery (for RiskFlow CNN Model)

**Source:** European Space Agency (ESA) Copernicus Open Access Hub

**Temporal Range:** 2017 – 2024

**Spatial Resolution:** 10m – 20m

The research focuses on Shispare Glacier through Sentinel-2 satellite images which compose its dataset. The European Space Agency (ESA) operates Sentinel-2 which delivers multispectral imagery at high definition through its satellites. Google Earth Engine processed the images followed by their export from the system.



*Figure 8 Region of Interest: Shispare Glacier*



## Sample image from dataset:




Figure 9 Sentinel-2, rendering of Bands B8, B11 and B12 on QGIS

## Bands Used:

The selection of Sentinel-2 bands B2, B3, B4, B8, B11, and B12 was based on their proven sensitivity to snow, ice, meltwater, and vegetation dynamics — all critical indicators for early prediction of Glacial Lake Outburst Flooding (GLOF) events

Table 2 Sentinel-2 Bands used in RiskFlow ( wavelength, resolution, Band Number)

Band Name	Band Number	Wavelength (nm)	Resolution	
Blue	B2	490 nm	10 m	
Green	B3	560 nm	10 m	
Red	B4	665 nm	10 m	
NIR (Near-Infrared)	B8	842 nm	10 m	
SWIR-1 (Shortwave Infrared 1)	B11	1610 nm	20 m	
SWIR-2 (Shortwave Infrared 2)	B12	2190 nm	20 m	

*Table 3 Reasons For Selection of Sentinel-2 bands for GLOF Detection*

Band(s)	Reason
<b>B2 (Blue)</b>	Very sensitive to <b>snow and ice reflectance</b> ; useful for detecting glaciers and frozen surfaces.
<b>B3 (Green)</b>	Captures <b>vegetation, bare soil, and surface features</b> contrast; helps distinguish glacier surroundings.
<b>B4 (Red)</b>	Key for <b>snow/ice monitoring</b> ; good contrast between glacier ice and bare earth.
<b>B8 (NIR)</b>	Strong separation between <b>vegetation, water, and ice</b> ; important for detecting meltwater or glacial lakes forming.
<b>B11 (SWIR-1)</b>	Sensitive to <b>snow grain size, melting ice, and moisture</b> ; helps monitor glacier melting and flood-prone areas.
<b>B12 (SWIR-2)</b>	Detects <b>liquid water, wet surfaces, and glacial lake formation</b> more clearly than visible bands.

## Classes

- **GLOF occurrence (label 1):** Images containing visible signs of glacial lake formation or outburst conditions.
- **NO-GLOF (label 0):** Images without signs of imminent outburst flooding.

## Description:

Sentinel-2 images were used to train the CNN model for detecting visual patterns of GLOF occurrence. These bands are sensitive to water, vegetation, snow, and ice—key features for glacial monitoring. Images were collected for the Shisper Glacier region, preprocessed (resized to 128×128, normalized), and labeled for binary classification: GLOF (1) or NO-GLOF (0).

## NASA ITS LIVE Glacier Velocity Data (for TerraFlow Model)

The TerraFlow model utilizes the **Inter-mission Time Series of Land Ice Velocity and Elevation (ITS LIVE)** dataset, developed by NASA's Jet Propulsion Laboratory (JPL). This publicly available dataset provides high-resolution measurements of glacier surface velocity across the globe, derived using feature tracking algorithms applied to optical satellite imagery such as Landsat 4, 5, 7, and 8.

### Temporal and Spatial Coverage:

- **Region of Interest:** Shisper Glacier, Karakoram Range, Pakistan
- **Coordinates:** Latitude 36.25°–36.50°, Longitude 74.50°–75.00°
- **Time Range:** January 1, 2000 – December 31, 2024
- **Temporal Resolution:** Daily observations
- **Raw Dataset Size:** ~73.6 million records globally
- **Filtered Dataset for TerraFlow:** ~11.9 million rows (Shisper region only)

*Table 4 Features Used for TerraFlow*

Variable Name	Description
latitude, longitude	Geographic coordinates of velocity point
date	Observation date, later encoded with cyclical features
vx, vy	Horizontal velocity components (m/day) along x and y directions
v	Magnitude of surface velocity, computed as $\sqrt{vx^2 + vy^2}$
velocity_avg, velocity_max	Aggregated daily mean and peak values per (lat, lon) point
day_of_year_sin, day_of_year_cos	Cyclical encodings of time to capture seasonal trends



The ITS LIVE dataset contains high-resolution velocity measurements of glacier surfaces, derived from feature tracking between satellite image pairs. For this project, ~11.9 million rows were processed from an initial 73.6 million, representing the Shisper Glacier. The data was cleaned, aggregated daily by coordinates, and used to predict velocity trends via the Transformer-based TerraFlow model.

## Data Characteristics and Challenges:

- **High Dimensionality:** The dataset spans over two decades of daily observations across many geolocations.
- **Heterogeneous Sampling:** Due to cloud cover, image quality, or satellite revisit limitations, some days are missing. These gaps were handled using interpolation or by masking the missing values.
- **Velocity Variability:** Glacier velocities ranged from 0 to >2 m/day, with seasonal peaks during summer. Anomalous spikes often correspond to surging activity or instability—important GLOF precursors.
- **Noise and Artifacts:** Optical feature-tracking introduces potential errors during rapid surface changes or image misalignment. These were mitigated by smoothing and filtering out extreme outliers.

---

## MODIS Land Surface Temperature (for TempFlow Model):

The TempFlow system leverages the **NASA MODIS Land Surface Temperature Product (MOD11A1 V6.1)**, a high-resolution dataset tailored for environmental monitoring. Below is a detailed breakdown of its characteristics:

### Source and Scope

- **Temporal Coverage:** February 24, 2000, to December 31, 2024, spanning 25 years.
- **Spatial Focus:** Shisper Glacier region (Latitude: 36.32°–36.47°, Longitude: 74.60°–74.90°), a critical area for studying glacier dynamics.
- **Resolution:** 1 km spatial resolution with daily temporal observations, yielding 98,879 data points across 11 unique coordinates and 8,989 unique dates (near-daily coverage).

## Key Variables

- **Primary Feature:** MOD11A1\_061\_LST\_Day\_1km, providing LST in Kelvin. Raw values range from 0–322 K, while quality-controlled (QC-filtered) data range from 250–330 K, reflecting typical surface temperatures in the region.
- **Quality Indicators:**
  - **QC\_Day\_MODLAND:** Flags indicating production quality (e.g., “LST produced, good quality”).
  - **QC\_Day\_LST\_Error\_Flag:** Error estimates categorized as  $\leq 1$  K,  $\leq 2$  K, or  $\leq 3$  K, with 78% of usable observations having errors  $\leq 2$  K.
  - **QC\_Day\_Emis\_Error\_Flag:** Metrics for emissivity-related errors, ensuring data reliability.

*Table 5 Key Variables for TempFlow*

Variable Name	Type	Description
LST_Day_1km	Numerical (float)	Land Surface Temperature in Kelvin (from MODIS MOD11A1 product)
QC_Day_MODLAND	Categorical	MODIS production quality flag (e.g., good, average, poor)
QC_Day_LST_Error_Flag	Categorical	Estimated LST error: $\leq 1$ K, $\leq 2$ K, or $\leq 3$ K
QC_Day_Emis_Error_Flag	Categorical	Emissivity error category (flag for radiative error levels)
latitude	Numerical (float)	Latitude of the pixel location
longitude	Numerical (float)	Longitude of the pixel location

date	DateTime	Date of observation
month_sin	Numerical (float)	Sine-transformed month value to capture annual seasonality
month_cos	Numerical (float)	Cosine-transformed month value to complement month_sin
quality_flag (derived)	Categorical	Combined quality classification: High, Medium, or Low (based on QC flags)
LST_scaled (derived)	Numerical (float)	Normalized or scaled version of LST for model stability

## Data Characteristics

The dataset exhibits pronounced seasonal patterns, with summer peaks (June–July) correlating with glacier melt and winter troughs (December–January) reflecting colder conditions. However, data quality poses a significant challenge:

- **Cloud Interference:** 61% of observations are flagged as low quality due to cloud cover, particularly during the melt season.
- **Quality Distribution:** 18% high quality, 22% medium quality, and 60% low quality, with high-quality data concentrated in winter months.





Figure 10 Time series of LST from 2000 to 2022

*Description:* The upper plot displays a time series of LST from 2000 to 2022, revealing daily fluctuations and seasonal peaks/troughs. The lower plot, a stacked time

series by year (2000–2023), illustrates monthly trends, highlighting a warming pattern consistent with the +0.8 K/decade trend.



*Figure 11 A kernel density plot showing LST distributions across high, medium, and low-quality*

**Description:** A kernel density plot showing LST distributions across high, medium, and low-quality categories, with peaks at approximately 280 K (high quality) and 290 K (medium quality), underscoring quality variability.

## Description:

This dataset was used to train the LSTM-based TempFlow model for temperature forecasting. Approximately 98,879 records were compiled from 11 coordinates over 9,000 unique dates in the Shisper Glacier area. The data exhibits strong seasonal cycles and long-term warming trends, which are critical for anticipating melt events. Low-quality readings (~60%) caused by cloud cover were filtered or imputed during preprocessing.

## Supplementary Labels and Ground Truth

**Source:** Manually curated from historical GLOF events, remote sensing literature, and visual inspection.

## Description:

Labels for the CNN model were created by visually identifying GLOF-related features in Sentinel-2 imagery near known historical GLOF events. Validation was done using literature, hydrological records (when available), and change detection over time.

*Table 6 Model Used, Respective Datasets and their Sources, Purpose*

Model	Dataset	Provider	Type	Years	Purpose
CNN	Sentinel-2 Imagery	ESA Copernicus	Satellite Image	2017–2024	GLOF detection
TerraFlow	ITS LIVE Velocity Data	NASA	Tabular Time Series	2000–2024	Glacier motion prediction
TempFlow	MODIS LST (MOD11A1)	NASA	Thermal Time Series	2000–2024	Surface temperature forecasting
All Models	Manually labeled GLOF Events	This project	Annotation	2000–2024	Ground truth for training

## TerraFlow Methodology

### Data Collection:

- Sources data from the NASA ITS\_LIVE dataset, containing 73.6 million rows of raw glacier velocity measurements (11 columns), reduced to 11.9 million rows and 9 columns post-preprocessing.
- Includes spatial coordinates (latitude, longitude), temporal features (dates), and velocity metrics.

### Preprocessing:

To prepare the dataset for sequence modeling with the Transformer-based TerraFlow architecture, the following preprocessing pipeline was applied:

## 1. Filtering by Region and Validity

Only entries within the bounding box of the Shisper Glacier were retained. Invalid or missing velocity values (NaNs or extreme outliers) were removed.

## 2. Temporal Aggregation

Because raw measurements could be sparse or noisy, daily aggregation was performed. For each (latitude, longitude) point, the average and maximum velocity values ( $v$ ) were computed for each day.

## 3. Feature Engineering

Temporal features were converted to cyclical encodings to preserve seasonality:

- $\text{day\_of\_year\_sin} = \sin(2\pi \times \text{day\_of\_year} / 365)$
- $\text{day\_of\_year\_cos} = \cos(2\pi \times \text{day\_of\_year} / 365)$

This allowed the model to learn seasonal patterns (e.g., summer glacier surge behavior).

## 4. Sequence Generation

The data was structured into time-series sequences of 32 days per sample. Each sequence consisted of:

- Daily velocity values ( $v$ )
- Spatial coordinates
- Temporal encodings

These sequences were used as input for the Transformer encoder.

## Model Processing:

- Processes sequences through 6 transformer encoder layers, each with 8-head multi-head attention, dropout (rate=0.2), and Add & Norm operations to model long-range dependencies.
- Applies feed-forward networks (FFNs) in each layer with additional dropout and normalization to learn complex patterns.

## Output Generation:

- Uses a final linear layer to output a single velocity prediction in meters per year.
- Identifies surge events (e.g., velocities  $>500$  m/yr) that may contribute to GLOF risk.

## Training:

12

- Trains with a batch size of 2048, 30 epochs, a learning rate of  $1e-5$ , and Mean Squared Error (MSE) loss.
- Uses Mean Absolute Error (MAE) as the evaluation metric, leveraging a CUDA-enabled GPU with mixed-precision training for efficiency.

# TempFlow Methodology

## Data Collection:

- Acquires data from the NASA MODIS Land Surface Temperature (LST) Product (MOD11A1 V6.1), spanning 2000-02-24 to 2024-12-31.
- Covers the Shispare Glacier with 98,879 observations at 1 km daily resolution, including MOD11A1\_061\_LST\_Day\_1km (250–330K) and quality indicators (QC\_Day\_MODLAND, QC\_Day\_LST\_Error\_Flag).

## Preprocessing:

To prepare the dataset for modeling, several preprocessing steps were applied:

- **Quality Filtering:**

Data was stratified into high (18%), medium (22%), and low (60%) quality categories. Low-quality, cloud-affected data were either excluded or imputed based on temporal patterns.

- **Temporal Encoding:**

Dates were converted to datetime objects, and cyclical features were introduced to capture seasonality:

- $\text{month\_sin} = \sin(2\pi \cdot \text{month}/12)$   $\text{month\_sin} = \sin(2\pi \cdot \text{month}/12)$
- $\text{month\_cos} = \cos(2\pi \cdot \text{month}/12)$   $\text{month\_cos} = \cos(2\pi \cdot \text{month}/12)$

### • Feature Selection:

Retained variables included LST values, QC flags, and spatial coordinates (latitude, longitude), while redundant metadata (e.g., MODIS\_Tile, bitmask details) were excluded to reduce dimensionality.

These steps enhance the dataset's usability, mitigating noise and aligning it with the LSTM's sequential input requirements.

### Data Structuring:

Structures data into 30-day sequences to align with the model's lookback window, capturing the  $\pm 15K$  annual cycle shown in the "Annual Temperature Cycle" plot.

### Model Processing:

- Processes sequences through three LSTM layers: LSTM(128) with ReLU and dropout (0.4), LSTM(64) with ReLU and dropout (0.4), and LSTM(32) with ReLU and dropout (0.2).
- Captures long-term trends (e.g., +0.8K/decade warming) and seasonal extremes.

### Output Generation:

- Outputs a temperature prediction in Kelvin via a dense layer, achieving a validation MSE of 12.3 K<sup>2</sup> and <3K error for seasonal extremes.
- Detects warming spikes (e.g., >150K) linked to GLOF risk, despite challenges with rapid melt events due to cloud cover.



# RiskFlow Methodology

## Data Collection:

70

- Collects Sentinel-2 imagery via Google Earth Engine, focusing on the Shispare Glacier.
- Utilizes six spectral bands: B2 (Blue), B3 (Green), B4 (Red), B8 (NIR), B11 (SWIR-1), and B12 (SWIR-2) for GLOF detection.

15

## Preprocessing:

The raw satellite images (.tif multiband files) underwent the following preprocessing steps:

- Band Selection:** Extracted only the 6 required bands.
- Normalization:** Pixel values scaled to [0, 1] by dividing by 255.
- Image Resizing:** All images resized to 128 × 128 dimensions for consistent input size.
- Augmentation:** Applied light augmentations to balance dataset and prevent overfitting:
  - Random horizontal and vertical flips
  - Random brightness adjustment
  - Random rotations
- Dataset Splitting:** Divided into training, validation, and testing sets.

5

## Dataset Splitting:

26

Splits data into 70% training, 15% validation, and 15% testing sets using train\_test\_split with random\_state=42 for reproducibility.

## Model Processing:

53

- Processes imagery through a Conv2d layer (128 filters, 3×3) with ReLU, followed by maxpooling (2×2).

4

- Applies a second Conv2d layer (32 filters,  $3 \times 3$ ) with ReLU and another maxpooling step to extract spatial features like glacial lakes.

5

- Passes features through a fully connected layer with ReLU (128 neurons) and a final dense layer with sigmoid activation.

### Output Generation and Evaluation:

- Outputs a GLOF probability (0–1), achieving a test accuracy of 94–95%, with precision (GLOF: 0.89, NO-GLOF: 0.93), recall (GLOF: 0.87, NO-GLOF: 0.94), and F1-scores (GLOF: 0.88, NO-GLOF: 0.935).
  - Trains with a batch size of 16, 5 epochs, Adam optimizer, and binary crossentropy loss, noting occasional false positives due to cloud cover or fresh snow.
-

## Chapter 5

# DETAILED DESIGN AND ARCHITECTURE

### TerraFlow Design and Architecture:

The TerraFlow model employs a Transformer-based architecture tailored for predicting glacier velocity, focusing on capturing complex spatiotemporal dependencies critical for identifying surge events in the Shispare Glacier.

#### Input Layer:

- Accepts sequences with a 32-day lookback window, incorporating 9 features: latitude, longitude, year, month\_sin, month\_cos, day\_sin, day\_cos, average velocity (avg\_velocity), and maximum velocity (max\_velocity). Input shape is (batch, 32, 9).

#### Embedding Layer:

- Projects the input features into a 256-dimensional space (d\_model=256) using a linear layer, creating a dense representation for Transformer processing.

#### Positional Encoding:

- Adds learnable positional encoding (shape: 32, 256) to preserve the temporal order of the sequence, essential since Transformers lack inherent sequential processing.

#### Transformer Encoder:

- Comprises 4 layers, each with 8-head multi-head attention (nhead=8) to capture long-range dependencies across spatial and temporal dimensions.
- Each layer includes a feed-forward network (FFN), dropout (rate=0.2), and Add & Norm operations to stabilize training and prevent overfitting.

- Outputs a sequence of shape (batch, 32, 256), capturing spatiotemporal patterns.

### Dropout and Output:

- Applies dropout (rate=0.2) to the last time step's output (shape: batch, 256), followed by a fully connected layer to produce a single velocity prediction in meters per year (shape: batch, 1).

### Model Parameters:

- Total trainable parameters: ~5.5 million, balancing capacity and computational efficiency.

### Optimizer and Loss:

- Uses the Adam optimizer with a learning rate of 1e-5 and Mean Squared Error (MSE) loss, suitable for regression tasks like velocity prediction.

The architecture leverages the NASA ITS\_LIVE dataset's preprocessed features, with cyclical encodings (day\_sin, day\_cos) ensuring the model captures seasonal velocity patterns, as evidenced by the Feature Correlation Matrix showing low correlation between cyclical features and velocity metrics, indicating their complementary roles.

```
SEQ_LENGTH = 32
BATCH_SIZE = 4096
NUM_EPOCHS = 50
LEARNING_RATE = 0.00001
D_MODEL = 512
NHEAD = 8
NUM_LAYERS = 6
DROPOUT = 0.2
```

Figure 12 TerraFlow Architecture (a)

```
class TransformerRegressor(nn.Module):
    """Transformer model for velocity prediction."""
    def __init__(self, input_dim, d_model, nhead, num_layers, dropout=0.1):
        super(TransformerRegressor, self).__init__()
        self.embedding = nn.Linear(input_dim, d_model)
        self.pos_encoder = nn.Parameter(torch.zeros(1, 1000, d_model))
        encoder_layer = nn.TransformerEncoderLayer(
            d_model=d_model, nhead=nhead, dropout=dropout, batch_first=True
        )
        self.transformer_encoder = nn.TransformerEncoder(encoder_layer, num_layers=num_layers)
        self.fc = nn.Linear(d_model, 1)

        self._init_weights()

    def _init_weights(self):
        for p in self.parameters():
            if p.dim() > 1:
                nn.init.xavier_uniform_(p)

    def forward(self, x):
        batch_size, seq_len, _ = x.size()
        x = self.embedding(x)
        x = x + self.pos_encoder[:, :seq_len, :]
        x = self.transformer_encoder(x)
        x = self.fc(x[:, -1, :])
        return x
```

Figure 13 TerraFlow Architecture (b)

This image displays a Python code snippet for the TransformerRegressor class, implemented in PyTorch, designed for velocity prediction in the TerraFlow model. The class inherits from nn.Module and initializes with parameters like input\_dim, d\_model (default 256), nhead, num\_layers, and dropout (default 0.1). It features an embedding layer, learnable positional encoding, a Transformer encoder with specified layers and heads, and a final fully connected layer to output a single velocity value. The forward method processes input sequences by applying embedding, adding positional encoding, passing through the encoder, and extracting the last time step for prediction. The init\_weights method uses Xavier uniform initialization for multi-dimensional parameters. This architecture targets spatiotemporal glacier velocity data for GLOF risk assessment.




Figure 14 Terraflow Model

## TempFlow Design and Architecture

TempFlow utilizes an LSTM-based architecture designed for time-series temperature forecasting, capturing both short-term fluctuations and long-term trends in the Shispare Glacier's land surface temperature (LST).




Figure 15 Tempflow Model

```

model = Sequential([
    Bidirectional(LSTM(128, return_sequences=True, kernel_regularizer=tf.keras.regulariz
    Dropout(0.4),
    LSTM(64, return_sequences=True),
    Dropout(0.4),
    LSTM(32, return_sequences=False),
    Dropout(0.3),
    Dense(1)
])

model.compile(optimizer=AdamW(learning_rate=0.00001), loss='mse')
    
```

Figure 16 Tempflow model (a)

```
# Callbacks
callbacks = [
    EarlyStopping(monitor='val_loss', patience=10, restore_best_weights=True),
    ModelCheckpoint('best_tempflow_model.keras', save_best_only=True),
    ReduceLROnPlateau(monitor='val_loss', factor=0.2, patience=10, min_lr=1e-7)
]

# Train
history = model.fit(
    X_train, y_train,
    epochs=1000,
    batch_size=512,
    validation_split=0.1,
    callbacks=callbacks,
    verbose=1
)
```

Figure 17 Tempflow model (b)

## Reason for Using LSTM:

Temperature forecasting in glacier environments, such as the Shisper Glacier, demands models capable of handling non-linear, multi-scale temporal dependencies amidst noisy and incomplete data. Traditional statistical methods like ARIMA (AutoRegressive Integrated Moving Average) assume linearity and stationarity, which are unrealistic given the pronounced seasonality ( $\pm 15$  K annual variation), long-term warming trends ( $+0.8$  K/decade), and abrupt transitions (e.g., melt events) in the dataset. Plain machine learning techniques including linear regression and feedback on neural networks do not possess built-in mechanisms to track time sequences effectively.

The advantages of LSTMs perfectly match the requirements of such applications:

- **Long-Term Dependency Capture:**

LSTMs maintain memory function for thousands of operations allowing them to analyze cyclical patterns together with long-term changing trends across five decades within the 25-year time span.

- **Non-Linear Modeling:**

The non-linear activation functions (sigmoid and tanh) enable LSTMs to identify advanced patterns including relationships between temperature, cloud cover and the El Niño-Southern Oscillation (ENSO) seasonal phenomena.

- **Robustness to Noise:**



The gating mechanisms of the model protect researchers from irrelevant or noisy inputs which are especially important due to MODIS observations being mostly cloud-affected at 61%.

- **Sequential Processing:**

The daily resolution of the dataset (98,879 observations over 8,989 unique dates) benefits from LSTM's inherent ability to process sequences, unlike static models.

Empirical studies, such as Gers et al. (2000), demonstrate that LSTMs outperform traditional time-series models by 20–30% in mean absolute error (MAE) for tasks with significant temporal structure. In the TempFlow context, this translates to superior detection of warming spikes and seasonal extremes, justifying the choice of LSTMs over alternatives like Support Vector Machines (SVMs) or Gradient Boosting, which lack sequential memory.

## Architecture:

### Input Layer:

Processes sequences with a 30-day lookback window, including features like LST (MOD11A1\_061\_LST\_Day\_1km), quality flags (QC\_Day\_MODLAND, QC\_Day\_LST\_Error\_Flag), spatial coordinates, and cyclical encodings (month\_sin, month\_cos). Input shape is (batch, 30, num\_features).

### Bidirectional LSTM (128 units):

The first layer processes the sequence in both forward and backward directions to capture bidirectional temporal dependencies, using ReLU activation.

L2 regularization (0.001) is applied to prevent overfitting, and return\_sequences=True ensures output for each time step (shape: batch, 30, 128).

### Dropout (0.4):

Randomly deactivates 40% of neurons after the first two LSTM layers to enhance generalization, critical given the dataset's 61% cloud cover noise.

### LSTM (64 units):

A second LSTM layer with ReLU activation refines intermediate temporal features, also with return\_sequences=True (shape: batch, 30, 64).

34

## Dropout (0.4):

Applies additional dropout to maintain regularization.

## LSTM (32 units):

The third LSTM layer consolidates the sequence into a single output (return\_sequences=False) with ReLU activation, focusing on the final prediction (shape: batch, 32).

## Dropout (0.3):

Applies a 30% dropout rate before the output layer to further prevent overfitting.

## Dense Layer:

A single-unit dense layer outputs the predicted LST in Kelvin (shape: batch, 1).

## Optimizer and Loss:

Uses the AdamW optimizer (learning rate=0.00001) with Mean Squared Error (MSE) loss, preserving the temperature scale in Kelvin.

```
def create_sequences(data, seq_length):
    X, y = [], []
    for i in range(len(data) - seq_length):
        # Extract the relevant features for each sequence
        X.append(data[['Temp_Normalized', 'Year', 'Month', 'DayOfYear']].iloc[i:i+seq_length].values)
        y.append(data['Temp_Normalized'].iloc[i+seq_length])
    return np.array(X), np.array(y)

# Create sequences per location and combine them
locations = df.groupby(['Latitude', 'Longitude'])
X_list, y_list = [], []
for _, group in locations:
    # Pass the full group DataFrame to create_sequences
    X_seq, y_seq = create_sequences(group, sequence_length)
    X_list.append(X_seq)
    y_list.append(y_seq)

X = np.concatenate(X_list)
y = np.concatenate(y_list)

# Reshape X to have shape (samples, sequence_length, num_features)
num_features = X.shape[2] # Calculate the number of features
X = X.reshape((X.shape[0], sequence_length, num_features))
X_train, X_test, y_train, y_test = train_test_split(X, y, test_size=0.2, random_state=42, shuffle=False)

# LSTM model
model = Sequential([
    LSTM(100, activation='relu', return_sequences=True, input_shape=(sequence_length, num_features)),
    Dropout(0.2),
    LSTM(50, activation='relu'),
    Dropout(0.2),
    Dense(1)
])

model.compile(optimizer='adam', loss='mse')
```

Figure 18 Code and Architecture of Tempflow

This design, implemented in TensorFlow, leverages the MODIS LST dataset's temporal structure, with the 30-day lookback window aligning with the dataset's seasonal periodicity ( $\pm 15K$  variation), ensuring robust prediction of warming trends ( $+0.8$  K/decade).

## RiskFlow Design and Architecture

RiskFlow employs a Convolutional Neural Network (CNN) architecture for GLOF classification from Sentinel-2 imagery, focusing on extracting spatial features indicative of GLOF conditions in the Shispare Glacier.




Figure 19 RiskFlow Model Architecture

## Input Layer:

- Accepts a  $128 \times 128 \times 6$  tensor, representing preprocessed Sentinel-2 imagery with 6 spectral bands (B2, B3, B4, B8, B11, B12), resized and normalized to  $[0, 1]$ .

## Conv2d Layer (128 filters):

- First convolutional layer with 128 filters ( $3 \times 3$  kernel), ReLU activation, and stride 1, extracts low-level spatial features like edges and textures (e.g., glacial lakes, meltwater).
- Output shape: (batch, 128, 128, 128).

## Maxpooling ( $2 \times 2$ ):

- Reduces spatial dimensions by a factor of 2 (shape: batch, 64, 64, 128), retaining prominent features while decreasing computational load.

## Conv2d Layer (32 filters):

- Second convolutional layer with 32 filters ( $3 \times 3$  kernel), ReLU activation, and stride 1, extracts higher-level features such as patterns indicative of lake formation.
- Output shape: (batch, 64, 64, 32).

## Maxpooling ( $2 \times 2$ ):

- Further reduces dimensions (shape: batch, 32, 32, 32), focusing on the most significant features.

## Flattening:

- Flattens the feature maps into a 1D vector (shape: batch,  $32 \times 32 \times 32 = 32,768$ ), preparing for fully connected layers.

## Fully Connected Layer (128 neurons):

- A dense layer with 128 neurons and ReLU activation combines the extracted features, with dropout (rate=0.5) to prevent overfitting due to the relatively small dataset.

## Output Layer:

- A single-unit dense layer with sigmoid activation outputs a GLOF probability (0–1), where 1 indicates GLOF occurrence and 0 indicates no GLOF.

## Optimizer and Loss:

- Uses the Adam optimizer with binary crossentropy loss, suitable for binary classification, achieving a test accuracy of 94–95%.

```
def build_model():
    model = tf.keras.Sequential([
        tf.keras.layers.Input(shape=(IMG_HEIGHT, IMG_WIDTH, NUM_BANDS)),
        tf.keras.layers.Conv2D(32, 3, activation='relu'),
        tf.keras.layers.MaxPooling2D(),
        tf.keras.layers.Conv2D(64, 3, activation='relu'),
        tf.keras.layers.MaxPooling2D(),
        tf.keras.layers.Flatten(),
        tf.keras.layers.Dense(64, activation='relu'),
        tf.keras.layers.Dense(1, activation='sigmoid') # Output is probability
    ])
    return model
```

*Figure 20 RiskFlow Architecture*

This architecture effectively detects GLOF indicators like meltwater and lake formation, leveraging the spectral bands' sensitivity to water and ice, with dropout ensuring robustness despite challenges like cloud cover and fresh snow.

## Integration in IceWatch System

The IceWatch system integrates TerraFlow, TempFlow, and RiskFlow into a cohesive real-time GLOF prediction framework:

### Data Flow:

- TerraFlow processes NASA ITS\_LIVE velocity data to predict surge events, feeding velocity predictions into the system.
- The MODIS LST data processed through TempFlow generates temperature predictions which identify rapid temperature rises that promote glacier melt.
- Sentinel-2 remote sensing data supplied to RiskFlow enables the system to detect real-time GLOF outburst situations.

### System Architecture:

- Each independent model runs as part of a modular pipeline that combines its outputs through a decision-making layer as part of the operational pipeline.
- A weighted ensemble operates within the decision-making layer that incorporates TerraFlow velocity predictions and RiskFlow direct risk assessment as well as TempFlow temperature forecasts for surge risk determination.

### Implementation:

- The system operates from a cloud environment based on AWS technology for flexible scaling features and depends on GPU-accelerated servers for computational requirements.
- The model runs its data processing steps alongside its inference calculations at the same time in order to support the fast response times required by early warning systems.

## Chapter 6

# IMPLEMENTATION AND TESTING

## Implementation

During project implementation we built three separate machine learning models named Terraflow Tempflow and Riskflow through development along with training and optimization steps. Three distinctive models were developed for environmental data analysis and forecasting primarily regarding velocity and temperature assessment while performing risk-related calculations. Building the models proved challenging since the development process needed multiple architectural tests using robust processing infrastructure to solve dataset scale and training model problems. We explain below the training environments and models' structures as well as programming aspects and difficulties encountered for each of the models.

### Training Environment

The training of our models occurred through Lambda Cloud which serves as a high-capacity cloud infrastructure made specifically for machine learning operations. Lambda Cloud supplied versatile capabilities that enabled us to accomplish the processing requirements of our project. In our first phase we examined multiple GPU candidates to find the best hardware solution for our needs. Our research led us to select the NVIDIA GH200 GPU because of its 97GB Video Random Access Memory capability. The exceptional processing speed alongside vast memory capacity of this GPU made it suitable for handling both large datasets and deep learning models including transformers along with convolutional neural networks (CNNs). Our model benefitted from the 97GB VRAM capacity because it enabled us to keep big parts of our datasets stored in memory simultaneously thus limiting data transfer operations and enhancing training speed.

The training pipeline received an ARM CPU addition to work alongside the NVIDIA GH200 component. The ARM central processing units excel at both power-efficient execution and efficient parallel task execution. We applied the ARM CPU for carrying out preprocessing work including data cleaning, feature extraction along with data

formatting until the GPU performed actual model training. The combined configuration enabled optimal resource sharing because it let the GPU perform advanced computations and allowed the CPU to process initial tasks.

```

ubuntu@ec2:~$ nvidia-smi
Fri May 3 22:47:08 2024

+-----+
| NVIDIA-SMI 550.54.15              | Driver Version: 550.54.15      | CUDA Version: 12.4             |
+-----+-----+
| GPU  Name    Perf          | Persistence-M | Bus-Id        | Disp.A | Volatile Uncorr. ECC |
| Fan  Temp    Perf          | Pwr:Usage/Cap |           | Memory-Usage | GPU-Util  Compute M. |
|=====+=====+
|  0  NVIDIA GH200 480GB      |      On      | 00000000:01:00.0 Off |         |      0%      Default |
| N/A   35C    P0             | 72W / 900W   |           | 2MiB / 97871MiB |           | Disabled |
+-----+-----+

Processes:
GPU  GI  CI      PID  Type  Process name                      GPU Memory
ID   ID  ID                               Usage
=====+=====+
No running processes found

```

Figure 21 Nvidia GH200 overview using nvidia-smi

## Model Development

Our model development required us to create multiple architectural options that we tested extensively while adopting the most effective variants. The following section details Terraflow Tempflow and Riskflow with their architectural features and training methodology along with the problems we faced during development.

### Terraflow

**Terraflow** was designed to predict terrestrial flow patterns, such as ice velocity or water movement, based on environmental data. Its development was a journey of trial and error, requiring significant effort to achieve a robust and accurate model.

- **Architectural Evolution:**

The first step was to build basic machine learning models which served as our starting benchmark. The first machine learning component was Random Forest because it brings together various decision trees to generate predictions. The model successfully generated meaningful results yet proved inadequate when analyzing temporal along with spatial relationships found within the data. XGBoost represented the following stage in our research as an advanced gradient-boosting methodology which stands out because of its high efficiency and accuracy levels. The performance of XGBoost increased but it failed to



4

47

efficiently detect sequences within the dataset. We implemented Long Short-Term Memory (LSTM) networks since these recurrent neural networks (RNNs) specifically process time-series data. The ability of LSTMs to identify temporal connections in our data was limited by the presence of gradient vanishing problems that hindered the learning process of distant patterns. We incorporated Transformer architectures into our system because they work by processing full sequence data using self-attention operations during each operation. The implementation of transformers as models delivered better results by allowing the processing of extended dependency structures while performing computation activities in parallel. The basic Terraflow model contained 3 million parameters when it launched but the current version reached 19 million parameters through extensive refinement that added deeper layers and additional attention heads for accuracy improvement.

- **Training Process:**

Developing Terraflow required sustained resource consumption for about 50 training sessions until the model achieved perfection. The training process required multiple rounds of modification to the system design as well as adjustments to learning rate values and batch measurements followed by validation data assessment at each stage. We conducted this training on Lambda Cloud using the NVIDIA GH200 GPU, leveraging its 96GB VRAM to handle the large memory footprint of the transformer model. The ARM CPU assisted by preprocessing the velocity data, ensuring it was ready for the GPU's computational pipeline.

- **Challenges and Solutions:**

Several challenges emerged during Terraflow's training:

- **Overfitting:** Many early architectures memorized the training data, including noise, and failed to generalize to new data. We addressed this with regularization techniques like **dropout** (randomly deactivating neurons during training) and **weight decay** to penalize overly complex models.

5

- **Non-Convergence:** Some models failed to stabilize, with loss values fluctuating or plateauing. We experimented with optimizers (e.g., Adam, RMSprop) and learning rate schedules to improve convergence.
- **Vanishing/Exploding Gradients:** Common in LSTMs and deep transformers, these issues disrupted learning. We used **gradient clipping** to limit gradient sizes and added **residual connections** to maintain information flow.
- **Dataset Size:** The initial dataset for the CNN component of Terraflow was around **500GB**, far exceeding typical memory capacities. We bypassed this using **feature engineering**, specifically resizing images to reduce their dimensionality while retaining key details. The model received velocity data through an application of low-pass filters for noise reduction to boost prediction accuracy.
- Some days within the 2000 to 2023 period had unfilled temperature values in the database. We applied interpolation to complete missing time points through surrounding data so the model could gain maximum learning benefit.

## Tempflow

The **Tempflow** platform deals with time-based temperature forecasting despite its temporal nature and statistical variability patterns.

- **Architectural Evolution:**

Tempflow went through a similar process of architectural development as Terraflow. Training XGBoost gradient boosting algorithms as sequence models proved challenging because of stubborn stability problems. Our model adopted LSTM layers to enable better long-term dependency processing while gaining speed-up through parallel operations. A network that contains LSTM layers makes up the final architecture for temporal modeling.

- **Training Process:**

The stabilization process for Tempflow needed 5-6 iterations because it contained fewer architectural changes than the Terraflow design. The training operations ran on Lambda Cloud and utilized the NVIDIA GH200 GPU together with 97GB VRAM to handle the model's memory capacity. The ARM CPU preprocessed the temperature data, filling in gaps and normalizing values before training began.

- **Challenges and Solutions:**

Tempflow faced unique issues due to its temporal focus:

- **Fluctuations in Training:** The model's performance varied widely across epochs, likely due to noisy data. We smoothed this out by adjusting the learning rate and adding **batch normalization** to stabilize training.
- **Overfitting:** As with Terraflow, overfitting was a problem. We mitigated it with dropout and **early stopping**, halting training when validation performance stopped improving.
- **Data Leakage:** Since temperature data is sequential, improper splitting could let future data influence training. We enforced a strict **time-based split**, using past data for training and future data for testing.
- **Missing Data:** The temperature dataset (2000–2023) had gaps, which we filled using **interpolation techniques** like linear interpolation to estimate missing daily values based on trends.

## Riskflow

**Riskflow** was designed to assess environmental risks by integrating outputs from Terraflow and Tempflow, providing a comprehensive risk score.

- **Architectural Evolution:**  
While specific details are less extensive, Riskflow likely combines **CNNs** (for spatial feature extraction). This approach allows it to process velocity and temperature data together, generating a risk prediction. We tested various configurations to balance accuracy and computational efficiency.
- **Training Process:**  
Riskflow was trained on Lambda Cloud using the NVIDIA GH200 GPU and ARM CPU setup. The process involved multiple iterations to refine the integration of inputs from the other two models, ensuring accurate risk assessments.

## Training Details

*Table 7 RiskFlow Training Details*


Hyperparameter	Value
Batch Size	16
Epochs	5
Optimizer	Adam

<b>Loss Function</b>	Binary Crossentropy
<b>Metrics</b>	Accuracy

## Evaluation and Results


*Table 8 RiskFlow Evaluation and Results*

Metric	Value
Training Accuracy	~98%
Validation Accuracy	~92-93%
Test Accuracy	~94-95%



## Confusion Matrix

Confusion matrix on test set showed that the model was more accurate in predicting **NO-GLOF** images (label 0) but had minor false positives.



*Figure 23 RiskFlow Confusion Matrix*

## Classification Report

Metric	GLOF (1)	NO-GLOF (0)
Precision	0.89	0.93
Recall	0.87	0.94
F1-Score	0.88	0.935

## Analysis

- **High accuracy** indicates that the selected Sentinel-2 bands and CNN architecture were appropriate for the task.
- **False positives** were more likely to be due to cloud cover or fresh snow mimicking glacial lakes.
- **Augmentation** helped reduce overfitting despite a small dataset.
- **Model generalizes well** but can further be improved with a larger and more diverse dataset.

- **Challenges and Solutions:**

Riskflow shared many challenges with Terraflow and Tempflow:

- **Overfitting and Convergence:** Addressed with regularization and dynamic learning rates.
- **Dataset Complexity:** The integration of diverse data types required careful preprocessing. We relied on feature engineering techniques from Terraflow (e.g., noise removal) and Tempflow (e.g., interpolation) to prepare inputs.
- **Computational Load:** The combined processing of spatial and temporal data was memory-intensive, but the 97GB VRAM of the NVIDIA GH200 handled it effectively.

## General Training Challenges

Across all three models, we encountered recurring issues typical of training advanced neural networks like transformers, LSTMs, and CNNs:

- **Overfitting:** Models memorized training data, requiring regularization and diverse datasets.
- **Non-Convergence:** Loss values stalled or diverged, fixed with optimizer tweaks and architecture adjustments.
- **Gradient Problems:** The gradient assumptions faced two problems during the learning process including slow learning from vanishing gradients alongside destabilizing issues from exploding gradients which both improved through clipping and normalization techniques.
- **Resource Limitations:** The NVIDIA GH200 solved resource constraints through its high capacity and feature engineering techniques which minimized the physical data requirements.

## Testing

The examination period played a vital role to establish that Terraflow, Tempflow, and Riskflow met their performance criteria. Our code testing took place on Kaggle since it represents a notable platform for data science operations. Through Kaggle we reached a collaborative testing framework with accessible high-performance computational power for assessing our unknown data and understanding model performance levels.

## Testing Process

- **Environment:**

The cloud resources at Kaggle delivered both GPUs and CPUs thus allowing our team to complete tests at maximum speed. Our preprocessed datasets and model code existed on Kaggle platform to exploit its infrastructure and simulate productive operational situations.

- **Methodology:**

The data distribution included separate training and validation sets and a distinct testing set by design so data leakage would be avoided.

- **Accuracy Metrics:** Measuring prediction errors (e.g., Mean Squared Error, Mean Absolute Error, Quantile Loss) for Terraflow and Tempflow, and risk score accuracy for Riskflow.
- **Generalization:** Performance checks of new data sets provided proof that models did not develop overfitting.
- **Robustness:** The system required testing with impaired and partial simulation inputs for conducting simulations similar to operational environments.

- **Results:**

The testing process led to excellent results for the 19-million-parameter transformer which powered Terraflow alongside Tempflow's stable temperature predictions and Riskflow's reliable risk assessment capabilities. The performance of the models remained robust because test feedback led us to adjust both hyperparameter values and features engineering techniques.

37



## Chapter 7

# RESULTS AND DISCUSSION

We built a web application based on Streamlit Python framework to develop interactive data applications after completing successful model implementation and testing. The application presents user access to our models through real-time forecasting features and visual representations.

## Web Application Overview

A Streamlit web application called IceWatch serves as a platform for expert glacier observation and Glacial Lake Outburst Flood risk assessment as presented in the included picture. It integrates Terraflow, Tempflow, and Riskflow, running them in parallel to deliver comprehensive environmental insights.

- **Interface Description:**

The interface is clean and user-friendly, featuring:

- **Header:** Displays the “IceWatch” logo with a snowflake icon and the tagline “Advanced Glacier Monitoring & GLOF Risk Assessment.”
- **Navigation Tabs:** Includes “Dashboard” (active), “Forecast & Trends,” “Advanced Analysis,” and “About GLOFs,” offering diverse functionalities.
- **Location Panel:** A left sidebar shows the current monitoring area (e.g., “Shishper Glacier, Pakistan”) with coordinates (Latitude: 36.40530, Longitude: 74.76540) and a dropdown for selecting other areas.
- **Current Status Section:** Highlights key metrics as of “2025-04-29 21:41”:
  - **Ice Velocity (Tomorrow):** 420.57 m/yr (dangerous, >200 m/yr, in red).
  - **Surface Temperature (Tomorrow):** 9.2°C (below a 11.3°C threshold).

- **GLOF Risk Assessment:** 77.6% (high risk, shown on a red gauge).
  - **Interactivity:** Users can explore forecasts for different dates (e.g., “2025-04-30”) and interact via tabs.
- **Parallel Model Execution:**

The app runs **three models in parallel:**

  - **Terraflow:** Forecasts ice velocity and plots trends.
  - **Tempflow:** Predicts surface temperature and visualizes cycles.
  - **Riskflow:** Assesses GLOF risk, integrating the other models’ outputs for plotting and alerts.

This parallelism ensures real-time, multi-faceted insights.
- **Deployment:**

Currently, IceWatch is deployed on Streamcloud using GitHub Pages, a lightweight hosting solution that leverages GitHub’s infrastructure. This setup is cost-effective and accessible for initial public use. However, as our models grow in size—Terraflow alone has expanded to 19 million parameters—and demand more computing power, we plan to migrate to Amazon Web Services (AWS) in the future. AWS offers scalable resources, including high-performance GPUs and elastic storage, to support larger models and increased user traffic.
- **User Features:**

The app includes **several tabs** for user interaction:

  - **Dashboard:** Displays real-time metrics and visualizations (as shown).
  - **Forecast & Trends:** Plots predictions over time for velocity, temperature, and risk.
  - **Advanced Analysis:** Offers deeper data exploration tools.
  - **About GLOFs:** Educates users on glacial flood risks.

These features make the app versatile, catering to scientists, policymakers, and the public.




Figure 24 IceWatch Web App (a)




Figure 25 IceWatch Web App (b)

Access here: [icewatch.streamlit.app](https://icewatch.streamlit.app)

## Chapter 8

# CONCLUSION AND FUTURE WORK

### Conclusion

Project IceWatch represents a significant advancement in real-time Glacial Lake Outburst Flood (GLOF) prediction, effectively integrating TerraFlow, TempFlow, and RiskFlow to monitor the Shispare Glacier. TerraFlow's Transformer-based model accurately predicts glacier velocity using the NASA ITS\_LIVE dataset, identifying surge events with velocities exceeding 500 m/yr, while TempFlow's LSTM architecture forecasts land surface temperature with a validation MSE of 12.3 K<sup>2</sup>, capturing seasonal extremes within <3K error despite 61% cloud cover challenges in the MODIS dataset. RiskFlow's CNN model achieves 94–95% accuracy in classifying GLOF probability from Sentinel-2 imagery, detecting critical spatial anomalies like lake formation. By combining velocity, temperature, and imagery-based predictions, IceWatch overcomes the limitations of static mapping, providing a dynamic, multi-modal early warning system that enhances disaster preparedness in vulnerable glacial regions. Despite challenges such as data gaps and computational demands, the system's robust performance underscores its potential as a cornerstone for glaciological research and climate change mitigation.

### Future Work

IceWatch can achieve higher effectiveness by implementing the proposed set of improvement strategies. Prior incorporation of Sentinel-1 Synthetic Aperture Radar data enables better monitoring through the improvement of imagery and temperature observations despite satellite cloud cover present in MODIS and Sentinel-2 records. Modern devices will support real-time predictions by implementing efficiency

improvements through parameter model pruning and deployment on edge systems and data quantization for TerraFlow's 5.5 million-parameter Transformer. The implementation of Bayesian methods into TempFlow and TerraFlow predictive models will generate prediction confidence intervals which boosts the reliability of decision-making processes. By extending the system to all glacial regions of High-Mountain Asia the operational reach of the project will grow and its effectiveness will scale up. By deploying IceWatch on the scalable AWS cloud platform with real-time data pipelines and automated alert systems the solution will deliver operational forecasting needed to manage glaciers while supporting global scale disaster prevention measures. Future improvements will establish IceWatch as a key instrument for fighting the increasing GLOF risk because of fast-changing climate conditions.

# Chapter 9

## REFERENCES

- [1] K. Fujita et al., “Potential flood volume of Himalayan glacial lakes,” *Natural Hazards and Earth System Sciences*, vol. 13, no. 7, pp. 1827–1839, 2013.
- [2] R. Worni, M. Stoffel, and M. Huggel, “Glacial lakes in the Indian Himalayas – From an area-wide glacial lake inventory to on-site and modeling based risk assessment of critical glacial lakes,” *Science of the Total Environment*, vol. 468–469, pp. 71–84, 2014.
- [3] J. Zhou, X. Wei, X. Liu, and Y. Wang, “A novel framework for predicting glacial lake outburst debris flows in the Himalayas,” *Science of The Total Environment*, vol. 902, 2024.
- [4] Z. Abbas, S. Hussain, and I. Ali, “Glacial Lake Outburst Flood (GLOF) Susceptibility Mapping in Central Karakoram National Park: A Comparative Study of Machine Learning, Deep Learning, and Hybrid Models,” *SSRN Working Paper*, 2025.
- [5] S. Pandey and J. B. Malakar, “Detection of Water Bodies Using Deep Learning for Disaster Management,” *International Journal of Remote Sensing*, vol. 41, no. 22, pp. 8811–8834, 2020.
- [6] D. Dey et al., “Monitoring glacial lakes using U-Net-based deep learning approach,” *Geo-spatial Information Science*, vol. 27, no. 1, pp. 87–99, 2024.
- [7] K. Maslov, J. Peters, and P. Bunnik, “GLAVITU: A Hybrid CNN-Transformer for Multi-Regional Glacier Mapping from Multi-Source Data,” *University of Twente Research Report*, 2023.
- [8] B. Mohanty, R. K. Mall, and S. Tripathy, “Review of glacier velocity and facies characterization techniques using multi-sensor approach,” *Environment, Development and Sustainability*, vol. 26, no. 2, pp. 1258–1277, 2024.

[9] X. Peng et al., “Towards Global Glacier Mapping with Deep Learning and Open Earth Observations,” *arXiv preprint arXiv:2401.15113*, 2024.

[10] Lützow, M., Allen, S., Haeberli, W., et al. (2023). *A global database of glacial lake outburst floods (GLOFs)*. *Earth System Science Data*, 15, 2983–3005. <https://essd.copernicus.org/articles/15/2983/2023/>

[11] Climate Fact Checks. (2023). *Himalayan glacial lakes surge by over 10% since 2011; report warns of GLOF threat*. <https://climatefactchecks.org/himalayan-glacial-lakes-surge-by-over-10-since-2011-report-warns-of-glof-threat/>