

School of Engineering and Applied Science (SEAS), Ahmedabad University

Probability and Stochastic Processes (MAT277)

Homework Assignment-4

Enrollment No: AU2140096

Name: Ansh Virani

1.

2. A random variable X is uniformly distributed over the interval $(0, 1)$ and related to Y by,

$$\tan\left(\frac{\pi Y}{2}\right) = e^X \implies Y = \frac{2}{\pi} \arctan(e^X)$$

$$\therefore \frac{dY}{dX} = \frac{2}{\pi} \cdot \frac{1}{1 + e^{2X}}$$

Applying the transformation rule, we get:

$$f_Y(y) = f_X(x) \left| \frac{dY}{dX} \right| = 1 \times \frac{2}{\pi} \cdot \frac{1}{1 + e^{2X}}$$

Since X is expressed in terms of Y through the initial transformation, $e^X = \tan\left(\frac{\pi Y}{2}\right)$, the *PDF* can be expressed in terms of Y as follows:

$$f_Y(y) = \left(\frac{2}{\pi}\right) \left(\frac{1}{1 + \tan^2\left(\frac{\pi y}{2}\right)} \right)$$

Using the identity $1 + \tan^2(z) = \sec^2(z)$, we get:

$$f_Y(y) = \left(\frac{2}{\pi}\right) \left(\frac{1}{\sec^2\left(\frac{\pi y}{2}\right)} \right) = \frac{2}{\pi} \cos^2\left(\frac{\pi y}{2}\right)$$

By solving for Y , computing the derivative with respect to X , and applying the transformation rule, the resulting *PDF* for Y is $f_Y(y) = \frac{2}{\pi} \cos^2\left(\frac{\pi y}{2}\right)$, valid for y in the interval $(0, 1)$.