

Permutations and Combinations: Takeaways



by Dataquest Labs, Inc. - All rights reserved © 2020

Concepts

- If we have an experiment E_1 (like flipping a coin) with a outcomes, followed by an experiment E_2 (like rolling a die) with b outcomes, then the total number of outcomes for the composite experiment E_1E_2 can be found by multiplying a with b (this is known as the **rule of product**):
- If we have an experiment E_1 with a outcomes, followed by an experiment E_2 with b outcomes, followed by an experiment E_n with z outcomes, the total number of outcomes for the composite experiment $E_1E_2 \dots E_n$ can be found by multiplying their individual outcomes:
- There are two kinds of arrangements:
 - Arrangements where the order matters, which we call **permutations**.
 - Arrangements where the order doesn't matter, which we call **combinations**.
- To find the number of permutations when we're sampling without replacement, we can use the formula:
- To find the number of permutations when we're sampling without replacement and taking only k objects from a group of n objects, we can use the formula:
- To find the number of combinations when we're sampling without replacement and taking only k objects from a group of n objects, we can use the formula:

Resources

- [A tutorial on calculating combinations when sampling with replacement](#), which we haven't covered in this mission

- [An easy-to-digest introduction to permutations and combinations](#)



Takeaways by Dataquest Labs, Inc. - All rights reserved © 2020