

Random Numbers

Anshul Sangrame
CS21BTECH11004

CONTENTS

1	Uniform Random Numbers	1
2	Central Limit Theorem	2
3	From Uniform to Other	4
4	Triangular Distribution	4
5	Maximum Likelihood	6
6	Gaussian to Other	9

1 UNIFORM RANDOM NUMBERS

Let U be a uniform random variable between 0 and 1.

- 1.1 Generate 10^6 samples of U using a C program and save into a file called uni.dat .

Solution:

Download the following files,

```
wget https://github.com/Anshul-Sangrame/
AI1110/blob/main/Assignment/solution/1/
coeffs.h
wget https://github.com/Anshul-Sangrame/
AI1110/blob/main/Assignment/solution
/1/1_1.c
```

Execute the above files using code,

```
gcc 1_1.c -lm
./a.out
```

- 1.2 Load the uni.dat file into python and plot the empirical CDF of U using the samples in uni.dat. The CDF is defined as

$$F_U(x) = \Pr(U \leq x) \quad (1.1)$$

Solution:

Download the following files

```
wget https://github.com/Anshul-Sangrame/
AI1110/blob/main/Assignment/solution
/1/1_2.py
```

Execute the code using command

```
python3 1_2.py
```

Plot 1.1 is obtained,




Fig. 1.1

- 1.3 Find a theoretical expression for $F_U(x)$.

Solution:

Pdf of Uniform distribution between $[0,1]$ is given by,

$$f_U(x) = \begin{cases} 1, & x \in [0, 1] \\ 0, & \text{otherwise} \end{cases} \quad (1.2)$$

$$F_U(x) = \int_{-\infty}^x f_U(x) dx \quad (1.3)$$

Case-1: $x < 0$,

$$F_U(x) = \int_{-\infty}^x 0 dx \quad (1.4)$$

$$= 0 \quad (1.5)$$

Case-2: $x \in [0,1]$,

$$F_U(x) = \int_{-\infty}^0 0dx + \int_0^x 1dx \quad (1.6)$$

$$= x \quad (1.7)$$

Case-3: $x > 1$,

$$F_U(x) = \int_{-\infty}^0 0dx + \int_0^1 1dx + \int_1^x 0dx \quad (1.8)$$

$$= 1 \quad (1.9)$$

Hence,

$$F_U(x) = \begin{cases} 0, & x < 0 \\ x, & x \in [0, 1] \\ 1, & x > 1 \end{cases} \quad (1.10)$$

1.4 The mean of U is defined as

$$E[U] = \frac{1}{N} \sum_{i=1}^N U_i \quad (1.11)$$

and its variance as

$$\text{var}[U] = E[U - E[U]]^2 \quad (1.12)$$

Write a C program to find the mean and variance of U .

Solution:

Download the following files,

```
wget https://github.com/Anshul-Sangrame/
AI1110/blob/main/Assignment/solution/1/
coeffs.h
wget https://github.com/Anshul-Sangrame/
AI1110/blob/main/Assignment/solution
/1/1_4.c
```

Execute the above files using code,

```
gcc 1_4.c -lm
./a.out
```

The following result is obtained

$$E[U] = 0.500007$$

$$\text{var}[U] = 0.083301$$

1.5 Verify your result theoretically given that

$$E[U^k] = \int_{-\infty}^{\infty} x^k dF_U(x) \quad (1.13)$$

Solution:

$F_U(x)$ for uniform distribution,

$$F_U(x) = \begin{cases} 0, & x < 0 \\ x, & x \in [0, 1] \\ 1, & x > 1 \end{cases} \quad (1.14)$$

$$E[U] = \int_{-\infty}^{\infty} x dF_U(x) \quad (1.15)$$

$$= \int_{-\infty}^0 0dx + \int_0^1 xdx + \int_1^{\infty} 0dx \quad (1.16)$$

$$= \frac{1}{2} \quad (1.17)$$

$$E[U^2] = \int_{-\infty}^{\infty} x^2 dF_U(x) \quad (1.18)$$

$$= \int_{-\infty}^0 0dx + \int_0^1 x^2 dx + \int_1^{\infty} 0dx \quad (1.19)$$

$$= \frac{1}{3} \quad (1.20)$$

$$E[U - E[U]]^2 = E[U^2] - [E[U]]^2 \quad (1.21)$$

From Equation (1.17) and (1.20),

$$E[U - E[U]]^2 = \frac{1}{3} - \left(\frac{1}{2}\right)^2 \quad (1.22)$$

$$= \frac{1}{3} - \frac{1}{4} \quad (1.23)$$

$$= \frac{1}{12} \approx 0.083 \quad (1.24)$$

2 CENTRAL LIMIT THEOREM

2.1 Generate 10^6 samples of the random variable

$$X = \sum_{i=1}^{12} U_i - 6 \quad (2.1)$$

using a C program, where $U_i, i = 1, 2, \dots, 12$ are a set of independent uniform random variables between 0 and 1 and save in a file called gau.dat

Solution:

Download the following files,

```
wget https://github.com/Anshul-Sangrame/
AI1110/blob/main/Assignment/solution/1/
coeffs.h
wget https://github.com/Anshul-Sangrame/
AI1110/blob/main/Assignment/solution
/2/2_1.c
```

Execute the above files using code,

```
gcc 2_1.c -lm
./a.out
```

- 2.2 Load gau.dat in python and plot the empirical CDF of X using the samples in gau.dat. What properties does a CDF have?

Solution:

Download the following files,

```
wget https://github.com/Anshul-Sangrame/
AI1110/blob/main/Assignment/solution
/2/2_2.py
```

Execute the above file using code,

```
python3 2_2.py
```

Plot 2.1 obtained is symmetric about (0,0.5)




Fig. 2.1

- 2.3 Load gau.dat in python and plot the empirical PDF of X using the samples in gau.dat. The PDF of X is defined as

$$p_X(x) = \frac{d}{dx} F_X(x) \quad (2.2)$$

What properties does the PDF have?

Solution:

Download the following files,

```
wget https://github.com/Anshul-Sangrame/
AI1110/blob/main/Assignment/solution
/2/2_3.py
```

Execute the above file using code,

```
python3 2_3.py
```




Fig. 2.2

Plot 2.2 obtained is symmetric about y-axis

- 2.4 Find the mean and variance of X by writing a C program.

Solution:

Download the following files,

```
wget https://github.com/Anshul-Sangrame/
AI1110/blob/main/Assignment/solution/1/
coeffs.h
wget https://github.com/Anshul-Sangrame/
AI1110/blob/main/Assignment/solution
/2/2_4.c
```

Execute the above files using code,

```
gcc 2_4.c -lm
./a.out
```

The result is,

$$E[U] = 0.000294 \quad (2.3)$$

$$\text{var}[U] = 0.999561 \quad (2.4)$$

- 2.5 Given that

$$p_X(x) = \frac{1}{\sqrt{2\pi}} \exp\left(-\frac{x^2}{2}\right), -\infty < x < \infty, \quad (2.5)$$

repeat the above exercise theoretically.

Solution:

$$E[U] = \int_{-\infty}^{\infty} u p_X(u) du \quad (2.6)$$

$$(2.7)$$

$up_X(u)$ is an odd function.

$$E[U] = 0 \quad (2.8)$$

$$(2.9)$$

$$E[U^2] = \int_{-\infty}^{\infty} \frac{u^2}{\sqrt{2\pi}} \exp\left(-\frac{u^2}{2}\right) du \quad (2.10)$$

$$= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} u \left(u \exp\left(-\frac{u^2}{2}\right) \right) du \quad (2.11)$$

Using integration by parts,

$$E[U^2] = -u \frac{1}{\sqrt{2\pi}} \exp\left(-\frac{u^2}{2}\right) \Big|_{-\infty}^{\infty} + \int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi}} \exp\left(-\frac{u^2}{2}\right) du \quad (2.12)$$

$$= 0 + \frac{1}{\sqrt{2\pi}} \sqrt{2\pi} \quad (2.13)$$

$$= 1 \quad (2.14)$$

Hence,

$$\text{var}[U] = E[U^2] - [E[U]]^2 \quad (2.15)$$

$$= 1 \quad (2.16)$$

3 FROM UNIFORM TO OTHER

3.1 Generate samples of

$$V = -2 \ln(1 - U) \quad (3.1)$$

and plot its CDF.

Solution:

Download the following files,

```
wget https://github.com/Anshul-Sangrame/
AI1110/blob/main/Assignment/solution/
coeffs.h
wget https://github.com/Anshul-Sangrame/
AI1110/blob/main/Assignment/solution
/3/3_1.c
wget https://github.com/Anshul-Sangrame/
AI1110/blob/main/Assignment/solution
/3/3_1.py
```

Execute the above files using code,

```
gcc 3_1.c -lm
./a.out
python3 3_1.py
```

Plot 3.1 is obtained

3.2 Find a theoretical expression for $F_V(x)$.




Fig. 3.1

Solution:

$$F_V(x) = \Pr(V \leq x) \quad (3.2)$$

$$= \Pr(-2 \ln(1 - U) \leq x) \quad (3.3)$$

$$= \Pr(U \leq 1 - e^{-\frac{x}{2}}) \quad (3.4)$$

$$F_V(x) = F_U(1 - e^{-\frac{x}{2}}) \quad (3.5)$$

$$= \begin{cases} 0, & 1 - e^{-\frac{x}{2}} < 0 \\ 1 - e^{-\frac{x}{2}}, & 1 - e^{-\frac{x}{2}} \in [0, 1] \\ 1, & 1 - e^{-\frac{x}{2}} > 1 \end{cases} \quad (3.6)$$

Now,

$$1 - e^{-\frac{x}{2}} < 0 \quad (3.7)$$

$$\Rightarrow x < 0 \quad (3.8)$$

$$0 \leq 1 - e^{-\frac{x}{2}} \leq 1 \quad (3.9)$$

$$\Rightarrow x \geq 0 \quad (3.10)$$

$$1 < 1 - e^{-\frac{x}{2}} \quad (3.11)$$

$$\Rightarrow x \in \phi \quad (3.12)$$

Hence,

$$F_V(x) = \begin{cases} 0, & x < 0 \\ 1 - e^{-\frac{x}{2}}, & x \geq 0 \end{cases} \quad (3.13)$$

4 TRIANGULAR DISTRIBUTION

4.1 Generate

$$T = U_1 + U_2 \quad (4.1)$$

Solution:

Download the following files,

```
wget https://github.com/Anshul-Sangrame/
AI1110/blob/main/Assignment/solution/
coeffs.h
wget https://github.com/Anshul-Sangrame/
AI1110/blob/main/Assignment/solution
/4/4_1.c
```

Execute the above files using code,

```
gcc 4_1.c -lm
./a.out
```

4.2 Find the CDF of T .**Solution:**

Download the following files,

```
wget https://github.com/Anshul-Sangrame/
AI1110/blob/main/Assignment/solution
/4/4_2.py
```

Execute the above files using code,

```
python3 4_2.py
```

Plot 4.1 is obtained




Fig. 4.1

4.3 Find the PDF of T .**Solution:**

Download the following files,

```
wget https://github.com/Anshul-Sangrame/
AI1110/blob/main/Assignment/solution
/4/4_3.py
```

Execute the above files using code,

```
python3 4_3.py
```

Plot 4.2 is obtained




Fig. 4.2

4.4 Find the theoretical PDF and CDF of T .**Solution:**

The CDF of T is given by

$$F_T(t) = \Pr(T \leq t) = \Pr(U_1 + U_2 \leq t) \quad (4.2)$$

Since $U_1, U_2 \in [0, 1] \implies U_1 + U_2 \in [0, 2]$
Therefore, if $t \geq 2$, then $U_1 + U_2 \leq t$ is always true and if $t < 0$, then $U_1 + U_2 \leq t$ is always false.

Now, fix the value of U_1 to be some x

$$x + U_2 \leq t \implies U_2 \leq t - x \quad (4.3)$$

If $0 \leq t \leq 1$, then x can take all values in $[0, t]$

$$F_T(t) = \int_0^t \Pr(U_2 \leq t - x) p_{U_1}(x) dx \quad (4.4)$$

$$= \int_0^t F_{U_2}(t - x) p_{U_1}(x) dx \quad (4.5)$$

$$0 \leq x \leq t \implies 0 \leq t - x \leq t \leq 1 \quad (4.6)$$

$$\implies F_{U_2}(t - x) = t - x \quad (4.7)$$

$$F_T(t) = \int_0^t (t - x) \cdot 1 \cdot dx \quad (4.8)$$

$$= tx - \frac{x^2}{2} \Big|_0^t \quad (4.9)$$

$$= \frac{t^2}{2} \quad (4.10)$$

If $1 < t < 2$, x can only take values in $[0, 1]$ as $U_1 \leq 1$

$$F_T(t) = \int_0^1 F_{U_2}(t-x) \cdot 1 \cdot dx \quad (4.11)$$

$$0 \leq x \leq t-1 \implies 1 \leq t-x \leq t \quad (4.12)$$

$$t-1 \leq x \leq 1 \implies 0 < t-1 \leq t-x \leq 1 \quad (4.13)$$

$$F_T(t) = \int_0^{t-1} 1 dx + \int_{t-1}^1 (t-x) dx \quad (4.14)$$

$$= t-1 + t(1-(t-1)) - \frac{1}{2} + \frac{(t-1)^2}{2} \quad (4.15)$$

$$= t-1 + 2t-t^2 - \frac{1}{2} + \frac{t^2}{2} + \frac{1}{2} - t \quad (4.16)$$

$$= -\frac{t^2}{2} + 2t - 1 \quad (4.17)$$

Therefore,

$$F_T(t) = \begin{cases} 0 & t < 0 \\ \frac{t^2}{2} & 0 \leq t \leq 1 \\ 2t - \frac{t^2}{2} - 1 & 1 < t < 2 \\ 1 & t \geq 2 \end{cases} \quad (4.18)$$

The PDF of T is given by

$$p_T(t) = \frac{d}{dt} F_T(t) \quad (4.19)$$

$$\therefore p_T(t) = \begin{cases} 0 & t < 0 \\ t & 0 \leq t \leq 1 \\ 2-t & 1 < t < 2 \\ 0 & t \geq 2 \end{cases} \quad (4.20)$$

4.5 Verify your results through a plot.

Solution:

Download the following files,

```
wget https://github.com/Anshul-Sangrame/
AI1110/blob/main/Assignment/solution
/4/4_5_1.py
wget https://github.com/Anshul-Sangrame/
AI1110/blob/main/Assignment/solution
/4/4_5_2.py
```

Execute the above files using code,

```
python3 4_5_1.py
```

```
python3 4_5_2.py
```

Plot 4.3 is obtained is pdf Plot 4.4 is obtained




Fig. 4.3

is cdf




Fig. 4.4

5 MAXIMUM LIKELIHOOD

5.1 Generate equiprobable $X \in \{1, -1\}$.

Solution:

Download the following files,

```
wget https://github.com/Anshul-Sangrame/
AI1110/blob/main/Assignment/solution
/5/5.1.c
wget https://github.com/Anshul-Sangrame/
AI1110/blob/main/Assignment/solution/
coeffs.h
```

Execute the above files using code,

```
gcc 5.1.c -lm
./a.out
```

5.2 Generate

$$Y = AX + N \quad (5.1)$$

where $A = 5$ dB, $X \in \{1, -1\}$ is Bernoulli and $N \sim \mathcal{N}(0, 1)$.

Solution:

Download the following files,

```
wget https://github.com/Anshul-Sangrame/
AI1110/blob/main/Assignment/solution
/5/5.2.c
wget https://github.com/Anshul-Sangrame/
AI1110/blob/main/Assignment/solution/
coeffs.h
```

Execute the above files using code,

```
gcc 5.2.c -lm
./a.out
```

5.3 Plot Y using a scatter plot.

Solution:

Download the following files,

```
wget https://github.com/Anshul-Sangrame/
AI1110/blob/main/Assignment/solution
/5/5.3.py
```

Execute the above files using code,

```
python3 5.3.py
```

Plot 5.1 is obtained




Fig. 5.1

5.4 Guess how to estimate X from Y .

Solution:

From the plot of Y , we see that the estimate model can be written as

$$\hat{X} = \begin{cases} 1 & Y > 0 \\ 0 & Y < 0 \end{cases} \quad (5.2)$$

5.5 Find

$$P_{e|0} = \Pr(\hat{X} = -1 | X = 1) \quad (5.3)$$

and

$$P_{e|1} = \Pr(\hat{X} = 1 | X = -1) \quad (5.4)$$

Solution:

Letting $X = 1$ and $X = -1$ respectively, we see the number of mismatched data points to compute the error probabilities. The simulation is coded in

```
$ wget https://github.com/Anshul-Sangrame/
AI1110/blob/main/Assignment/solution
/5/5.5.py
```

and can be run by typing

```
$ python3 5.5.py
```

The results are

$$P_{e|0} = 5.02 \times 10^{-4} \quad (5.5)$$

$$P_{e|1} = 5.24 \times 10^{-4} \quad (5.6)$$

5.6 Find P_e assuming that X has equiprobable symbols.

Solution:

Here, $\Pr(X = 1) = \Pr(X = -1) = 0.5$. Thus,

$$P_e = \Pr(X = 1) P_{e|1} + \Pr(X = -1) P_{e|0} \quad (5.7)$$

$$= \frac{1}{2} (P_{e|0} + P_{e|1}) = 5.13 \times 10^{-4} \quad (5.8)$$

5.7 Verify by plotting the theoretical P_e wrt A from 0 dB to 10 dB.

Solution:

$$P_{e|0} = \Pr(\hat{X} = 1 | X = -1) \quad (5.9)$$

$$= \Pr(Y > 0 | X = -1) \quad (5.10)$$

$$= \Pr(AX + N > 0 | X = -1) \quad (5.11)$$

$$= \Pr(N > A) = Q(A) \quad (5.12)$$

since X and N are independent. Writing a

similar expression for $P_{e|1}$ and noting that

$$\Pr(N < -A) = \Pr(N > A) = Q(A) \quad (5.13)$$

it follows that $P_e = Q(A)$. This is the idea used to plot the theoretical P_e . The plot is coded both in the rectangular axes and the semilog-y axes. Download the relevant codes using

```
wget https://github.com/Anshul-Sangrame/
AI1110/blob/main/Assignment/solution
/5/5.7.py
wget https://github.com/Anshul-Sangrame/
AI1110/blob/main/Assignment/solution
/5/5.7_semilog.py
```

Execute the above files using code,

```
python3 5.7.py
python3 5.7_semilog.py
```

Plot 5.2 obtained in rectangular axis




Fig. 5.2

Plot 5.3 obtained in semilog-y axis

5.8 Now, consider a threshold δ while estimating X from Y . Find the value of δ that maximizes the theoretical P_e .

Solution:

To estimate X from Y , we now consider the following:

$$X = \begin{cases} 1, & Y > \delta \\ -1, & Y < \delta \end{cases} \quad (5.14)$$




Fig. 5.3

Therefore,

$$P_e = \Pr(X = -1) Q(A + \delta) + \Pr(X = 1) Q(A - \delta) \quad (5.15)$$

$$= \frac{1}{2} (Q(A + \delta) + Q(A - \delta)) \quad (5.16)$$

To minimise P_e , we differentiate the above equation wrt δ :

$$0 = \frac{d}{d\delta} \left(\frac{1}{2} (Q_N(A - \delta) + Q_N(A + \delta)) \right) \quad (5.17)$$

$$= \frac{1}{2} \left(\frac{1}{\sqrt{2\pi}} e^{-\frac{(\delta-A)^2}{2}} - \frac{1}{\sqrt{2\pi}} e^{-\frac{(A+\delta)^2}{2}} \right) \quad (5.18)$$

Therefore,

$$(\delta - A)^2 = (A + \delta)^2 \quad (5.19)$$

$$\Rightarrow \delta = 0 \quad (5.20)$$

5.9 Repeat the above exercise when

$$p_x(0) = p$$

Solution:

$$P_e = P_{e|0}p + P_{e|1}(1-p) \quad (5.21)$$

$$= pQ_N(A - \delta) + (1-p)Q_N(A + \delta) \quad (5.22)$$

Differentiating as before, we get:

$$0 = p \frac{1}{\sqrt{2\pi}} e^{-\frac{(\delta-A)^2}{2}} - (1-p) \frac{1}{\sqrt{2\pi}} e^{-\frac{(A+\delta)^2}{2}} \quad (5.23)$$

Taking ln on both sides we have:

$$\ln p - \frac{(\delta-A)^2}{2} = \ln(1-p) - \frac{(A+\delta)^2}{2} \quad (5.24)$$

$$\implies 2\delta A = \ln \frac{1-p}{p} \quad (5.25)$$

$$\implies \delta = \frac{1}{2A} \ln \frac{1-p}{p} \quad (5.26)$$

5.10 Repeat the above exercise using the MAP criterion.

Solution:

Using theorem of conditional probabilities,

$$p_{X|Y}(x|y) = \frac{p_{Y|X}(y|x) \times p_X(x)}{p_Y(y)} \quad (5.27)$$

$$p_{X|Y}(x|y) = \frac{p_X(x) Pr(Ax + N = y)}{p_Y(y)} \quad (5.28)$$

$$p_{X|Y}(x|y) = \frac{p_X(x) Pr(N = y - Ax)}{p_Y(y)} \quad (5.29)$$

$$p_{X|Y}(x|y) = \frac{p_X(x) p_N(y - Ax)}{p_Y(y)} \quad (5.30)$$

$$p_{X|Y}(x|y) = \frac{\frac{1}{\sqrt{2\pi}} p_X(x) e^{-\frac{(y-Ax)^2}{2}}}{p_Y(y)} \quad (5.31)$$

$$(5.32)$$

When $X = 1$, we have:

$$p_{X|Y}(1|y) = \frac{p_{Y|X}(y|1) \times p_X(1)}{p_Y(y)} \quad (5.33)$$

$$= \frac{(1-p) \frac{e^{-\frac{(y-A)^2}{2}}}{\sqrt{2\pi}}}{p \frac{e^{-\frac{(y+A)^2}{2}}}{\sqrt{2\pi}} + (1-p) \frac{e^{-\frac{(y-A)^2}{2}}}{\sqrt{2\pi}}} \quad (5.34)$$

$$= \frac{(1-p) e^{2yA}}{p + (1-p) e^{2yA}} \quad (5.35)$$

Similarly when $X = -1$,

$$p_{X|Y}(-1|y) = \frac{p}{p + (1-p) e^{2yA}} \quad (5.36)$$

Therefore, when $p_{X|Y}(1|y) > p_{X|Y}(-1|y)$, we

have:

$$\frac{(1-p) e^{2yA}}{p + (1-p) e^{2yA}} > \frac{p}{p + (1-p) e^{2yA}} \quad (5.37)$$

$$e^{2yA} > \frac{p}{(1-p)} \quad (5.38)$$

$$y > \frac{1}{2A} \ln \frac{p}{(1-p)} \quad (5.39)$$

Therefore, when Eq. (5.39), we can assert that $X = 1$, and $X = -1$ otherwise.

6 GAUSSIAN TO OTHER

6.1 gaytam singh