

Follow-up response components for all 31 matched compounds

Ordered by follow-up marker score. Component values are diagnostic and do not change the fixed endpoint.

SNS-032	100%	0%	0%	100%	0%	0%	0%
Flavopiridol	100%	0%	0%	75%	50%	0%	0%
Halofuginone	100%	0%	0%	100%	0%	0%	0%
Bazedoxifene HCl	25%	0%	25%	0%	100%	25%	0%
FCCP	0%	100%	0%	100%	25%	0%	0%
Tyrphostin 9	0%	100%	0%	100%	25%	0%	0%
Bazedoxifene Acetate	0%	0%	25%	25%	75%	0%	0%
Alectinib hydrochloride	0%	75%	25%	75%	50%	50%	0%
WAY-600	0%	75%	0%	50%	50%	75%	0%
Rotenone	0%	0%	0%	50%	0%	0%	0%
WAY-657095	0%	50%	25%	50%	100%	100%	0%
RG2833	0%	75%	100%	100%	75%	100%	0%
Bimiralisib	0%	100%	75%	75%	25%	25%	0%
STF-62247	0%	100%	50%	50%	75%	75%	0%
GSK1324726A	0%	100%	0%	100%	25%	0%	0%
Thiothixene	0%	75%	25%	50%	100%	100%	0%
VPS34 inhibitor 1	0%	50%	25%	100%	75%	100%	0%
Tinostamustine	0%	0%	50%	25%	75%	100%	0%
WAY-658979	0%	0%	0%	25%	75%	100%	0%
PIK-III	0%	100%	25%	75%	100%	100%	0%
Duloxetine	0%	75%	25%	75%	75%	100%	0%
Chloroquine Phosphate	0%	100%	75%	75%	75%	25%	0%
Lonafarnib	0%	100%	75%	75%	100%	100%	0%
Trifluoperazine	0%	50%	0%	50%	50%	100%	0%
UNC2025	0%	100%	50%	100%	0%	0%	0%
SKL2001	0%	50%	0%	50%	75%	100%	0%
Vorinostat	0%	100%	50%	0%	50%	50%	0%
Ensartinib dihydrochloride	0%	100%	25%	100%	0%	0%	0%
Sertraline HCl	0%	75%	0%	100%	75%	75%	0%
Simvastatin	0%	100%	25%	50%	75%	100%	0%
Entinostat	0%	100%	100%	25%	75%	75%	0%
	α-synuclein	MAP2 neurites	TH neurites	TH intensity	Living fraction	Living count	Joint

Marker score ≥ 0.5 ; normalized neuronal features and living fraction ≥ -0.5 control SD; living count $\geq 80\%$ of vehicle.
A compound needs $\geq 75\%$ jointly passing wells. These are operational cutoffs, not validated toxicity or safety thresholds.
The corrected controls also fail the joint rule, so a red cell cannot by itself be interpreted as biological harm.