

**Table 1:** Key units in Physical Chemistry

$N_{\text{Av}}$ :	$6.02214 \times 10^{23}$	$\text{mol}^{-1}$		
1 amu:	$1.6605 \times 10^{-27}$	kg		
$k_{\text{B}}$ :	$1.38065 \times 10^{-23}$	$\text{J K}^{-1}$	$8.61734 \times 10^{-5}$	$\text{eV K}^{-1}$
$R$ :	8.314472	$\text{J K}^{-1} \text{mol}^{-1}$	$8.2057 \times 10^{-2}$	$\text{l atm mol}^{-1} \text{K}^{-1}$
$\sigma_{\text{SB}}$ :	$5.6704 \times 10^{-8}$	$\text{J s}^{-1} \text{m}^{-2} \text{K}^{-4}$		
$c$ :	$2.99792458 \times 10^8$	$\text{m s}^{-1}$		
$h$ :	$6.62607 \times 10^{-34}$	$\text{J s}$	$4.13566 \times 10^{-15}$	$\text{eV s}$
$\hbar$ :	$1.05457 \times 10^{-34}$	$\text{J s}$	$6.58212 \times 10^{-16}$	$\text{eV s}$
$hc$ :	1239.8	$\text{eV nm}$		
$e$ :	$1.60218 \times 10^{-19}$	C		
$m_e$ :	$9.10938215 \times 10^{-31}$	kg	1: 0.5109989	$\text{MeV c}^{-2}$
$\epsilon_0$ :	$8.85419 \times 10^{-12}$	$\text{C}^2 \text{J}^{-1} \text{m}^{-1}$	$5.52635 \times 10^{-3}$	$e^2 \text{\AA}^{-1} \text{eV}^{-1}$
$e^2/4\pi\epsilon_0$ :	$2.30708 \times 10^{-28}$	$\text{J m}$	14.39964	$\text{eV \AA}$
$a_0$ :	$0.529177 \times 10^{-10}$	m	0.529177	$\text{\AA}$
$E_{\text{H}}$ :	1	Ha	27.212	eV

## 1 The Classical Foundations

### 1.1 Lecture 0: Introduction

1. Burning lighter
2. Foundations of Physical Chemistry
  - (a) Quantum mechanics
  - (b) Statistical mechanics
  - (c) Thermodynamics, kinetics, spectroscopy
  - (d) Physical and chemical properties of matter

### 1.2 Lecture 1: Basic statistics

1. Discrete probability distributions—Coin flip
  - (a) Example of Bernoulli trial,  $2^n$  possible outcomes from  $n$  flips
  - (b) Number of ways to get  $i$  heads in  $n$  flips,  ${}_nC_i = n!/i!(n-i)!$
  - (c) Probability of  $i$  heads  $P_i \propto {}_nC_i$
  - (d) Normalized probability,  $\tilde{P}_i = P_i / \sum_i P_i = {}_nC_i / 2^n$
  - (e) Expectation value  $\langle i \rangle = \sum_i i \tilde{P}_i$
2. Continuous distributions—temperature
  - (a) Probability density  $P(x)$  has units  $1/x$
  - (b) Normalized  $\tilde{P}(x) = P(x) / \int P(x) dx$
  - (c) (Unitless) probability  $a < x < b = \int_a^b \tilde{P}(x) dx$

(d) Expectation value  $\langle f(x) \rangle = \int f(x) \tilde{P}(x) dx$

(e) Mean =  $\langle x \rangle$

(f) Mean squared =  $\langle x^2 \rangle$

(g) Variance  $\sigma^2 = \langle x^2 \rangle - \langle x \rangle^2$

(h) Standard deviation  $\Delta x = \sigma$

### 3. Boltzmann distribution

(a)  $P(E) \propto e^{-E/k_B T}$ , in some sense the definition of temperature

(b) Energy and its units

(c) Absolute temperature and its units

(d)  $k_B T$  as an energy scale, 0.026 eV at 298 K

(e) Gravity example

i.  $E(h) = mgh$ , linear, continuous energy spectrum

ii. molecule vs car in a gravitational field (Table 2)

iii. Barometric law for gases,  $P = P_0 e^{-mgh/k_B T}$

(f) Kinetic energy in 1-D example

i.  $KE = \frac{1}{2} m v_x^2$

ii.  $P_{1D}(v_x) = \left( \frac{m}{2\pi k_B T} \right)^{1/2} \exp \left( -\frac{m|v_x|^2}{2k_B T} \right)$

iii. Gaussian distribution, mean  $\mu$ , variance  $\sigma^2$

$$G(x) = \frac{1}{\sigma\sqrt{2\pi}} \exp \left( -\frac{(x - \mu)^2}{2\sigma^2} \right)$$

iv. By inspection,  $\mu = \langle v_x \rangle = 0$ ,  $\sigma^2 = \langle v_x^2 \rangle = k_B T/m$

v. Molecule vs car again

(g) Equipartition – energy freely exchanged between all degrees of freedom

**Table 2:** Car vs gas molecule at the earth's surface

	car	gas molecule
$m$	1000 kg	$1 \times 10^{-26}$ kg
$h$	1 m	1 m
$mgh$	9800 J	$9.8 \times 10^{-26}$ J
	$6.1 \times 10^{22}$ eV	$6.1 \times 10^{-7}$ eV
$T$	298 K	298 K
$k_B T$	0.026 eV	0.026 eV
$mgh/k_B T$	$2.4 \times 10^{24}$	$2.3 \times 10^{-5}$
$P(1 \text{ m})/P(0)$	$e^{-2.4 \times 10^{-24}}$	0.99998
$\langle h \rangle$	0 m	42 km
$\langle v_x \rangle^{1/2}$	$2 \times 10^{-12}$ m/s	640 m/s

**Table 3:** Energy conversions and correspondences

	J	eV	Hartree	$\text{kJ mol}^{-1}$	$\text{cm}^{-1}$
1 J =	1	$6.2415 \times 10^{18}$	$2.2937 \times 10^{17}$	$6.0221 \times 10^{20}$	$5.0340 \times 10^{22}$
1 eV =	$1.6022 \times 10^{-19}$	1	0.036748	96.485	8065.5
1 Ha =	$4.3598 \times 10^{-18}$	27.212	1	2625.6	219474.6
1 $\text{kJ mol}^{-1}$ =	$1.6605 \times 10^{-21}$	0.010364	$3.8087 \times 10^{-4}$	1	83.5935
1 $\text{cm}^{-1}$ =	$1.986410^{-23}$	$1.23984 \times 10^{-4}$	$4.55623 \times 10^{-6}$	0.011963	1

---

```

1  import numpy as np
2  import matplotlib.pyplot as plt
3
4  RO = 8.31441    J/mol K
5  mass = 28. /1000  kg/mol N2
6
7  def Boltzmann(E,T):
8      return np.exp(E/(RO*T))/(RO*T)
9
10 def MB1D(v,T):
11     return np.sqrt(mass/(2*np.pi*RO*T))*np.exp((mass*v*v)/(2*RO*T))
12
13 def MB(c,T):
14     K = 0.5 * mass * c * c
15     degeneracy = 4 * np.pi * c * c
16     normalization = (mass/(2*np.pi*RO*T))**1.5
17     return normalization*Boltzmann(K,T)
18
19 energy = np.linspace(0,3000,1500)
20 velocity = np.linspace(1000,1000,1000)
21 speed = np.linspace(0,1500,1000)
22
23 plt.figure()
24 for Temperature in [100,300,1000]:
25     Probability = Boltzmann(energy,Temperature)
26     plt.plot(Probability,energy,label=0 K.format(Temperature))
27
28 legend = plt.legend()
29
30 plt.ylabel(Energy (J/mol))
31 plt.xlabel(Probability (mol/J))
32 plt.title(Boltzmann distribution at various temperatures)
33 plt.savefig(./Images/Boltzmann.png)
34
35 plt.figure()
36 for Temperature in [100,200,300,400,500]:
37     Probability = MB1D(velocity,Temperature)
38     plt.plot(velocity,Probability,label= K.format(Temperature))
39
40 legend=plt.legend()
41 plt.xlabel(Velocity (m/s))
42 plt.ylabel(Probability (1/(m/s)))
43 plt.title(Boltzmann distribution at various temperatures)
44 plt.savefig(./Images/MB1D.png)
45
46 plt.figure()
47 for Temperature in [100,200,300,400,500]:
48     Probability = MB(speed,Temperature)
49     plt.plot(speed,Probability,label= K.format(Temperature))
50
51 legend=plt.legend()
52 plt.xlabel(Speed (m/s))
53 plt.ylabel(Probability (1/(m/s)))


```

---


```

54 plt.title(Boltzmann distribution at various temperatures)
55 plt.savefig('./Images/MB.png')

```



**Figure 1:** Boltzmann distribution at various temperatures



**Figure 2:** One-dimensional (Gaussian) velocities of N<sub>2</sub> gas


### 1.3 Lecture 2: Kinetic theory of gases

#### 1. Postulates

- (a) Gas is composed of molecules in constant random, thermal motion
- (b) Molecules only interact by perfectly elastic collisions
- (c) Volume of molecules is  $\ll$  total volume

#### 2. Maxwell-Boltzmann distribution of molecular speeds

- (a) Speed  $v = \sqrt{v_x^2 + v_y^2 + v_z^2}$



**Figure 3:** Maxwell-Boltzmann speed distribution of  $N_2$  gas

(b)  $P_{MB}(v)dv = P_{1D}(v_x)P_{1D}(v_y)P_{1D}(v_z) * \text{degeneracy}(v)dv$

(c) mean speeds  $\propto \sqrt{T}$

(d) mean energy  $U = \frac{3}{2}RT$  and heat capacity  $C_v = \frac{3}{2}R$

### 3. Flux and pressure

(a) Velocity flux  $j(v_x)dv_x = v_x \frac{N}{V} P(v_x)dv_x$ , molecules /area /time / $v_x$

(b) Wall collisions,  $J_w$ , total collisions /area /time


(c) Momentum exchange, pressure, ideal gas law

### 4. Collisions and mean free path

(a) Collision cross section  $\sigma = \pi d^2$ , size of molecule

(b) Molecular collisions,  $z$  per molecule and  $z_{AA}$  per volume

(c) Mean free path,  $\lambda$ , mean distance between collisions



**Figure 4:** Diffusional spreading,  $\sqrt{\langle x^2 \rangle} = \sqrt{2Dt}$

**Table 4:** Kinetic theory of gases key equations

Boltzmann distribution ( $g(E)$ : degeneracy of $E$ )	$P(E) = g(E)e^{-E/k_B T}$
Maxwell-Boltzmann distribution	$P_{\text{MB}}(v) = 4\pi v^2 \left(\frac{m}{2\pi k_B T}\right)^{3/2} \exp\left(-\frac{mv^2}{2k_B T}\right)$
Mean and RMS speeds	$\langle v \rangle = \left(\frac{8k_B T}{\pi m}\right)^{1/2} \quad \langle v^2 \rangle^{1/2} = \left(\frac{3k_B T}{m}\right)^{1/2}$
Pressure	$\langle P \rangle = \frac{\Delta p}{\Delta t} = m \frac{N}{V} \frac{1}{3} \langle v^2 \rangle = \frac{N k_B T}{V} = \frac{n R T}{V}$
Wall collision frequency	$J_W = \frac{1}{4} \frac{N}{V} \langle v \rangle = \frac{P}{(2\pi m k_B T)^{1/2}}$
Molecular collision frequency	$z = \sqrt{2} \sigma \langle v \rangle \frac{N}{V} = \frac{4\sigma P}{(\pi m k_B T)^{1/2}}$
Total collisions	$z_{AA} = \frac{1}{2} \frac{N}{V} z$
Mean free path	$\lambda = \frac{\langle v \rangle}{z} = \frac{V}{\sqrt{2} \sigma N}$
Graham's effusion law	$\frac{dN}{dt} = \text{Area} \cdot J_w \propto 1/m^{1/2}$
Effusion from a vessel	$P = P_0 e^{-t/\tau}, \tau = \frac{V}{A} \left(\frac{2\pi m}{k_B T}\right)^{1/2}$
Self-diffusion constant	$D_{11} = \frac{1}{3} \langle v \rangle \lambda$
Diffusion rate	$\langle x^2 \rangle^{1/2} = \sqrt{2Dt} \quad \langle r^2 \rangle^{1/2} = \sqrt{6Dt}$
Einstein-Smoluchowski equation	$D_{11} = \frac{\delta^2}{2\tau}$
Stokes-Einstein equation for liquids	$D_{11} = \frac{k_B T}{4\pi\eta r}$ "Slip" boundary
	$D_{\text{Brownian}} = \frac{k_B T}{6\pi\eta r}$ "Stick" boundary

## 1.4 Lecture 3: Transport

1. Effusion and Graham's law, effusion rate  $\propto MW^{-1/2}$
2. Fick's first law: net flux proportional to concentration gradient
  - (a)  $j_x = -D \frac{dc}{dx}$
  - (b) Self-diffusion constant,  $D = \frac{1}{3} \lambda \langle v \rangle$
3. Knudsen diffusion,  $D = \frac{1}{3} l \langle v \rangle$
4. Fick's second law: time evolution of concentration gradient
  - (a) Continuity with no advection:  $\frac{\partial c}{\partial t} = -\nabla \cdot \vec{j} + \text{gen}$
  - (b) One-dimension:  $\frac{dc}{dt} = D \frac{d^2 c}{dx^2}$
  - (c) Diffusion has Gaussian probability distribution:  $c(x, t)/c_0 = [2\sqrt{\pi Dt}]^{-1} \exp(-x^2/4Dt)$
5. Seeing is believing—Brownian motion
  - (a) Seemingly random motion of large particles (“dust”) due to “kicks” from invisible molecules
  - (b) Einstein receives Nobel Prize for showing:
    - i. Motion follows same Gaussian diffusion behavior
    - ii. From steady-state arguments in a field, diffusion constant is ratio of Boltzmann energy,  $k_B T$ , to mobility
    - iii. Mobility inversely related to viscosity
  - (c) Stokes-Einstein equation
  - (d) Allows measurement of Avogadro's number, final proof of kinetic theory
  - (e) Similar model for diffusion of liquid molecules, slip boundary
6. Random walk model of diffusion
  - (a) Binomial distribution
  - (b) Large  $N$  and Stirling approximation
  - (c) Einstein-Smoluchowski relation

## 2 Quantum Mechanics: Blurred Lines Between Particles and Waves

### 2.1 Lecture 4: Duality and demise of classical physics

#### 2.1.1 Properties of waves

1. traveling waves,  $\psi(x, t) = A \sin(kx - \omega t)$ ,  $k = 2\pi/\lambda$ ,  $\omega = 2\pi\nu$
2. standing waves,  $\psi(x, t) = A \sin(kx) \cos(\omega t)$
3. interference, diffraction
4. energy proportional to amplitude squared
5. Expected energy of a classical oscillator,  $\langle \epsilon \rangle_\nu = k_B T$  for all  $\nu$

### 2.1.2 Blackbody radiation

1. Hohlraum spectrum (like the sun) empirically observed to obey:
  - (a) Stefan-Boltzmann law, total irradiance
  - (b) Wien's displacement law
2. Rayleigh-Jeans predicts spectrum using classical physics
  - (a) standing waves + classical oscillators  $\rightarrow$  ultraviolet catastrophe
3. Planck model
  - (a) Energy spectrum of oscillators are *quantized*,  $\epsilon_\nu = nh\nu$
  - (b) Expected energy of a quantized oscillator,  $\langle \epsilon \rangle_\nu = h\nu / (e^{h\nu/k_B T} - 1)$
  - (c) Correctly reproduces Stefan-Boltzmann and Wien Laws!

### 2.1.3 Heat capacities of solids

1. Law of DuLong and Pettite,  $C_v = 3R$ , fails at low  $T$
2. Einstein model
  - (a) Atomic vibrations are *quantized*,  $\epsilon_n = nh\nu$
  - (b) Heat capacity goes to zero at low  $T$

### 2.1.4 Photoelectric effect

1. Stopping potential and work function,  $E_{\text{kinetic}} = h\nu - W$
2. Kinetic energy varies with light frequency, number of electrons varies with light intensity

### 2.1.5 Compton effect

1. light scattering of electrons changes  $\lambda$
2. Photon properties,  $\epsilon = h\nu, p = h/\lambda$

### 2.1.6 Wave-particle duality

### 2.1.7 Rutherford, planetary model of atom

1. Inconsistent with Maxwell's equations

### 2.1.8 Bohr model of H atom

1. Discrete H energy spectrum and Rydberg formula
2. Bohr model (the old quantum mechanics)
  - (a) Stable electron "orbits," quantized angular momentum
  - (b) Light emission corresponds to orbital jumps,  $\nu = \Delta E/h$



- (c) Bohr equations
- (d) Comparison with Rydberg formula
- (e) Failure for larger atoms

### 2.1.9 de Broglie relation

1.  $\lambda = h/p$  universally
2. Relation to Bohr orbits
3. Davison and Germer experiment,  $e^-$  diffraction off Ni

---

```

1  import numpy as np
2  import matplotlib.pyplot as plt
3
4  hc = 1239.8          eV nm
5  c = 2.9979e8 * 1.e9  nm/s
6  k = 8.61734e5       eV /K
7  hck = hc/k          nm K
8
9  def Irrad(wl,T):
10     return (8. * np.pi * hc * c * wl**5) / (np.exp(hck/(wl*T))1)
11  def PlanckEnergy(wl,T):
12     return (hc/wl) / (np.exp(hck/(wl*T))1)
13
14  plt.figure()
15  wl=np.linspace(100,5000,1000)
16  for T in [1000.,2000.,3000.,4000.,5000.]:
17      Intensity = Irrad(wl,T)
18      plt.plot(wl,Intensity,label= K.format(T))
19
20  legend=plt.legend()
21  plt.xlabel(Wavelength (nm))
22  plt.ylabel(Irradiance (eV/nm3/s))
23  plt.title(Boltzmann distribution at various temperatures)
24  plt.savefig(./Images/BlackBody.png)
25
26  plt.figure()
27  color=[red,orange,green,blue,violet]
28  wl=np.linspace(100,20000,1000)
29  for T in [1000.,2000.,3000.,4000.,5000.]:
30      Energy = PlanckEnergy(wl,T)
31      plt.plot(wl,Energy,label= K.format(T),color=color[0])
32      kT = k*T
33      plt.plot([100,max(wl)], [kT,kT],ls=,color=color.pop(0))
34
35  legend=plt.legend()
36  plt.xlabel(Wavelength (nm))
37  plt.ylabel(Energy (eV))
38  plt.title(Boltzmann distribution at various temperatures)
39  plt.savefig(./Images/Planck.png)

```



---

## 2.2 Lecture 5: Postulates of quantum mechanics

### 2.2.1 Schrödinger equation describes wave-like properties of matter

#### 2.2.2 Born interpretation

1. wavefunction is a probability amplitude
2. wavefunction squared is probability density

**Figure 5:** Blackbody irradiance**Figure 6:** Average energy of a Planck quantized oscillator

**Table 5:** The new physics

Stefan-Boltzmann Law	$\int I(\lambda, T) d\lambda = \sigma_{\text{SB}} T^4$
Wien's Law	$\lambda_{\text{max}} T = 2897768 \text{ nm K}$
Rayleigh-Jeans eq	$I(\lambda, T) = \frac{8\pi}{\lambda^4} k_B T c$
Blackbody irradiance	$I(\lambda, T) = \frac{8\pi}{\lambda^5} \frac{hc^2}{e^{hc/\lambda k_B T} - 1}$
Einstein crystal	$C_v = 3R \left( \frac{h\nu}{k_B T} \right)^2 \frac{e^{h\nu/k_B T}}{(e^{h\nu/k_B T} - 1)^2}$
Photon energy	$\epsilon = h\nu = hc/\lambda$
Rydberg equation	$\nu = R_H c \left( 1/n^2 - 1/k^2 \right)$
Bohr equations	$l_n = n\hbar$
$n = 1, 2, \dots$	$r_n = n^2 \left( \frac{4\pi\epsilon_0 \hbar^2}{e^2 m_e} \right) = n^2 a_0$
	$E_n = -\frac{m_e e^4}{8\epsilon_0^2 h^2} \frac{1}{n^2} = -\frac{E_H}{2} \frac{1}{n^2}$
	$p_n = \frac{e^2}{4\pi\epsilon_0} \frac{m_e}{\hbar} \frac{1}{n} = p_0 \frac{1}{n}$
de Broglie equation	$\lambda = \frac{h}{p}$

### 2.2.3 Postulates

1. Wavefunction contains all information about a system
2. Operators used to extract that information
  - (a) QM operators are *Hermitian*
  - (b) Have eigenvectors and real eigenvalues,  $\hat{O}\psi_i = o\psi_i$
  - (c) Are orthogonal,  $\langle\psi_i|\psi_j\rangle = \delta_{ij}$
  - (d) Always observe an eigenvalue when making an observation
3. Expectation values
4. Energy-invariant wavefunctions given by Schrödinger equation
5. Uncertainty principle

### 2.2.4 Particle in a box illustrations

## 2.3 Lecture 6: Particle in a box model

### 2.3.1 Particle between infinite walls, electron confined in a wire

### 2.3.2 Classical solution, either stationary or uniform bouncing back and forth

### 2.3.3 One-dimensional QM solutions

1. Schrödinger equation and boundary conditions
2. discrete, quantized solutions
3. standing waves,  $\lambda = 2L/n$ ,  $n - 1$  nodes, non-uniform probability
4. Ho paper, STM of Pd wire
5. zero point energy and uncertainty
6. correspondence principle
7. superpositions

### 2.3.4 Finite walls and tunneling

1. Potential well of finite depth  $V_0$
2. Finite number of bound states
3. Classical region,  $\psi(x) e^{ikx} + e^{-ikx}$ ,  $k = \sqrt{2mE}/\hbar$
4. “Forbidden” region,  $\psi(x) e^{\kappa x} + e^{-\kappa x}$ ,  $\kappa = \sqrt{2m(V_0 - E)}/\hbar$
5. Non-zero probability to “tunnel” into forbidden region
6. Tunneling between two adjacent wells: chemical bonding, STM, nanoelectronics
7. H atom tunneling:  $\text{NH}_3$  inversion, H transfer, kinetic isotope effect

**Table 6:** Postulates of Non-relativistic Quantum Mechanics

**Postulate 1: The physical state of a system is completely described by its wavefunction  $\Psi$ .** In general,  $\Psi$  is a complex function of the spatial coordinates and time.  $\Psi$  is required to be:

- I. Single-valued
- II. continuous and twice differentiable
- III. square-integrable ( $\int \Psi^* \Psi d\tau$  is defined over all finite domains)
- IV. For bound systems,  $\Psi$  can always be normalized such that  $\int \Psi^* \Psi d\tau = 1$

**Postulate 2:** To every physical observable quantity  $M$  there corresponds a Hermitian operator  $\hat{M}$ . **The only observable values of  $M$  are the eigenvalues of  $\hat{M}$ .**

Physical quantity	Operator	Expression
Position $x, y, z$	$\hat{x}, \hat{y}, \hat{z}$	$x, y, z$
Linear momentum $p_x, \dots$	$\hat{p}_x, \dots$	$-i\hbar \frac{\partial}{\partial x}, \dots$
Angular momentum $l_x, \dots$	$\hat{p}_x, \dots$	$-i\hbar \left( y \frac{\partial}{\partial z} - z \frac{\partial}{\partial y} \right), \dots$
Kinetic energy $T$	$\hat{T}$	$-\frac{\hbar^2}{2m} \nabla^2$
Potential energy $V$	$\hat{V}$	$V(\mathbf{r}, t)$
Total energy $E$	$\hat{H}$	$-\frac{\hbar^2}{2m} \nabla^2 + V(\mathbf{r}, t)$

**Postulate 3:** If a particular observable  $M$  is measured many times on many identical systems in a state  $\Psi$ , the average results will be the expectation value of the operator  $\hat{M}$ :

$$\langle M \rangle = \int \Psi^* (\hat{M} \Psi) d\tau$$


**Postulate 4:** The energy-invariant states of a system are solutions of the equation

$$\begin{aligned} \hat{H} \Psi(\mathbf{r}, t) &= i\hbar \frac{\partial}{\partial t} \Psi(\mathbf{r}, t) \\ \hat{H} &= \hat{T} + \hat{V} \end{aligned}$$

The time-independent, stationary states of the system are solutions to the equation

$$\hat{H} \Psi(\mathbf{r}) = E \Psi(\mathbf{r})$$

**Postulate 5:** (The **uncertainty principle**.) Operators that do not commute ( $\hat{A}(\hat{B}\Psi) \neq \hat{B}(\hat{A}\Psi)$ ) are called *conjugate*. Conjugate observables cannot be determined simultaneously to arbitrary accuracy. For example, the standard deviation in the measured positions and momenta of particles all described by the same  $\Psi$  must satisfy  $\Delta x \Delta p_x \geq \hbar/2$ .

**Table 7:** Particle-in-a-box model

### 2.3.5 Multiple dimensions

1. separation of variables, one quantum number for each dimension

### 2.3.6 Introduce Pauli principle for fermions?

## 2.4 Lecture 7: Harmonic oscillator

### 2.4.1 Classical harmonic oscillator

1. Hooke's law,  $F = -k(x - x_0)$ ,  $k$  spring constant
2. Continuous sinusoidal motion
3.  $x(t) = A \sin(\frac{k}{\mu})^{1/2} t$ ,  $\nu = \frac{1}{2\pi}(\frac{k}{\mu})^{1/2}$ ,  $E = \frac{1}{2}kA^2$
4. Exchanging kinetic and potential energies

### 2.4.2 Quantum harmonic oscillator


1. Solutions like P-I-A-B, waves, nodes
2. Zero-point energy

3. Expectation values  $\langle x^2 \rangle = \alpha^2(v + 1/2)$ ,  $\langle V(x) \rangle = \frac{1}{2}h\nu(v + \frac{1}{2})$
4. Classical turning point and tunneling
5. Classical limiting behavior

### 2.4.3 HCl example

1. Reduced mass,  $\frac{1}{\mu} = \frac{1}{m_A} + \frac{1}{m_B}$
2. Anharmonicity, Morse potential

**Table 8:** Harmonic oscillator model



## 2.5 Lecture 8: Vibrational spectroscopy

### 2.5.1 Spectroscopy is quantitative measurement of interaction of light with matter

1. Observed  $I(\nu)/I(\nu_0)$

2. Bohr condition,  $|E_f - E_i|/h = \nu = c\tilde{\nu} = c/\lambda$
3. Intensities determined by state populations and transition probabilities

### 2.5.2 Einstein coefficients

1. Stimulated absorption,  $dn_1/dt = -n_1 B\rho(\nu)$
2. Stimulated emission,  $dn_2/dt = -n_2 B\rho(\nu)$
3. Spontaneous emission,  $dn_2/dt = -n_2 A$ ,  $A = \left(\frac{8\pi h\nu^3}{c^3}\right) B$
4.  $1/A = \text{lifetime}$

### 2.5.3 Transition probability

1. Einstein coefficient  $B_{if} = \frac{|\mu_{if}|^2}{6\epsilon_0 h^2}$
2. Classical electric dipole,  $\vec{\mu} = q \cdot \vec{r}$ , quantum dipole operator  $\hat{\mu} = e \cdot \vec{r}$
3. Transition dipole moment,  $\mu_{if} = \left(\frac{d\mu}{dx}\right) \langle \psi_i | \hat{\mu} | \psi_f \rangle$
4. Selection rules—conditions that make  $\mu_{if}$  non-zero, “allowed” vs “forbidden” transitions

### 2.5.4 Vibrational transitions

1. Gross selection rule: dynamic dipole  $d\mu/dx$  non-zero
2. Specific selection rule: dipole integral  $\langle \psi_v | \hat{\mu} | \psi_{v'} \rangle = 0$  unless  $\Delta v = \pm 1$
3. Allowed  $\Delta E = h\nu$
4. Boltzmann,  $v = 1$  states dominate at normal  $T$

### 2.5.5 Vibrational spectroscopy

1. Diatomics, homo- vs. heteronuclear
2. Polyatomics,  $3n - 6$  ( $3n - 5$  for linear polyatomic) vibrational modes
3. CO<sub>2</sub> example


### 2.5.6 Raman spectroscopy

## 2.6 Lecture 9: Rigid Rotor

### 2.6.1 Classical rigid rotor

1. Rotation about an axis vs linear motion
2. Moment of inertia  $I = \mu r^2$
3. Angular momentum,  $\mathbf{l} = I\omega = \mathbf{r} \times \mathbf{p}$ ,  $T = \frac{1}{2} I \omega^2$
4. Angular momentum and energy continuous variable



**Table 9:** 2-D rigid rotor model

### 2.6.2 Quantum rotor in a plane

1. Angular momentum and kinetic energy operators in polar coordinates,  $\hat{l}_z = -i\hbar \frac{d}{d\phi}$
2. Eigenfunctions and energy spectrum
3. No zero point energy
4. Angular momentum eigenfunctions,  $l_z = m_l \hbar$
5. Energy superpositions and localization

### 2.6.3 Quantum rotor in 3-D

1. Angular momentum and kinetic energy operators in spherical coordinates

2. Spherical harmonic solutions,  $Y_{lm_l}$
3. Azimuthal QN  $l=0, 1, \dots$
4. Magnetic QN  $m_l = -l, -l+1, \dots, l$
5. Energy spectrum,  $2l+1$  degeneracy
6. Vector model - can only know total  $|L|$  and  $L_z$
7. Wavefunctions look like atomic orbitals,  $l$  nodes

#### 2.6.4 Particle angular momentum

1. Fermions, mass, half-integer spin
  - (a) Electron,  $s = 1/2, m_s = \pm 1/2$
2. Bosons, force-carrying, integer spin

#### 2.6.5 Diatomic rotational spectroscopy

1. Rotational constant  $B = \hbar/4\pi I c \text{ cm}^{-1}$ ,  $I = \mu R^2$
2. Gross selection rule: dipole moment non-zero
3. Specific selection rule:  $\Delta l = \pm 1$ ,  $\Delta m_l = 0, \pm 1$
4.  $\Delta \tilde{E}_l = 2B(l+1) \text{ cm}^{-1}$
5. Rotational state populations

#### 2.6.6 Polyatomic rotational spectroscopy

#### 2.6.7 Vibration-rotation spectroscopy

1. Harmonic oscillator + rigid rotor
2. Selection rules:  $\Delta v = \pm 1, \Delta l = \pm 1$
3.  $R$  branch:  $\Delta \tilde{E} = \tilde{\nu} + 2B(l+1), \Delta l = 1$
4.  $P$  branch:  $\Delta \tilde{E} = \tilde{\nu} - 2B(l), \Delta l = -1$

### 3 Statistical Mechanics: The Bridge from the Tiny to the Many

**Table 10:** 3-D rigid rotor model

$$V(\theta, \phi) = 0, 0 \leq \phi \leq 2\pi, 0 \leq \theta < \pi$$

$$\hat{L}^2 = -\hbar^2 \left[ \frac{1}{\sin^2 \theta} \frac{\partial^2}{\partial \phi^2} + \frac{1}{\sin \theta} \frac{\partial}{\partial \theta} \left( \sin \theta \frac{\partial}{\partial \theta} \right) \right]$$

$$\hat{H}_{\text{rot}} = \frac{1}{2I} \hat{L}^2$$

$$Y_{lm_l}(\theta, \phi) = N_l^{|m_l|} P_l^{|m_l|}(\cos(\theta)) e^{im_l \phi}$$

$$l = 0, 1, 2, \dots, \quad m_l = 0, \pm 1, \dots, \pm l$$

$$E_l = \frac{\hbar^2}{2I} l(l+1)$$

$$|L| = \hbar \sqrt{l(l+1)}, L_z = m_l \hbar$$

