

JRC TECHNICAL REPORT

Methane as a greenhouse gas

From 'unconventional' methane production and recovery to biological mitigation options: A literature review relying on text-mining tools

Marmier, A; Schosger J.-P.

2020



This publication is a technical report by the Joint Research Centre (JRC), the European Commission's science and knowledge service. It aims to provide evidence-based scientific support to the European policymaking process. The scientific output expressed does not imply a policy position of the European Commission. Neither the European Commission nor any person acting on behalf of the Commission is responsible for the use that might be made of this publication. For information on the methodology and quality underlying the data used in this publication for which the source is neither Eurostat nor other Commission services, users should contact the referenced source. The designations employed and the presentation of material on the maps do not imply the expression of any opinion whatsoever on the part of the European Union concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries.

Contact information

Name: Alain Marmier

Address: European Commission – Joint Research Centre, P.O. Box 2, 1755 ZG, Petten, the Netherlands

Email: alain.marmier@ec.europa.eu

Tel. +31 22456-5147

EU Science Hub

<https://ec.europa.eu/jrc>

JRC121995

EUR 30392

PDF

ISBN 978-92-76-22969-8

ISSN 1831-9424

doi:10.2760/09370

Luxembourg: Publications Office of the European Union, 2020

© European Union, 2020



The reuse policy of the European Commission is implemented by the Commission Decision 2011/833/EU of 12 December 2011 on the reuse of Commission documents (OJ L 330, 14.12.2011, p. 39). Except otherwise noted, the reuse of this document is authorised under the Creative Commons Attribution 4.0 International (CC BY 4.0) licence (<https://creativecommons.org/licenses/by/4.0/>). This means that reuse is allowed provided appropriate credit is given and any changes are indicated. For any use or reproduction of photos or other material that is not owned by the EU, permission must be sought directly from the copyright holders.

All content © European Union, 2020, except cover page, Copyrights: Željko Radojko, Source: Fotolia 59639069; page 7, figure 1, Copyrights: Jagersma, 2006. Source: WUR, <https://www.wur.nl/en/show/Anaerobic-oxidation-of-methane.htm>.

How to cite this report: Marmier, A., Schosger, J.-P., *Methane as greenhouse gas, From 'unconventional' methane production and recovery to biological mitigation options: A literature review relying on text-mining tools*, Publications Office of the European Union, Luxembourg, 2020, ISBN 978-92-76-22969-8, doi:10.2760/09370, JRC121995.

Contents

| | |
|--|----|
| Foreword..... | 1 |
| Acknowledgements | 2 |
| Abstract | 3 |
| 1. Introduction..... | 4 |
| 2. Methane recovery technologies and controlled biomethane production, starting from an economic perspective | 5 |
| 2.1. Current processes and technologies for methane concentration and upgrading..... | 5 |
| 2.2. Technologies behind biomethane production..... | 6 |
| 3. Towards biological remediation strategies | 7 |
| 4. Recommendations and open questions..... | 10 |
| 4.1. Knowledge gathering and sharing | 10 |
| 4.2. Specific emissions and atmospheric concentration mitigation options | 10 |
| 4.3. Ethical aspects | 10 |
| 5. Conclusions | 11 |
| References | 12 |
| List of abbreviations and definitions | 19 |
| List of figures | 20 |
| Annexes | 21 |
| Annex 1: Literature mapping – Clustering of publications | 21 |
| Annex 2: JRC activities on methane | 26 |

Foreword

Methane is making the news. And the news is not always positive:

- *Nature Climate Change*, 10 June 2020: 'Remote sensing northern lake methane ebullition', <https://www.nature.com/articles/s41558-020-0762-8.pdf>, referred to by the NASA Earth Observatory: 'Satellites size up bubbles of methane in lake ice', <https://earthobservatory.nasa.gov/images/146940/satellites-size-up-bubbles-of-methane-in-lake-ice>
- *Nature*, 14 July 2020: 'Global methane levels soar to record high', <https://www.nature.com/articles/d41586-020-02116-8>
- *The Guardian*, 22 July 2020: 'First active leak of sea-bed methane discovered in Antarctica', <https://www.theguardian.com/environment/2020/jul/22/first-active-leak-of-sea-bed-methane-discovered-in-antarctica>
- *Science*, 28 July 2020: 'Siberia's "gateway to the underworld" grows as record heat wave thaws permafrost', <https://www.sciencemag.org/news/2020/07/siberia-s-gateway-underworld-grows-record-heat-wave-thaws-permafrost>
- Phys.org, 29 July 2020: 'New space satellite pinpoints industrial methane emissions', <https://phys.org/news/2020-07-space-satellite-industrial-methane-emissions.html>
- Space.com, 29 July 2020: 'Strange gas signature on Mars may help explain methane mystery', <https://www.space.com/mars-atmosphere-gas-detections-methane-mystery.html>
- Helmholtz Centre for Ocean Research Kiel (GEOMAR), 30 July 2020: 'New study confirms extensive gas leaks in the North Sea', https://www.eurekalert.org/pub_releases/2020-07/hcfo-nsc073020.php

This report builds on JRC's horizon scanning capacities, led within the JRC by the JRC.I2 unit. It dives deeper into the emerging issue of global warming, focusing on methane as a recovered or biogenerated primary energy source and on biological mitigation processes of methane, a potent greenhouse gas. It follows JRC.C7 involvement in negative emissions technologies in the framework of the Future and Emerging Technologies work package of the Low Carbon Energy Observatory, which highlighted the significant efforts to remove carbon dioxide from the atmosphere. Motivations for further work on methane are summarised in the JRC Science behind the Debate fact sheet JRC121991, while an overview of JRC activities related to methane is available in Annex 2.

Acknowledgements

The authors acknowledge the reviews provided by Elisa Boelman and Laura Contor (JRC.C7) and Robert Ziegler (JRC.C3).

Authors

Marmier, A; Schosger J.-P.

Abstract

This report aims to address climate change by focusing on methane (CH₄), a greenhouse gas with high global warming potential. Methane is the second most important greenhouse gas in terms of concentration and impact on the climate. This highlights the importance of focusing on processes that are able to lower the methane concentration in the atmosphere, where it is considered a significant contributor to climate change. This can be achieved by (a combination of) emissions avoidance, recovery, mitigation, capture and combustion or use.

This review, in itself, is also significant in terms of scientific process. The narrow literature review allowed for the (manual) definition of a mapping structure. By using text-mining tools, this structure can now be applied to a broader set of scientific documents. Such scaling up will enrich the mapping and help to enhance the mapping structure. This enhanced knowledge gathering will in turn lead to better mapping of ongoing activities and thus mapping of the gaps within.

The review analysed 109 scientific communications, mapped along various dimensions:

- by type of document: 17 documents reviewed specific aspects of methane;
- by environment (e.g. atmosphere, soil, freshwater and saline water) and by source of methane held in these environments;
- by process: methane production through methanogenesis, methane absorption for separation or for storage, genetic approaches, as well as methanotrophy, methane oxidation and methane conversion;
- by by- and co-products, such as carbonates, chromate, copper, iron, manganese, nitrous compounds (N₂O, NO_x), ammonia, nitrate, sulfur and sulfate, but also fuels and chemicals, such as lactic acid, methanol and acetate;
- by technologies relying on living organisms and micro-organisms, such as biochar, biotrickling filtration, digesters, microbial fuel cells and relevant genetic technologies (e.g. next-generation sequencing), or relying on ((in)organic) materials such as membranes, sorbents or pressure swing absorption;
- by stage of development: demonstration or commercial projects, patents, estimates of the potential for methane emission reduction, quantified costs, other economic aspects, programmes and partnerships.

The review provides a list of possible technologies and constitutes a step towards finding the most cost-effective approaches to methane emission mitigation. It identifies processes and technologies for the biological oxidation of methane, as well as for methane recovery and controlled biomethane production. However, it also highlights how broad the topic is and the numerous questions that remain. As such, this review calls for additional and more targeted investigations. The most salient investigations are as follows:

- the ongoing efforts to thoroughly identify the spatial/geographical distribution of greenhouse gas sources to better understand the contributions of each greenhouse gas, source and sector to global warming;
the role of microorganisms in methane production and mitigation and the ethics of genetically modifying such natural processes;
- the added value of joint anaerobic oxidation of methane and denitrification processes, leading to both nitrous oxide (N₂O) and CH₄ consumption, which are powerful greenhouse gases;
- the feasibility of developing the air capture/consumption of greenhouse gases besides and/or coupled to carbon dioxide (CO₂).

Technology is an enabler. Should the technological bottlenecks above be removed, the economic, environmental and social benefits could be realised.

1. Introduction

Three gases (carbon dioxide or CO₂, methane or CH₄ and nitrous oxide or N₂O) account for about 98 % of the global annual emissions of greenhouse gases (GHGs). CH₄, formed when four hydrogen atoms bond to one atom of carbon, is a molecule that is lighter than air. It is the primary component of natural gas, which generates roughly 22 % of the world's electricity, making it the largest primary energy source for electricity after coal.

In order to compare the warming potential of various GHGs over a certain period of time with that of CO₂, the Intergovernmental Panel on Climate Change (IPCC) has set up a common scale for measuring the climate effects of different gases. These are calculated as the global warming potential (GWP) and correspond to the mass of CO₂ needed to produce a similar warming effect over a given period of time, typically 20 and 100 years

In its Fifth Assessment Report (AR5) ⁽¹⁾, the IPCC estimates the GWP of CH₄ to be 84 over a period of 20 years and 28 over a period of 100 years. Therefore, 1 tonne of emitted CH₄ can be considered to be equivalent to 28 tonnes of emitted CO₂ when looking at its warming impact over 100 years.

The current atmospheric concentrations of CO₂ and CH₄ are about 400 ppm and 2 ppm, respectively; therefore, the concentration of CH₄ is 200 times lower than that of CO₂. Nevertheless, this tiny fraction of CH₄ has a warming potential equivalent to $2 \times 28 = 56$ ppm of CO₂ over 100 years and $2 \times 84 = 168$ ppm of CO₂ over 20 years. Therefore, even though CH₄ is 200 times less concentrated than CO₂, its warming potential is only about seven times lower than that of CO₂ over a period of 100 years and about 40 % of the CO₂ warming potential over a period of 20 years.

Therefore, rising CH₄ concentrations in the atmosphere deserve special attention because of CH₄'s significant potency as a GHG. Efforts to identify the various sources of CH₄ and mitigate its warming effects should be strengthened.

The objective of this study is to map CH₄ recovery and emission mitigation processes and technologies (including their current level of development), as well as CH₄ sources. While this mapping is not exhaustive, it nonetheless remains instrumental for issuing recommendations on emission reductions and atmospheric concentration reductions and in the effort to limit the impact of climate change and global warming.

⁽¹⁾ IPCC, *Climate change 2014: Synthesis report. Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change (IPCC)*, IPCC, Geneva, Switzerland, 2014, available online at: <https://www.ipcc.ch/assessment-report/ar5/>, accessed 4.9.2020.

2. Methane recovery technologies and controlled biomethane production, starting from an economic perspective

Methane is considered to be a primary energy source and an energy carrier: an economic asset subject to its own market and geopolitics. Outside the traditional natural gas industry, which is generally not considered in this review, numerous technologies, including separation and concentration, target the economic production of CH₄. Patents (Air Liquide targets landfill gas-to-energy, 2007), demonstration or commercial projects (Bhattarai et al., 2017; Fei et al., 2020; He et al., 2018; Hur et al., 2017; Liu et al., 2020b; Kosmack et al., 2008; Rice, 1980; Utaki, 2010; Yang et al., 2016), programmes (Bracmort et al., 2014; Chianese et al., 2015) or partnerships (Industry news, 2004) refer primarily to the **fossil fuel industry** (e.g. from CH₄ recovery from coal mines to purification for injection into the natural gas grid). **Coal bed methane** can be used as a high-energy fuel. The concentration of CH₄ is, however, critical for its use (Liu et al., 2020a). Technologies for separating the gas mixtures from coal mines with low concentrations of CH₄ have been developed (Moiseev et al., 2016b). Besides fossil fuels, CH₄ is emitted from a number of sources (Bracmort et al., 2014), linked directly or indirectly to human activities. Some of these sources can be used for the controlled production of biomethane:

1. **animal residues** are a source of biogas (Wentworth et al., 1979), especially when coupled to digesters, a proven technology for the capture and use of the biomethane produced (Chianese et al., 2015);
2. the recovery and treatment of CH₄ generated by sanitary **landfills** is also a commercial reality (Rice, 1980), including the production of pipeline-grade biomethane from landfill gas (Cavenati et al., 2005) ⁽²⁾.

Hydrate-based gas separation technologies for CH₄ recovery from biogas have also gained prominence (Zang and Liang, 2018).

2.1. Current processes and technologies for methane concentration and upgrading

Various **physico-chemical (adsorption, absorption, cryogenic and membrane separations) and biological (in situ and ex situ) processes** exist for biogas upgrading (Kapoor et al., 2019). Important considerations for biogas cleaning and upgrading technologies include product purity and impurities; CH₄ recovery and loss; upgrading efficiency; and the investment and operating costs. In addition, the potential utilisation of the biogas and the corresponding requirements for gas quality are also important. Recommendations on the appropriate technology rely on comparisons between the technical features of the **upgrading technologies**, the specific requirements for different gas utilisations and the relevant investment and operating costs (Sun et al., 2015). Suitable technologies for CH₄ upgrading include **membrane separation, chemical adsorption** (Vignali and Vitale, 2017) and water scrubbing ⁽³⁾.

Adsorbents are already used for adsorption and enrichment of CH₄ from low-concentration gas mixtures (Liu et al., 2020a). Solid adsorbents have also been developed and tested to capture CH₄ emissions, ideally under atmospheric conditions, with the objective of developing new yet artificial remediation technologies (Delgado et al., 2018). Indeed, novel nanomaterials are being used as adsorbents for CO₂ and CH₄ with the objective of mitigating their emissions (Alonso et al., 2017). Nanomaterials used for CH₄ adsorption are divided into non-carbonaceous materials (e.g. zeolites, metal-organic frameworks and porous polymers) and carbonaceous materials (e.g. activated carbons, ordered porous carbons and activated carbon fibres) (Choi et al., 2016). Carbon-based materials, such as activated carbons, are promising as **storage** adsorbents for natural gas because of their high surface area, high porosity and high volumetric storage capacity (Poomisitiporn et al., 2016). **Technologies based on adsorbents** include the vacuum pressure swing adsorption unit (Cavenati et al., 2005; Utaki, 2010; Yang et al., 2018) and adsorption with displacement chromatography (Yang et al., 2016) for the recovery of ventilation air methane from coal mines. **Membranes for gas separation** have also been studied and are already used for the separation of CH₄ from air. Economics are dictated by the CH₄ concentration, making the technology less suited for mine ventilation exhaust (2 vol. per cent or less) than for

⁽²⁾ 'Horizon 2020 Store&Go project demonstrates new methanation technologies at three pilot sites', available online at: <https://www.storeandgo.info/demonstration-sites/>, accessed 4.9.2020.

⁽³⁾ Ofori-Boateng, C., Kwofie, E.M., 'Water scrubbing: a better option for biogas purification for effective storage', *World Applied Sciences Journal*, 5, pp. 122–125, available online at: https://www.researchgate.net/publication/265180428_Water_Scrubbing_A_Better_Option_for_Biogas_Purification_for_Effective_Storage, accessed 16.10.2020; Chandra, R., Vijay, V.K., Subbarao, P.M.V., 'Vehicular quality biomethane production from biogas by using an automated water scrubbing system', *International Scholarly Research Notices*, 2012, 904167, available online at: <https://www.hindawi.com/journals/isrn/2012/904167/>, accessed 4.9.2020.

gob hole ⁽⁴⁾ (mining cavities that have collapsed) methane-air mixtures with higher CH₄ concentrations (from 30 to 100 vol. per cent) (Garcia and Cervik, 1988). Each process is site and case specific (Kapoor et al., 2019).

2.2. Technologies behind biomethane production

Whether produced from manure or landfill or as fossil fuel, CH₄ is the product of biological processes and is derived from biological material. Digesters are a proven emission reduction technology and allow the capture and use of CH₄ produced by the anaerobic digestion of manures (Chianese et al., 2015). Further, anaerobic digestion technology and its economics make biogas production from selected animal residues worthy of development (Wentworth et al., 1979) and a reality ⁽⁵⁾. Conventional anaerobic digestion, together with landfill gas recovery, have also been investigated for reducing emissions from municipal solid waste. Improved technology for enhancing CH₄ recovery from landfills appears to have greater potential than using conventional sludge digestion technology (Wise et al., 1981). It remains that these technologies rely on microorganisms and could therefore benefit from a biotrickling filter (BTF), which allows for the enrichment of microorganisms (Cassarini et al., 2019a) (see Chapter 3).

⁽⁴⁾ Saki, S.A., *Gob ventilation borehole design and performance optimisation for longwall coal mining using computational fluid dynamics*, PhD thesis, January 2016, doi:10.13140/RG.2.1.5119.9129.


⁽⁵⁾ Strauch, S., Krassowski, J., Singhal, A., *Biomethane Guide for Decision Makers – Policy guide on biogas injection into the natural gas grid*, WP2/D2.3, April 2013, available online at: https://www.dena.de/fileadmin/dena/Dokumente/Themen_und_Projekte/Erneuerbare_Energien/GreenGasGrids/Policy_Guide_for_Decision_Makers.pdf, accessed 4.9.2020.

3. Towards biological remediation strategies

Digestion stems from methanogenesis, a process in which microorganisms synthesise methane. This process takes place in most environments (soil (Gutknecht et al., 2006), including permafrost (Li et al., 2020), and water (Martinez-Cruz et al., 2017)). Together with other processes (such as nitrification and denitrification), methanogenesis influences atmospheric chemistry (Gutknecht et al., 2006).

Methanotrophs are bacteria capable of using CH_4 as their sole carbon source (Strong et al., 2015). Three major functional types of methanotrophs are known to consume CH_4 : (1) anaerobic methane-oxidising archaea, also called anaerobic methanotrophs (ANME); (2) anaerobic methane-oxidising bacteria (NC10 phylum); and (3) aerobic methane-oxidising bacteria (Proteobacteria and Verrucomicrobia). These conduct different types of methanotrophy (Kalyuzhnaya et al., 2018), as shown in Figure 1.

Figure 1. Illustration of CH_4 oxidation (vertical air-water cross-section. The blue line represents the air-water interface). Anaerobic CH_4 oxidation is visualised by the red and green consortium, representing ANME and sulfate-reducing bacteria, respectively.



Source: WUR ⁽⁶⁾.

Considering CH_4 as a GHG, some of the above processes could be artificially enhanced and become instrumental in the reduction of emissions and atmospheric concentrations. These could be complemented by natural and industrial processes and technologies.

It should be noted that all biological oxidation processes oxidise CH_4 into CO_2 , which is also a GHG, but with a significantly lower warming potential than CH_4 .

Some of the newly discovered physiological types of anaerobic CH_4 oxidisers challenge the view of obligate syntrophy for the anaerobic oxidation of methane (AOM) (Bhattarai et al., 2019). It appears, for instance, that

⁽⁶⁾ Wageningen University and Research, 'Anaerobic oxidation of methane', available online at: <https://www.wur.nl/en/show/Anaerobic-oxidation-of-methane.htm>, accessed 4.9.2020.

ANME can also produce CH₄, but only during net CH₄ oxidation (i.e. enzymatic back-flux) (Timmers et al., 2017).

AOM is catalysed by ANME via a **reverse and modified methanogenesis pathway**. The reversibility of the methanogenesis pathway and essential differences between ANME and methanogens have been subject to **genetic investigations** (Timmers et al., 2017). Other potential technologies for CH₄ consumption linked to genetic investigations include the following:

- Next-generation sequencing (NGS) opens up possibilities for improving our knowledge of microorganisms (Cruaud et al., 2014).
- The *pmoA* gene sequence diversity highlights the diversity of cultivated and uncultivated aerobic methanotrophic bacteria (Knief, 2015).

This knowledge will enable progress to be made towards **engineering** native methanotrophs for aerobic and anaerobic CH₄ utilisation and synthetic methylotrophs for methanol utilisation (Bennett et al., 2018) or the production of other chemicals (Lieven et al., 2018).

The study of hydrate environments has allowed the recovery of 124 379 bacterial and 130 351 archaeal reads. This has provided insights into the **distributions and capacities of microbial communities** (Lin et al., 2014).

Regarding microbial capacities, the reduction of sulfate, nitrite/nitrate and iron/manganese can be coupled with AOM (Luo et al., 2019). Net AOM is actually exergonic when coupled to an external electron acceptor (Timmers et al., 2017). This redox process leads to the formation of two main biomineral by-products: calcium carbonates and iron sulfides (Wrede et al., 2013a). However, methanotrophic bacteria have numerous potential **biotechnological applications**, thereby generating value while using CH₄ as a carbon source (Strong et al., 2015). These include **microbial fuel cells** (McAnulty et al., 2017) for electricity production; metal-AOM (He et al., 2018) for the production of iron and manganese, carbonates (Chen et al., 2014; Wrede et al., 2013a,b), chromate (Lv et al., 2019), copper (Kampman et al., 2014; Ro and Rosenzweig, 2018; Semrau et al., 2010; Ve et al., 2012), nitrous compounds (He et al., 2018; Kampman et al., 2014; Kolb and Horn, 2012; Li et al., 2020; Lin et al., 2014; Luo et al., 2019; McAnulty et al., 2017; Mei et al., 2019; Menyailo et al., 2008; Modin et al., 2008; Ren et al., 2018; Rissanen et al., 2018; Shi et al., 2019; Strong et al., 2015; Sun et al., 2013; Timmers et al., 2017; Urbanová et al., 2011; Vaksmaa et al., 2016; van Grinsven et al., 2020; Winkel et al., 2018; Xie et al., 2018), sulfur (Aromokeye et al., 2020; Bar-Or et al., 2017; Bhattarai et al., 2017; Bhattarai et al., 2018, 2019; Bomberg et al., 2015; Cassarini et al., 2019a,b,c; Chen et al., 2014; Cruaud et al., 2015; Gutknecht et al., 2006; Hatzenpichler et al., 2016; He et al., 2018; Lin et al., 2014; Luo et al., 2019; McAnulty et al., 2017; Mei et al., 2019; Ren et al., 2018; Scheller et al., 2016; Skennerton et al., 2017; Timmers et al., 2017; Valenzuela et al., 2017; van Grinsven et al., 2020; Vigneron et al., 2019; Winkel et al., 2018; Wrede et al., 2013a,b) or the transformation of CH₄ in fuels and chemicals (Bennett et al., 2018), lactic acid (Fei et al., 2020), methanol (Hur et al., 2017; Ito et al., 2018; Wrede et al., 2013b) and acetate (McAnulty et al., 2017); and other numerous applications (Strong et al., 2015).

Regarding their distribution, ANME groups occur widely in association with gas hydrates, cold seeps and organic-rich thermal sediments, as well as in anoxic water bodies such as the Black Sea and Cariaco Basin (Jiao et al., 2012). Methanotrophy (at large) takes place in various environments, such as wetlands (Gutknecht et al., 2006) and the deep oceans and marine sediments (Jiao et al., 2012). However, climate change and global warming stem from the accumulation of GHGs in the **atmosphere**. There is thus an interest in focusing on the capture of CH₄ under conditions related to the atmospheric environment.

- The microbiota of acidic wetlands seem to consume both atmospheric CH₄ (likely through *Methylocystis*-related species) and atmospheric N₂O (through denitrifier communities) (Kolb and Horn, 2012).
- Although afforestation is beneficial for atmospheric CO₂ assimilation, it shows negative effects on CH₄ uptake from well-aerated grassland in Siberia (Menyailo et al., 2008).
- Airborne methanotrophs are also found in air and rainwater. Once enriched, they have been successfully used for the degradation of atmospheric CH₄ (Šantl-Temkiv et al., 2013).

A BTF enables the enrichment of microorganisms, such as those mediating AOM (Cassarini et al., 2019a; Ito et al., 2018). This technology may potentially support natural processes in the consumption of CH₄. A biotrickling filtration system composed of methane-consuming bacteria has been modelled to assess the utility of these systems in removing CH₄ from the atmosphere. While the current atmospheric CH₄ concentration does not allow use of this technology, this approach could be applied to other environments with higher CH₄ concentrations,

such as landfills or factory farms (Yoon et al., 2009). Although there are indeed limitations of ANME laboratory-based cultivation systems, as well as advantages and potential improvements to be made (Bhattarai et al., 2019), such systems enable the study of the characteristics of CH₄ bio-oxidation and methane-oxidising microorganisms (Mei et al., 2019), facilitating progress in their development for the consumption of CH₄.

The biotrickling filtration system (Yoon et al., 2009) is one of the few technologies to have been assessed from the perspectives of its **CH₄ emission reduction potential** (Annachhatre and Khanna, 1987; Aromokeye et al., 2020; Hur et al., 2017; Industry news, 2004; Kosmack et al., 2008; Li et al., 2020; Lima et al., 2008; Liu et al., 2020b; Mei et al., 2019; Sanchis-Perucho et al., 2020; Serrano-Silva et al., 2014; Shi et al., 2020; Utaki, 2010; Vaksmaa et al., 2016; Valenzuela et al., 2017; Winkel et al., 2018; Yang et al., 2016; Yoon et al., 2009; Zhang et al., 2019) and its **economics/costs** (Fei et al., 2020; Yoon et al., 2009). This demonstrates that such CH₄ emission mitigation technologies are at an early stage of development. However, innovation is ongoing, for instance work on CH₄ cracking/pyrolysis for the production of clean hydrogen and solid carbon for storage ⁽⁷⁾.

⁽⁷⁾ TNO, 'EMBER methane pyrolysis technology produces hydrogen without CO₂ emissions', available online at: <https://www.tno.nl/en/focus-areas/energy-transition/roadmaps/towards-co2-neutral-fuels-and-feedstock/hydrogen-for-a-sustainable-energy-supply/optimising-production-hydrogen/ember-methane-pyrolysis/>, accessed 4.9.2020.

4. Recommendations and open questions

This review, together with its accompanying fact sheet ⁽⁸⁾, identified the following areas for further investigation.

4.1. Knowledge gathering and sharing

- Develop a list of definitions, abbreviations and acronyms linked to the topic.
- (Further support ongoing efforts ⁽⁹⁾ to) Identify spatial/geographical distributions of GHG sources to better understand the contributions of each GHG/source/sector to global warming, possibly through satellite imaging.
- Identify GHG consumption processes and technologies, also possibly linked to GHG sources.
- (Linked to the discussion on afforestation) Assess the environment and sources of emissions at GWP level, instead of focusing on individual GHGs separately.
- This review is significant in terms of scientific process. The narrow literature review allowed for the (manual) definition of a mapping structure (see the structure presented above). By using text-mining tools, this structure can now be applied to a broader set of scientific documents. Such scaling up will enrich the mapping and help to enhance the mapping structure. This enhanced knowledge gathering will in turn lead to better mapping of ongoing activities and thus mapping of the gaps within.

4.2. Specific emissions and atmospheric concentration mitigation options

- Develop more accurate CH₄ emission measurement methodologies to enable convergence of the current bottom-up and top-down approaches in order to significantly reduce uncertainties ⁽⁸⁾.
- Identify procedures to identify CH₄ leakages in industrial installations and upgrade installations to prevent these leakages and promote the recovery of the leaked CH₄ ⁽⁸⁾.
- Set up monitoring procedures for former fossil fuel exploitation sites (decommissioned coal mines, decommissioned oil and gas wells) to ensure long-term control of potential CH₄ emissions ⁽⁸⁾.
- Increase knowledge of CH₄ production as a by-product of agricultural processes (ruminant animals, flooded rice fields, animal waste and biomass) in order to control and mitigate such emissions.
- Investigate biological processes for joint AOM and denitrification, leading to both N₂O and CH₄ consumption.
- Investigate the feasibility of developing the air capture/consumption of GHGs besides and/or coupled to CO₂.
- Identify possible/more targeted topics for future fact sheets, such as CH₄ cracking. This process was not identified through this literature review, although it has potential for the decarbonisation of the hydrogen production process.

4.3. Ethical aspects

- Reflect on (ongoing) progress, objectives and, above all, risks linked to gene editing for modifying microorganisms. Considering the limited knowledge of methanogenesis and natural processes for CH₄ consumption (reverse methanogenesis and methanotrophy), such techniques may have significant potential for addressing (natural) CH₄ emissions. Gene editing, however, remains a sensitive topic.
- Carry out a benefit–risk analysis of the use of genetically modified organisms in controlled and uncontrolled conditions to mitigate CH₄ emissions compared with the consequences of CH₄ emissions for climate change.

⁽⁸⁾ JRC, 'The special case of methane emissions, a potent greenhouse gas', JRC121991, 2020.

⁽⁹⁾ European Space Agency, 'Mapping methane emissions on a global scale', 2020, available at: https://www.esa.int/Applications/Observing_the_Earth/Copernicus/Sentinel-5P/Mapping_methane_emissions_on_a_global_scale, accessed 20.10.2020; IEA, 'Global methane emissions from oil and gas', 2020, available at: <https://www.iea.org/articles/global-methane-emissions-from-oil-and-gas>, accessed 20.10.2020; NASA, 'New 3D view of methane tracks sources and movement around the globe', 2020, available at: <https://www.nasa.gov/feature/goddard/2020/new-3d-view-of-methane-tracks-sources-and-movement-around-the-globe>, accessed 16.10.2020.

5. Conclusions

Methane is a primary energy source that fuels one-quarter of the modern economy. It is mostly burned to produce heat and power. This process leads to massive CO₂ emissions that trigger climate change. As well as producing CO₂ emissions, CH₄ itself is also a powerful GHG and therefore is worthy of focused attention.

Methane emissions should be prevented because of CH₄'s high GWP. The CH₄ concentration in the atmosphere has doubled since the start of the industrial age in 1750, probably because of leakages and its production as a by-product of the fossil fuel industries, as well as its production as a result of biological processes in agriculture. Efforts should be made to control and mitigate these emissions.

This review identifies processes and technologies for the biological oxidation of CH₄, as well as for CH₄ recovery and controlled biomethane production. It provides a list of possible options for CH₄ emission mitigation, to slow down the increase in CH₄ atmospheric concentrations or even reduce them. In order to foster the deployment of such technologies, further (techno-economic) analysis is recommended. This, however, requires a change of mindset: while CH₄ can be useful, it is also harmful to the environment.

Further, this review, in itself, is significant in terms of scientific process: The narrow literature review allowed the manual definition of a mapping structure. By using text-mining tools, this structure can now be applied to a broader set of scientific documents. Such scaling up will enrich the mapping and help to enhance the mapping structure. This enhanced knowledge gathering will in turn lead to better mapping of ongoing activities and thus a mapping of the gaps within.

References

The following references were retrieved from Scopus using searches for CH₄ capture-related articles.

With the objective of assessing the field, the (generic) query TITLE-ABS-KEY ((methan OR methane OR ch4) AND (greenhouse AND gases OR greenhouse AND gas OR ghg)) retrieved 14 338 results, continuously increasing from four in 1980 to 1 400 in 2019.

For the purpose of this study, the literature search was narrowed along two axes:

- With the objective of identifying technologies for 'CH₄ removal', the query TITLE-ABS-KEY ((methan OR methane OR ch4) W/5 (capture OR recovery OR adsorption) AND technology) retrieved 896 results.
- With the objective of investigating (human enhancement of) the natural process of 'methanotrophy', the query ALL (artific* W/5 methano*) retrieved 521 results.

The large number of articles retrieved in the initial searches was reduced by focusing on those listed below:

- those from the 'CH₄ removal' list that referred to technology in their abstract;
- those from the 'methanotrophy' list that referred to methanotrophs in their abstract; the relevance of the remaining articles was then assessed.

In addition to the 109 references listed below, eight additional references are provided as footnotes in the report. These eight references were not identified through the above searches; however, they do provide information of relevance for this report.

Abbott, L.J., Hart, K.E., Larsen, G.S., Lin, P., Colina, C.M., 'Novel nanoporous polymers: In silico design', 2011, paper presented at 242nd ACS National Meeting, Denver, CO, 28 August–1 September.

Air Liquide targets landfill gas-to-energy, 2007, *Process Engineering*, 88(2), pp. 43.

Ajayi, A., Shehu, H., Gobina, E., 'Recovery of VOC from onshore and offshore shuttle tankers using structured membranes', in Ao, S.I., Gelman, L., Hukins, D.W.L., Hunter, A., Korsunsky, A.M. (eds.), *Proceedings of the 2015 World Congress on Engineering and Computer Science (WCECS 2015)*, 21–23 October 2015, San Francisco, USA, Lecture notes in engineering and computer science, 2220, Newswood Limited, Hong Kong, pp. 637–642.

Alonso, A., Moral-Vico, J., Abo Markeb, A., Busquets-Fité, M., Komilis, D., Puentes, V., Sánchez, A., Font, X., 'Critical review of existing nanomaterial adsorbents to capture carbon dioxide and methane', 2017, *Science of the Total Environment*, 595, doi:10.1016/j.scitotenv.2017.03.229.

Annachatre, A.P., Khanna, P., 'Methane recovery from water hyacinth through whole-cell immobilization technology', 1987, *Biotechnology and Bioengineering*, 29, doi:10.1002/bit.260290703.

Aromokeye, D.A., Kulkarni, A.C., Elvert, M., Wegener, G., Henkel, S., Coffinet, S., Eickhorst, T., Oni, O.E., Richter-Heitmann, T., Schnakenberg, A., Taubner, H., Wunder, L., Yin, X., Zhu, Q., Hinrichs, K.-U., Kasten, S., Friedrich, M.W., 'Rates and microbial players of iron-driven anaerobic oxidation of methane in methanic marine sediments', 2020, *Frontiers in Microbiology*, 10, doi:10.3389/fmicb.2019.03041.

Bar-Or, I., Elvert, M., Eckert, W., Kushmaro, A., Vigderovich, H., Zhu, Q., Ben-Dov, E., Sivan, O., 'Iron-coupled anaerobic oxidation of methane performed by a mixed bacterial-archaeal community based on poorly reactive minerals', 2017, *Environmental Science and Technology*, 51, doi:10.1021/acs.est.7b03126.

Bennett, R.K., Steinberg, L.M., Chen, W., Papoutsakis, E.T., 'Engineering the bioconversion of methane and methanol to fuels and chemicals in native and synthetic methylotrophs', 2018, *Current Opinions in Biotechnology*, 50, doi:10.1016/j.copbio.2017.11.010.

Bhattarai, S., Cassarini, C., Gonzalez-Gil, G., Egger, M., Slomp, C.P., Zhang, Y., Esposito, G., Lens, P.N.L., 'Anaerobic methane-oxidizing microbial community in a coastal marine sediment: Anaerobic methanotrophy dominated by ANME-3', 2017, *Microbial Ecology*, 74, doi:10.1007/s00248-017-0978-y.

Bhattarai, S., Cassarini, C., Rene, E.R., Kümmel, S., Esposito, G., Lens, P.N.L., 'Enrichment of ANME-2 dominated anaerobic methanotrophy from cold seep sediment in an external ultrafiltration membrane bioreactor', 2018, *Engineering in Life Sciences*, 18, doi:10.1002/elsc.201700148.

Bhattarai, S., Cassarini, C., Lens, P.N.L., 'Physiology and distribution of archaeal methanotrophs that couple anaerobic oxidation of methane with sulfate reduction', 2019, *Microbiology and Molecular Biology Reviews*, 83, doi:10.1128/MMBR.00074-18.

- Bomberg, M., Nyyssönen, M., Pitkänen, P., Lehtinen, A., Itävaara, M., 'Active microbial communities inhabit sulphate-methane interphase in deep bedrock fracture fluids in Olkiluoto, Finland', 2015, *BioMed Research International*, 2015, doi:10.1155/2015/979530.
- Bracmort, K., Ramseur, J.L., McCarthy, J.E., Folger, P., Marples, D.J., 'Methane capture: Options for greenhouse gas emission reduction', in Davis, G.L. (ed.), *New trends in environmental science*, Nova Science Publishers, New York, 2014, pp. 211–228.
- Casey, E., Rishell, S., Glennon, B., Hamer, G., 'Engineering aspects of a mixed methanotrophic culture in a membrane-aerated biofilm reactor', 2004, *Water Science and Technology*, 49.
- Cashman, S., Ma, X., Mosley, J., Garland, J., Crone, B., Xue, X., 'Energy and greenhouse gas life cycle assessment and cost analysis of aerobic and anaerobic membrane bioreactor systems: Influence of scale, population density, climate, and methane recovery', 2018, *Bioresource Technology*, 254, doi:10.1016/j.biortech.2018.01.060.
- Cassarini, C., Bhattarai, S., Rene, E.R., Vogt, C., Musat, N., Esposito, G., Lens, P.N.L., 'Enrichment of anaerobic methanotrophs in biotrickling filters using different sulfur compounds as electron acceptor', 2019a, *Environmental Engineering Science*, 36, doi:10.1089/ees.2018.0283.
- Cassarini, C., Rene, E.R., Bhattarai, S., Vogt, C., Musat, N., Lens, P.N.L., 'Anaerobic methane oxidation coupled to sulfate reduction in a biotrickling filter: Reactor performance and microbial community analysis', 2019b, *Chemosphere*, 236, doi:10.1016/j.chemosphere.2019.07.021.
- Cassarini, C., Zhang, Y., Lens, P.N.L., 'Pressure selects dominant anaerobic methanotrophic phylotype and sulfate reducing bacteria in coastal marine Lake Grevelingen sediment', 2019c, *Frontiers in Environmental Science*, 6, doi:10.3389/fenvs.2018.00162.
- Cavenati, S., Grande, C.A., Rodrigues, A.E., 'Upgrade of methane from landfill gas by pressure swing adsorption', 2005, *Energy Fuels*, 19, doi:10.1021/ef050072h.
- Chen, Y., Li, Y.-L., Zhou, G.-T., Li, H., Lin, Y.-T., Xiao, X., Wang, F.-P., 'Biomining mediated by anaerobic methane-consuming cell consortia', 2014, *Scientific Reports*, 4, doi:10.1038/srep05696.
- Chianese, D.S., Kim, D.T., Daugherty, D.D., 'An avenue for mitigation: Lessons from California's offsets program', 2015, ASABE 1st Climate Change Symposium: Adaptation and Mitigation Conference Proceedings 152124156, pp. 140–142.
- Choi, P.-S., Jeong, J.-M., Choi, Y.-K., Kim, M.-S., Shin, G.-J., Park, S.-J., 'A review: Methane capture by nanoporous carbon materials for automobiles', 2016, *Carbon Letters*, 17, doi:10.5714/CL.2016.17.1.018.
- Cramer, T.A., Johnson, D.W., Urynowicz, M.A., 'Membrane gas transfer of methane and carbon dioxide in submerged coal deposits', 2009, *Environmental Technology*, 30, doi:10.1080/09593330802421425.
- Cruaud, P., Vigneron, A., Lucchetti-Miganeh, C., Ciron, P.E., Godfroy, A., Cambon-Bonavita, M.-A., 'Influence of DNA extraction method, 16S rRNA targeted hypervariable regions, and sample origin on microbial diversity detected by 454 pyrosequencing in marine chemosynthetic ecosystems', 2014, *Applied and Environmental Microbiology*, 80, doi:10.1128/AEM.00592-14.
- Cruaud, P., Vigneron, A., Pignet, P., Caprais, J.-C., Lesongeur, F., Toffin, L., Godfroy, A., Cambon-Bonavita, M.-A., 'Microbial communities associated with benthic faunal assemblages at cold seep sediments of the Sonora Margin, Guaymas Basin', 2015, *Frontiers in Marine Science*, 2, doi:10.3389/fmars.2015.00053.
- Delgado, B., Lagace, R., Godbout, S., Valverde, J.L., Giroir-Fendler, A., Ramirez, A.A., 'Chemical and adsorptive characterization of adsorbents to capture greenhouse gases under atmospheric conditions of temperature and pressure', 2018, *ACS Symposium Series*, 1302, doi:10.1021/bk-2018-1302.ch006.
- Fei, Q., Liang, B., Tao, L., Tan, E.C.D., Gonzalez, R., Henard, C.A., Guarnieri, M.T., 'Biological valorization of natural gas for the production of lactic acid: Techno-economic analysis and life cycle assessment', 2020, *Biochemical Engineering Journal*, 158, doi:10.1016/j.bej.2020.107500.
- Garcia, F., Cervik, J., *Review of membrane technology for methane recovery from mining operations*, Information circular volume 9174, U.S. Department of the Interior, Bureau of Mines, Washington, D.C., 1988.
- Gutknecht, J.L.M., Goodman, R.M., Balser, T.C., 'Linking soil process and microbial ecology in freshwater wetland ecosystems', 2006, *Plant and Soil*, 289, doi:10.1007/s11104-006-9105-4.

- Hatzenpichler, R., Connon, S.A., Goudeau, D., Malmstrom, R.R., Woyke, T., Orphan, V.J., 'Visualizing in situ translational activity for identifying and sorting slow-growing archaeal – bacterial consortia', 2016, *Proceedings of the National Academy of Sciences U.S.A.*, 113, doi:10.1073/pnas.1603757113.
- He, Z., Zhang, Q., Feng, Y., Luo, H., Pan, X., Gadd, G.M., 'Microbiological and environmental significance of metal-dependent anaerobic oxidation of methane', 2018, *Science of the Total Environment*, 610–611, doi:10.1016/j.scitotenv.2017.08.140.
- Henares, M., Izquierdo, M., Penya-Roja, J.M., Martínez-Soria, V., 'Comparative study of degassing membrane modules for the removal of methane from expanded granular sludge bed anaerobic reactor effluent', 2016, *Separation and Purification Technology*, 170, doi:10.1016/j.seppur.2016.06.024.
- Hur, D.H., Na, J.-G., Lee, E.Y., 'Highly efficient bioconversion of methane to methanol using a novel type I *Methylobacter* sp. DH-1 newly isolated from brewery waste sludge', 2017, *Journal of Chemical Technology and Biotechnology*, 92, doi:10.1002/jctb.5007.
- Industry news: U.S. joins international methane partnership, 2004, *Pollution Engineering*, 36, 10, pp. 10.
- Ito, H., Kondo, R., Yoshimori, K., Kamachi, T., 'Methane hydroxylation with water as an electron donor under light irradiation in the presence of reconstituted membranes containing both photosystem II and a methane monooxygenase', 2018, *ChemBioChem*, 19, doi:10.1002/cbic.201800324.
- Jiang, H., Duan, C., Luo, M., Xing, X.-H., 'Enrichment and characteristics of mixed methane-oxidizing bacteria from a Chinese coal mine', 2016, *Applied Microbiology and Biotechnology*, 100, doi:10.1007/s00253-016-7738-7.
- Jiao, N., Zhang, C., 'Primary production and carbon cycling in the deep sea: Archaeal carbon fixation and methane oxidation', in Bailey, D.R., Howard, S.E. (eds), *Deep-sea: Marine biology, geology, and human impact*, Nova Science Publishers, New York, 2012, pp. 149–170.
- Kalyuzhnaya, M.G., Collins, D., Chistoserdova, L., 'Microbial cycling of methane', in Schmidt, T.M. (ed.), *Encyclopedia of Microbiology*, Elsevier, Cambridge, 2018, pp. 115–124.
- Kampman, C., Temmink, H., Hendrickx, T.L.G., Zeeman, G., Buisman, C.J.N., 'Enrichment of denitrifying methanotrophic bacteria from municipal wastewater sludge in a membrane bioreactor at 20 °C', 2014, *Journal of Hazardous Material*, 274, doi:10.1016/j.jhazmat.2014.04.031.
- Kapoor, R., Ghosh, P., Kumar, M., Vijay, V.K., 'Evaluation of biogas upgrading technologies and future perspectives: A review', 2019, *Environmental Science and Pollution Research*, 26, doi:10.1007/s11356-019-04767-1.
- Knief, C., 'Diversity and habitat preferences of cultivated and uncultivated aerobic methanotrophic bacteria evaluated based on pmoA as molecular marker', 2015, *Frontiers in Microbiology*, 6, doi:10.3389/fmicb.2015.01346.
- Kolb, S., Horn, M.A., 'Microbial CH₄ and N₂O consumption in acidic wetlands', 2012, *Frontiers in Microbiology*, 3, 78, doi:10.3389/fmicb.2012.00078.
- Kosmack, D.A., Zak, K.P., 'First U.S. field trial of oxidation technology for coal mine ventilation air methane', 2008, paper presented at 25th Annual International Pittsburgh Coal Conference 2008, 29 September–2 October, Pittsburgh, PA.
- Kostoglou, N., Koczwara, C., Prehal, C., Terziyska, V., Babic, B., Matovic, B., Constantinides, G., Tampaxis, C., Charalambopoulou, G., Steriotis, T., Hinder, S., Baker, M., Polychronopoulou, K., Dumanidis, C., Paris, O., Mitterer, C., Rebholz, C., 'Nanoporous activated carbon cloth as a versatile material for hydrogen adsorption, selective gas separation and electrochemical energy storage', 2017, *Nano Energy*, 40, doi:10.1016/j.nanoen.2017.07.056.
- Kwon, M.J., Beulig, F., Ilie, I., Wildner, M., Küsel, K., Merbold, L., Mahecha, M.D., Zimov, N., Zimov, S.A., Heimann, M., Schuur, E.A.G., Kostka, J.E., Kolle, O., Hilke, I., Göckede, M., 'Plants, microorganisms, and soil temperatures contribute to a decrease in methane fluxes on a drained Arctic floodplain', 2017, *Global Change Biology*, 23, doi:10.1111/gwebcb.13558.
- Leisle, A.V., Kovalski, E.R., 'Complex extraction of methane and coal from thick coal seams', 2016, *Research Journal of Pharmaceutical, Biological and Chemical Science*, 7.
- Li, M., Xue, L., Zhou, B., Duan, J., He, Z., Wang, X., Xu, X., Yang, L., 'Effects of domestic sewage from different sources on greenhouse gas emission and related microorganisms in straw-returning paddy fields', 2020, *Science of the Total Environment*, 718, doi:10.1016/j.scitotenv.2020.137407.

- Lieven, C., Herrgård, M.J., Sonnenschein, N., 'Microbial methylotrophic metabolism: Recent metabolic modeling efforts and their applications in industrial biotechnology', 2018, *Biotechnology Journal*, 13, doi:10.1002/biot.201800011.
- Lima, I.B.T., Ramos, F.M., Bambace, L.A.W., Rosa, R.R., 'Methane emissions from large dams as renewable energy resources: A developing nation perspective', 2008, *Mitigation and Adaptation Strategies for Global Change*, 13, doi:10.1007/s11027-007-9086-5.
- Lin, L.-H., Wu, L.-W., Cheng, T.-W., Tu, W.-X., Lin, J.-R., Yang, T.F., Chen, P.-C., Wang, Y., Wang, P.-L., 'Distributions and assemblages of microbial communities along a sediment core retrieved from a potential hydrate-bearing region offshore southwestern Taiwan', 2014, *Journal of Asian Earth Sciences*, 92, doi:10.1016/j.jseaes.2014.02.014.
- Liu, H., Ding, W., Zhou, F., Yang, G., Du, Y., 'An overview and outlook on gas adsorption: For the enrichment of low concentration coalbed methane', 2020a, *Separation Science and Technology*, 55, doi:10.1080/01496395.2019.1585454.
- Liu, W., Xiao, H., Ma, H., Li, Y., Adyel, T.M., Zhai, J., 'Reduction of methane emissions from manganese-rich constructed wetlands: Role of manganese-dependent anaerobic methane oxidation', 2020b, *Chemical Engineering Journal*, 387, doi:10.1016/j.cej.2019.123402.
- Luo, J.-H., Wu, M., Liu, J., Qian, G., Yuan, Z., Guo, J., 'Microbial chromate reduction coupled with anaerobic oxidation of methane in a membrane biofilm reactor', 2019, *Environment International*, 130, doi:10.1016/j.envint.2019.104926.
- Lv, P.-L., Shi, L.-D., Wang, Z., Rittmann, B., Zhao, H.-P., 'Methane oxidation coupled to perchlorate reduction in a membrane biofilm batch reactor', 2019, *Science of the Total Environment*, 667, doi:10.1016/j.scitotenv.2019.02.330.
- Mahmoudian, L., Rashidi, A., Dehghani, H., Rahighi, R., 'Single-step scalable synthesis of three-dimensional highly porous graphene with favorable methane adsorption', 2016, *Chemical Engineering Journal*, 304, doi:10.1016/j.cej.2016.07.015.
- Martinez-Cruz, K., Leewis, M.-C., Herriott, I.C., Sepulveda-Jauregui, A., Anthony, K.W., Thalasso, F., Leigh, M.B., 'Anaerobic oxidation of methane by aerobic methanotrophs in sub-Arctic lake sediments', 2017, *Science of the Total Environment*, 607–608, doi:10.1016/j.scitotenv.2017.06.187.
- McAnulty, M.J., Poosarla, V.G., Kim, K.-Y., Jasso-Chávez, R., Logan, B.E., Wood, T.K., 'Electricity from methane by reversing methanogenesis', 2017, *Nature Communications*, 8, doi:10.1038/ncomms15419.
- Mei, J., Wu, Y., Qian, F., Chen, C., Shen, Y., Zhao, Y., 'Methane-oxidizing microorganism properties in landfills', 2019, *Polish Journal of Environmental Studies*, 28, doi:10.15244/pjoes/96239.
- Menyailo, O.V., Hungate, B.A., Abraham, W.-R., Conrad, R., 'Changing land use reduces soil CH₄ uptake by altering biomass and activity but not composition of high-affinity methanotrophs', 2008, *Global Change Biology*, 14, doi:10.1111/j.1365-2486.2008.01648.x.
- Modin, O., Fukushima, K., Nakajima, F., Yamamoto, K., 'A membrane biofilm reactor achieves aerobic methane oxidation coupled to denitrification, AME-D, with high efficiency', 2008, *Water Science and Technology*, 58, doi:10.2166/wst.2008.648.
- Moiseev, V.A., Andrienko, V.G., Piletskii, V.G., Chentsov, A.V., 'Technology of efficient recovering of highly concentrated methane and mechanical impurities from ventilation and degasification emissions from coal mines', 2016a, *International Journal of Applied Engineering Research*, 11.
- Moiseev, V.A., Piletskii, V.G., Andrienko, V.G., Tailakov, O.V., Chentsov, A.V., 'The technological and design solutions for energy vortex gas separation of the ventilation and degasification emissions from coal mines using gradient separators CJSC COMPOMASH-TEK', 2016b, *Global Journal of Pure and Applied Mathematics*, 12.
- Myung, J., Flanagan, J.C.A., Waymouth, R.M., Criddle, C.S., 'Expanding the range of polyhydroxyalkanoates synthesized by methanotrophic bacteria through the utilization of omega-hydroxyalkanoate co-substrates', 2017, *AMB Express*, 7, doi:10.1186/s13568-017-0417-y.
- Nazaries, L., Karunaratne, S.B., Delgado-Baquerizo, M., Campbell, C.D., Singh, B.K., 'Environmental drivers of the geographical distribution of methanotrophs: Insights from a national survey', 2018, *Soil Biology and Biochemistry*, 127, doi:10.1016/j.soilbio.2018.08.014.

- Orphan, V.J., House, C.H., Hinrichs, K.-U., McKeegan, K.D., DeLong, E.F., 'Multiple archaeal groups mediate methane oxidation in anoxic cold seep sediments', 2002, *Proceedings of the National Academy of Sciences U.S.A.*, 99, doi:10.1073/pnas.072210299.
- Petersen, S.O., Ambus, P., 'Methane oxidation in pig and cattle slurry storages, and effects of surface crust moisture and methane availability', 2006, *Nutrient Cycling in Agroecosystems*, 74, doi:10.1007/s10705-005-3822-6.
- Poomisitiporn, K., Rangsunvigit, P., Kitiyanan, B., Kulprathipanja, S., 'Competitive adsorption of methane and carbon dioxide on different activated carbons', 2016, *Chemical Engineering Transactions*, 52, doi:10.3303/CET1652021.
- Rashidi, A.M., Lotfi, R., Nouralishahi, A., Khodagholi, M.A., Zare, M., Eslamipour, F., 'Nanoporous carbons as promising novel methane adsorbents for natural gas technology', 2011, *Journal of Natural Gas Chemistry*, 20, doi:10.1016/S1003-9953(10)60239-3.
- Reddy, K.R., Rai, R.K., Green, S.J., Chetri, J.K., 'Effect of pH on methane oxidation and community composition in landfill cover soil', 2020, *Journal of Environmental Engineering*, 146, doi:10.1061/(ASCE)EE.1943-7870.0001712.
- Ren, G., Ma, A., Zhang, Y., Deng, Y., Zheng, G., Zhuang, X., Zhuang, G., Fortin, D., 'Electron acceptors for anaerobic oxidation of methane drive microbial community structure and diversity in mud volcanoes', 2018, *Environmental Microbiology*, 20, doi:10.1111/1462-2920.14128.
- Rice, F.C., 'Second generation technology for the production of pipeline quality gas from sanitary landfills', 1980, paper presented at Society of Petroleum Engineers (SPE) Unconventional Gas Recovery Symposium (UGR), Pittsburgh, PA, 18-21 May 1980, pp. 225-227.
- Rissanen, A.J., Saarenheimo, J., Tirola, M., Peura, S., Aalto, S.L., Karvinen, A., Nykänen, H., 'Gammaproteobacterial methanotrophs dominate methanotrophy in aerobic and anaerobic layers of boreal lake waters', 2018, *Aquatic Microbial Ecology*, 81, doi:10.3354/ame01874.
- Ro, S.Y., Rosenzweig, A.C., 'Recent advances in the genetic manipulation of *Methylosinus trichosporium* OB3b', 2018, *Methods in Enzymology*, 605, doi:10.1016/bs.mie.2018.02.011.
- Rubin-Blum, M., Antony, C.P., Sayavedra, L., Martínez-Pérez, C., Birgel, D., Peckmann, J., Wu, Y.-C., Cardenas, P., MacDonald, I., Marcon, Y., Sahling, H., Hentschel, U., Dubilier, N., 'Fueled by methane: deep-sea sponges from asphalt seeps gain their nutrition from methane-oxidizing symbionts', 2019, *ISME Journal*, 13, doi:10.1038/s41396-019-0346-7.
- Sanchis-Perucho, P., Robles, Á., Durán, F., Ferrer, J., Seco, A., 'PDMS membranes for feasible recovery of dissolved methane from AnMBR effluents', 2020, *Journal of Membrane Science*, 604, doi:10.1016/j.memsci.2020.118070.
- Šantl-Temkiv, T., Finster, K., Hansen, B.M., Pašić, L., Karlson, U.G., 'Viable methanotrophic bacteria enriched from air and rain can oxidize methane at cloud-like conditions', 2013, *Aerobiologia*, 29, doi:10.1007/s10453-013-9287-1.
- Scheller, S., Yu, H., Chadwick, G.L., McGlynn, S.E., Orphan, V.J., 'Artificial electron acceptors decouple archaeal methane oxidation from sulfate reduction', 2016, *Science*, 351, doi:10.1126/science.aad7154.
- Semrau, J.D., Dispirito, A.A., Yoon, S., 'Methanotrophs and copper', 2010, *FEMS Microbiology Reviews*, 34, doi:10.1111/j.1574-6976.2010.00212.x.
- Serrano-Silva, N., Valenzuela-Encinas, C., Marsch, R., Dendooven, L., Alcántara-Hernández, R.J., 'Changes in methane oxidation activity and methanotrophic community composition in saline alkaline soils', 2014, *Extremophiles*, 18, doi:10.1007/s00792-014-0641-1.
- Shi, L.-D., Wang, M., Li, Z.-Y., Lai, C.-Y., Zhao, H.-P., 'Dissolved oxygen has no inhibition on methane oxidation coupled to selenate reduction in a membrane biofilm reactor', 2019, *Chemosphere*, 234, doi:10.1016/j.chemosphere.2019.06.138.
- Shi, Q., Wang, J., Shang, H., Bai, H., Zhao, Y., Yang, J., Dong, J., Li, J., 'Effective CH₄ enrichment from N₂ by SIM-1 via a strong adsorption potential SOD cage', 2020, *Separation and Purification Technology*, 230, doi:10.1016/j.seppur.2019.115850.

Skenneron, C.T., Chourey, K., Iyer, R., Hettich, R.L., Tyson, G.W., Orphan, V.J., 'Methane-fueled syntrophy through extracellular electron transfer: Uncovering the genomic traits conserved within diverse bacterial partners of anaerobic methanotrophic archaea', 2017, *mBio*, 8, doi:10.1128/mBio.00530-17.

Strong, P.J., Xie, S., Clarke, W.P., 'Methane as a resource: Can the methanotrophs add value?', 2015, *Environmental Science and Technology*, 49, doi:10.1021/es504242n.

Sun, F.-Y., Dong, W.-Y., Shao, M.-F., Lv, X.-M., Li, J., Peng, L.-Y., Wang, H.-J., 'Aerobic methane oxidation coupled to denitrification in a membrane biofilm reactor: Treatment performance and the effect of oxygen ventilation', 2013, *Bioresource Technology*, 145, doi:10.1016/j.biortech.2013.03.115.

Sun, Q., Li, H., Yan, J., Liu, L., Yu, Z., Yu, X., 'Selection of appropriate biogas upgrading technology-a review of biogas cleaning, upgrading and utilisation', 2015, *Renewable and Sustainable Energy Reviews*, 51, doi:10.1016/j.rser.2015.06.029.

Terent'ev, P.A., Gyl'maliev, A.M., Ruban, V.A., Malysenko, V.S., 'Coal adsorbents using for gaseous effluents purification', 1996, *Khimiya Tverdogo Topliva*, 1, pp. 59–66.

Timmers, P.H.A., Welte, C.U., Koehorst, J.J., Plugge, C.M., Jetten, M.S.M., Stams, A.J.M., 'Reverse methanogenesis and respiration in methanotrophic archaea', 2017, *Archaea*, 2017, doi:10.1155/2017/1654237.

Urbanová, Z., Pícek, T., Bárta, J., 'Effect of peat re-wetting on carbon and nutrient fluxes, greenhouse gas production and diversity of methanogenic archaeal community', 2011, *Ecological Engineering*, 37, doi:10.1016/j.ecoleng.2010.07.012.

Ursueguía, D., Díaz, E., Ordóñez, S., 'Adsorption of methane and nitrogen on basolite MOFs: Equilibrium and kinetic studies', 2020, *Microporous and Mesoporous Materials*, 298, doi:10.1016/j.micromeso.2020.110048.

Utaki, T., 'Development of coal mine methane concentration technology for reduction of greenhouse gas emissions', 2010, *Science China Technological Sciences*, 53, doi:10.1007/s11431-009-0419-3.

Vaksmas, A., Lüke, C., van Alen, T., Valè, G., Lupotto, E., Jetten, M.S.M., Ettwig, K.F., 'Distribution and activity of the anaerobic methanotrophic community in a nitrogen-fertilized Italian paddy soil', 2016, *FEMS Microbiology Ecology*, 92, doi:10.1093/femsec-fiw181.

Valenzuela, E.I., Prieto-Davó, A., López-Lozano, N.E., Hernández-Eligio, A., Vega-Alvarado, L., Juárez, K., García-González, A.S., López, M.G., Cervantes, F.J., 'Anaerobic methane oxidation driven by microbial reduction of natural organic matter in a tropical wetland', 2017, *Applied and Environmental Microbiology*, 83, doi:10.1128/AEM.00645-17.

van Grinsven, S., Sinninghe Damsté, J.S., Abdala Asbun, A., Engelmann, J.C., Harrison, J., Villanueva, L., 'Methane oxidation in anoxic lake water stimulated by nitrate and sulfate addition', 2020, *Environmental Microbiology*, 22, doi:10.1111/1462-2920.14886.

Ve, T., Mathisen, K., Helland, R., Karlsen, O.A., Fjellbirkeland, A., Røhr, Å.K., Andersson, K.K., Pedersen, R.-B., Lillehaug, J.R., Jensen, H.B., 'The *Methylococcus capsulatus*, bath, secreted protein, MopE*, binds both reduced and oxidized copper', 2012, *PLOS ONE*, 7, doi:10.1371/journal.pone.0043146.

Vignali, G., Vitale G., 'Life cycle assessment of bio-methane obtained by valorization of food waste', 2017, *Proceedings of the XXII Summer School 'Francesco Turco'*, Palermo, Italy, 13–15 September, pp. 136–142.

Vigneron, A., Alsop, E.B., Cruaud, P., Philibert, G., King, B., Baksmaty, L., Lavalley, D., Lomans, B.P., Elie-Fadrosh, E., Kypides, N.C., Head, I.M., Tsismetis, N., 'Contrasting pathways for anaerobic methane oxidation in Gulf of Mexico cold seep sediments', 2019, *mSystems*, 4, doi:10.1128/mSystems.00091-18.

Wentworth, R.L., Ashare, E., Wise, D.L., 'Fuel gas production from animal residue part I. Technical perspective', 1979, *Resource Recovery and Conservation*, 3, doi:10.1016/0304-3967(79)90021-0.

Winkel, M., Mitzscherling, J., Overduin, P.P., Horn, F., Winterfeld, M., Rijkers, R., Grigoriev, M.N., Knoblauch, C., Mangelsdorf, K., Wagner, D., Liebner, S., 'Anaerobic methanotrophic communities thrive in deep submarine permafrost', 2018, *Scientific Reports*, 8, doi:10.1038/s41598-018-19505-9.

Wise, D.L., Kispert, R.G., Langton, E.W., 'A review of bioconversion systems for energy recovery from municipal solid waste part II: Fuel gas production', 1981, *Resources and Conservation*, 6, doi:10.1016/0166-3097(81)90063-8.

Wrede, C., Kokoschka, S., Dreier, A., Heller, C., Reitner, J., Hoppert, M., 'Deposition of biogenic iron minerals in a methane oxidizing microbial mat', 2013a, *Archaea*, 2013, doi:10.1155/2013/102972.

- Wrede, C., Krukenberg, V., Dreier, A., Reitner, J., Heller, C., Hoppert, M., 'Detection of metabolic key enzymes of methane turnover processes in cold seep microbial biofilms', 2013b, *Geomicrobiology Journal*, 30, doi:10.1080/01490451.2012.665150.
- Xie, T., Yang, Q., Winkler, M.K.H., Wang, D., Zhong, Y., An, H., Chen, F., Yao, F., Wang, X., Wu, J., Li, X., 'Perchlorate bioreduction linked to methane oxidation in a membrane biofilm reactor: Performance and microbial community structure', 2018, *Journal of Hazardous Materials*, 357, doi:10.1016/j.jhazmat.2018.06.011.
- Yang, Y., Wu, Y., Liu, H., Ribeiro, A.M., Li, P., Yu, J., Rodrigues, A.E., 'Enrichment of ventilation air methane by adsorption with displacement chromatography technology: Experiment and numerical simulation', 2016, *Chemical Engineering Science*, 149, doi:10.1016/j.ces.2016.04.030.
- Yang, X., Liu, Y., Li, Z., Zhang, C., Xing, Y., 'Vacuum exhaust process in pilot-scale vacuum pressure swing adsorption for coal mine ventilation air methane enrichment', 2018, *Energies*, 11, doi:10.3390/en11051030.
- Yoon, S., Carey, J.N., Semrau, J.D., 'Feasibility of atmospheric methane removal using methanotrophic biotrickling filters', 2009, *Applied Microbiology and Biotechnology*, 83, doi:10.1007/s00253-009-1977-9.
- Zang, X., Liang, D., 'Experimental study on hydrate formation with synthesized CH₄/CO₂/N₂ ternary mixtures', 2018, *Energy Fuels*, 32, doi:10.1021/acs.energyfuels.8b02021.
- Zhai, J., Li, Y., He, M., Ma, H., Dai, Y., 'Review on the research progress of archaeal anaerobic methanotrophs in freshwater system', 2019, *Chinese Journal of Environmental Engineering*, 13, doi:10.12030/j.cjee.201812098.
- Zhang, X., Xia, J., Pu, J., Cai, C., Tyson, G.W., Yuan, Z., Hu, S., 'Biochar-mediated anaerobic oxidation of methane', 2019, *Environmental Science and Technology*, 53, doi:10.1021/acs.est.9b01345.
- Zhu, N., An, P., Krishnakumar, B., Zhao, L., Sun, L., Mizuochi, M., Inamori, Y., 'Effect of plant harvest on methane emission from two constructed wetlands designed for the treatment of wastewater', 2007, *Journal of Environmental Management*, 85, doi:10.1016/j.jenvman.2006.11.004.

List of abbreviations and definitions

| | |
|------------------|--|
| ANME | anaerobic methanotroph (anaerobic methane-oxidizing archaea) |
| AOM | anaerobic oxidation of methane |
| BTF | biotrickling filter |
| CH ₄ | methane |
| CO ₂ | carbon dioxide |
| GHG | greenhouse gas |
| GWP | global warming potential |
| IPCC | Intergovernmental Panel on Climate Change |
| JRC | Joint Research Centre |
| MOF | metal–organic framework |
| N ₂ O | nitrous oxide |
| NGS | next-generation sequencing |

List of figures

| | |
|---|----|
| Figure 1. Illustration of CH ₄ oxidation (vertical air-water cross-section. The blue line represents the air-water interface). Anaerobic CH ₄ oxidation is visualised by the red and green consortium, representing ANME and sulfate-reducing bacteria, respectively. | 7 |
| Figure 2. JRC outputs related to CH ₄ per JRC directorate and output type | 26 |
| Figure 3. JRC outputs related to CH ₄ per unit in JRC Directorates C and D | 26 |

Annexes

Annex 1: Literature mapping – Clustering of publications

This review analysed 109 scientific communications, mapped along various dimensions:

- by type of document: 17 documents reviewed specific aspects of CH₄;
- by environment (e.g. atmosphere, soil, freshwater and saline water) and by source of methane held in these environments;
- by process: methane production through methanogenesis, methane absorption for separation or for storage, genetic approaches, as well as methanotrophy, methane oxidation and methane conversion;
-
- by by- and co-products, such as carbonates, chromate, copper, iron, manganese, nitrous compounds (N₂O, NO_x), ammonia, nitrate, sulfur and sulfate, but also fuels and chemicals, such as lactic acid, methanol and acetate;
- by technologies relying on living organisms and microorganisms, such as biochar, biotrickling filtration, digesters, microbial fuel cells and relevant genetic technologies (e.g. NGS), or relying on ((in)organic) materials such as membranes, sorbents or pressure swing absorption;
- by stage of developments: demonstration or commercial projects, patents, estimates of the potential for CH₄ emission reduction, quantified costs, other economic aspects, programmes and partnerships.

Types of documents

Few references review specific aspects of CH₄ production, concentration and consumption (Alonso et al., 2017 ; Bennett et al., 2018; Bhattarai et al., 2019; Choi et al., 2016; Garcia and Cervik, 1988; Gutknecht et al., 2006; He et al., 2018; Jiao et al., 2012; Kapoor et al., 2019; Knief, 2015; Lieven et al., 2018; Liu et al., 2020a; Strong et al., 2015; Sun et al., 2015; Timmers et al., 2017; Wentworth et al., 1979; Wise et al., 1981). These references are highlighted in bold in the rest of this chapter.

Environments and sources of methane

Atmosphere (Delgado et al., 2018; Kolb and Horn, 2012; Menyailo et al., 2008; Šantl-Temkiv et al., 2013; Yoon et al., 2009).

Soil:

- coal (Bracmort et al., 2014; Cramer et al., 2009; **Garcia and Cervik, 1988**; Industry news, 2004; Kosmack et al., 2008; Leisle and Kovalski, 2018; **Liu et al., 2020a**; Moiseev et al., 2016a,b; Poomisitiporn et al., 2016; Shi et al., 2020; Ternet'ev et al., 1996; Utaki, 2010; Yang et al., 2016, 2018);
- farms (animals, digestion, manure) (Bracmort et al., 2014; Chianese et al., 2015; Petersen and Ambus, 2006; **Wentworth et al., 1979**; Yoon et al., 2009);
- forest and grassland (Menyailo et al., 2008);
- hydrates (**Jiao et al., 2012**; Lin et al., 2014; Moiseev et al., 2016b; Zang and Liang, 2018);
- landfills (Air Liquide targets landfill gas-to-energy, 2007; Bracmort et al., 2014; Cavenati et al., 2005; Industry news, 2004; Mei et al., 2019; Reddy et al., 2020; Rice, 1980; **Wise et al., 1981**; Yoon et al., 2009);
- mud volcanoes (Bhattarai et al., 2018; Ren et al., 2018);
- other fossil fuels (oil and gas production, natural and shale gas, gob gas) (Ajayi et al., 2015; **Bennett et al., 2018**; Bracmort et al., 2014; Fei et al., 2020; **Garcia and Cervik, 1988**; Hur et al., 2017; Industry news, 2004; Moiseev et al., 2016b; Poomisitiporn et al., 2016; Shi et al., 2020);
- paddy fields (Li et al., 2020; Vaksmaa et al., 2016);
- peatlands (Urbanová et al., 2011);
- solid waste (Vignali and Vitale, 2017; Reddy et al., 2020; **Wise et al., 1981**);

- Undefined soil (**Gutknecht et al., 2006**; Jiang et al., 2016; Kolb and Horn, 2012; Kwon et al., 2017; Li et al., 2020; Menyailo et al., 2008; Nazaries et al., 2018; Reddy et al., 2020; Šantl-Temkiv et al., 2013; Serrano-Silva et al., 2014; Urbanová et al., 2011; Vaksmaa et al., 2016; Zhang et al., 2019);
- wetlands (**Gutknecht et al., 2006**; Kolb and Horn, 2012; Kwon et al., 2017; Liu et al., 2020b; Valenzuela et al., 2017; Zhu et al., 2007).

Freshwaters:

- dams (Lima et al., 2008);
- lakes (Bar-Or et al., 2017; Bhattarai et al., 2017; Cassarini et al., 2019c; Martinez-Cruz et al., 2017; Rissanen et al., 2018; van Grinsven et al., 2020);
- wastewater (Cashman et al., 2018; Cassarini et al., 2019a; Henares et al., 2016; Kampman et al., 2014; Li et al., 2020; Liu et al., 2020b; Modin et al., 2008; Sanchis-Perucho et al., 2020; **Strong et al., 2015**);
- water hyacinths (Annachatre and Khanna, 1987).

Saline waters:

- oceans (Chen et al., 2014; Hatzenpichler et al., 2016; **Jiao et al., 2012**; Semrau et al., 2010; Skennerton et al., 2017; Zhai et al., 2019);
- marine environments (Aromokeye et al., 2020; Bhattarai et al., 2017; **Bhattarai et al., 2019**; Bomberg et al., 2015; Cassarini et al., 2019c; Chen et al., 2014; Cruaud et al., 2014, 2015; **Jiao et al., 2012**; Lin et al., 2014; Orphan et al., 2002; Rubin-Blum et al., 2019; Scheller et al., 2016; Vigneron et al., 2019; Winkel et al., 2018);
- saline lakes (Serrano-Silva et al., 2014);
- seas (Aromokeye et al., 2020; Bhattarai et al., 2018; **Bhattarai et al., 2019**; Cassarini et al., 2019a,b; Chen et al., 2014; Cruaud et al., 2014, 2015; **Jiao et al., 2012**; Rubin-Blum et al., 2019; Scheller et al., 2016; Vigneron et al., 2019; Wrede et al., 2013a,b).

Processes

CH₄ production (methanogenesis) (Bhattarai et al., 2017; Cruaud et al., 2014; **Gutknecht et al., 2006**; Kwon et al., 2017; Li et al., 2020; Liu et al., 2020b; Martinez-Cruz et al., 2017; Urbanová et al., 2011; Zhu et al., 2007).

Genetic approaches (Bhattarai et al., 2017; Bomberg et al., 2015; Cassarini et al., 2019b; Cruaud et al., 2014, 2015; Hatzenpichler et al., 2016; **Knief, 2015**; Kolb and Horn, 2012; Li et al., 2020; Lin et al., 2014; Luo et al., 2019; Lv et al., 2019; Mei et al., 2019; Nazaries et al., 2018; Orphan et al., 2002; Reddy et al., 2020; Ren et al., 2018; Rissanen et al., 2018; Ro and Rosenzweig, 2018; Rubin-Blum et al., 2019; Šantl-Temkiv et al., 2013; Semrau et al., 2010; Serrano-Silva et al., 2014; Shi et al., 2019; Skennerton et al., 2017; Vaksmaa et al., 2016; van Grinsven et al., 2020; Vigneron et al., 2019; Winkel et al., 2018; Xie et al., 2018; Zhang et al., 2019).

Absorption (Abbott et al., 2011; **Alonso et al., 2017**; Cavenati et al., 2005; **Choi et al., 2016**; Delgado et al., 2018; **Kapoor et al., 2019**; Kostoglou et al., 2017; **Liu et al., 2020a**; Mahmoudian et al., 2016; Moiseev et al., 2016b; Poomisitiporn et al., 2016; Rashidi et al., 2011; Shi et al., 2020; Ternet'ev et al., 1996; Ursueguía et al., 2020; Utaki, 2010; Vignali and Vitale, 2017; Yang et al., 2018; Yang et al., 2016):

- for separation (Abbott et al., 2011; **Alonso et al., 2017**; Cavenati et al., 2005; **Kapoor et al., 2019**; Kostoglou et al., 2017; Moiseev et al., 2016b; Shi et al., 2020; Ternet'ev et al., 1996; Vignali and Vitale, 2017; Yang et al., 2016, 2018);
- for storage (Abbott et al., 2011; **Alonso et al., 2017**; **Choi et al., 2016**; Kosmack et al., 2008; Kostoglou et al., 2017; Mahmoudian et al., 2016; Poomisitiporn et al., 2016; Rashidi et al., 2011).

Separation processes not explicitly based on adsorption (Air Liquide targets landfill gas-to-energy, 2007; Ajayi et al., 2015; Annachatre and Khanna, 1987; Cramer et al., 2009; Fei et al., 2020; **Garcia and Cervik, 1988**; Leisle and Kovalski, 2018; Moiseev et al., 2016a; Rice, 1980; **Sun et al., 2015**; Zang and Liang, 2018).

CH₄ consumption is mainly carried out by oxidation, but also by conversion:

- conversion (**Bennett et al., 2018**; Fei et al., 2020; Hur et al., 2017; Ito et al., 2018; McAnulty et al., 2017; **Strong et al., 2015**; Wrede et al., 2013b);

- methanotrophy (Aromokeye et al., 2020; Bar-Or et al., 2017; **Bennett et al., 2018**; Bhattarai et al., 2017, 2018; **Bhattarai et al., 2019**; Bomberg et al., 2015; Casey et al., 2004; Cassarini et al., 2019a,b,c; Chen et al., 2014; Cruaud et al., 2014, 2015; Fei et al., 2020; **Gutknecht et al., 2006**; Hatzenpichler et al., 2016; **He et al., 2018**; Hur et al., 2017; Ito et al., 2018; Jiang et al., 2016; **Jiao et al., 2012**; Kalyuzhnaya et al., 2018; Kampman et al., 2014; **Knief, 2015**; Kolb and Horn, 2012; Kwon et al., 2017; Li et al., 2020; **Lieven et al., 2018**; Lin et al., 2014; Liu et al., 2020b; Luo et al., 2019; Lv et al., 2019; Martinez-Cruz et al., 2017; McNulty et al., 2017; Mei et al., 2019; Menyailo et al., 2008; Modin et al., 2008; Myung et al., 2017; Nazaries et al., 2018; Orphan et al., 2002; Petersen and Ambus, 2006; Reddy et al., 2020; Ren et al., 2018; Rissanen et al., 2018; Ro and Rosenzweig, 2018; Rubin-Blum et al., 2019; Šantl-Temkiv et al., 2013; Scheller et al., 2016; Semrau et al., 2010; Serrano-Silva et al., 2014; Shi et al., 2019; Skennerton et al., 2017; **Strong et al., 2015**; Sun et al., 2013; **Timmers et al., 2017**; Urbanová et al., 2011; Vaksmaa et al., 2016; Valenzuela et al., 2017; van Grinsven et al., 2020; Ve et al., 2012; Vigneron et al., 2019; Winkel et al., 2018; Wrede et al., 2013a,b; Xie et al., 2018; Yoon et al., 2009; Zhai et al., 2019; Zhang et al., 2019; Zhu et al., 2007);
- methylotrophy (**Bennett et al., 2018**; Jiang et al., 2016; **Lieven et al., 2018**; Martinez-Cruz et al., 2017; Reddy et al., 2020; Rubin-Blum et al., 2019; Wrede et al., 2013b; Xie et al., 2018);
- other oxidation, not explicitly linked to reduction (**Bhattarai et al., 2019**; Casey et al., 2004; Jiang et al., 2016; Kalyuzhnaya et al., 2018; Kosmack et al., 2008; **Knief, 2015**; Kwon et al., 2017; Mei et al., 2019; Menyailo et al., 2008; Orphan et al., 2002; Petersen and Ambus, 2006; Reddy et al., 2020; Šantl-Temkiv et al., 2013; Semrau et al., 2010; Sun et al., 2013; Ve et al., 2012) and including:
 - hydroxylation (Ito et al., 2018);
 - photooxidation (Šantl-Temkiv et al., 2013);
- redox (oxidation and reduction) (Aromokeye et al., 2020; Bar-Or et al., 2017; Bhattarai et al., 2017, 2018; Cassarini et al., 2019a,b,c; Chen et al., 2014; Hatzenpichler et al., 2016; **He et al., 2018**; Liu et al., 2020b; Luo et al., 2019; Lv et al., 2019; Martinez-Cruz et al., 2017; Modin et al., 2008; Ren et al., 2018; Rissanen et al., 2018; Scheller et al., 2016; Serrano-Silva et al., 2014; Shi et al., 2019; Skennerton et al., 2017; **Timmers et al., 2017**; Vaksmaa et al., 2016; Valenzuela et al., 2017; van Grinsven et al., 2020; Vigneron et al., 2019; Winkel et al., 2018; Wrede et al., 2013a,b; Xie et al., 2018; Zhai et al., 2019; Zhang et al., 2019);
- reduction, not explicitly linked to oxidation (Cruaud et al., 2015; **Gutknecht et al., 2006**; Kolb and Horn, 2012; McNulty et al., 2017);
- reverse methanogenesis (**Lieven et al., 2018**; Lv et al., 2019; **Timmers et al., 2017**; Wrede et al., 2013b).

Both conversion and reduction are linked to by- and co-products.

By- and co-products:

- carbonates (Chen et al., 2014; Wrede et al., 2013a,b);
- chromate (Lv et al., 2019);
- copper (Kampman et al., 2014; Ro and Rosenzweig, 2018; Semrau et al., 2010; Ve et al., 2012);
- iron (Aromokeye et al., 2020; Bar-Or et al., 2017; **Gutknecht et al., 2006**; Martinez-Cruz et al., 2017; Ren et al., 2018; Wrede et al., 2013a);
- manganese (**He et al., 2018**; Liu et al., 2020b; Luo et al., 2019; Rissanen et al., 2018; Winkel et al., 2018);
- nitrous compounds (N₂O, NO_x, nitrous, ammonia, nitrate, nitrous, (de)nitrifi(cation/er)) (**He et al., 2018**; Kampman et al., 2014; Kolb and Horn, 2012; Li et al., 2020; Lin et al., 2014; Luo et al., 2019; McNulty et al., 2017; Mei et al., 2019; Menyailo et al., 2008; Modin et al., 2008; Ren et al., 2018; Rissanen et al., 2018; Shi et al., 2019; **Strong et al., 2015**; Sun et al., 2013; **Timmers et al., 2017**; Urbanová et al., 2011; Vaksmaa et al., 2016; van Grinsven et al., 2020; Winkel et al., 2018; Xie et al., 2018);
- sulfur/sulphur/sulfate (Aromokeye et al., 2020; Bar-Or et al., 2017; Bhattarai et al., 2017, 2018; **Bhattarai et al., 2019**; Bomberg et al., 2015; Cassarini et al., 2019a,b,c; Chen et al., 2014; Cruaud et al., 2015; **Gutknecht et al., 2006**; Hatzenpichler et al., 2016; **He et al., 2018**; Lin et al., 2014; Luo et al., 2019; McNulty et al., 2017; Mei et al., 2019; Ren et al., 2018; Scheller et al., 2016; Skennerton et al., 2017; **Timmers et al., 2017**; Valenzuela et al., 2017; van Grinsven et al., 2020; Vigneron et al., 2019; Winkel et al., 2018; Wrede et al., 2013a,b).

Further, the conversion process indicated above leads to the transformation of CH₄ in fuels and chemicals (**Bennett et al., 2018**), lactic acid (Fei et al., 2020), methanol (Hur et al., 2017; Ito et al., 2018; Wrede et al., 2013b) and acetate (McAnulty et al., 2017), and has other numerous applications (**Strong et al., 2015**).

Technologies

CH₄ technologies can be divided into two categories: those based on living organisms and those of a mineral nature.

Numerous living organisms support the processes of methanotrophy, methylotrophy and (reverse) methanogenesis, including Euryarchaeota (phylum), *Candidatus Methanoperedens* (genus), *Candidatus Methanoperedens nitroreducens* (species), *Methanosarcina mazei* (species), *Candidatus Bathyarchaeota* (phylum), Thaumarchaeota (phylum), *Moheibacter* (genus), Proteobacteria (phylum), *Hyphomicrobium* (genus), *Methylocystis* (genus), *Methylosinus* (genus), *Methylosinus trichosporium* OB3b (species), *Cupriavidus* (genus), *Variovorax* (genus), *Methylophilus* (genus), *Methylobacter* (genus), *Methylocaldum* (genus), *Methylococcus capsulatus* (species), *Methylomicrobium alcaliphilum* (species), *Methylomonas* (genus), *Candidatus Methyloiumidiphilus alinensis* (species), Verrucomicrobia (class), *Candidatus Methyloimabilis oxyfera* (species)⁽¹⁰⁾ (Aromokeye et al., 2020; Bar-Or et al., 2017; Bhattarai et al., 2017; Bomberg et al., 2015; Cassarini et al., 2019a,b; Hatzenpichler et al., 2016; Jiang et al., 2016; **Jiao et al., 2012**; Kalyuzhnaya et al., 2018; Kampman et al., 2014; Knief, 2015; Kolb and Horn, 2012; Liu et al., 2020b; Luo et al., 2019; Lv et al., 2019; Martinez-Cruz et al., 2017; McAnulty et al., 2017; Mei et al., 2019; Orphan et al., 2002; Reddy et al., 2020; Rissanen et al., 2018; Ro and Rosenzweig, 2018; Rubin-Blum et al., 2019; Šantl-Temkiv et al., 2013; Serrano-Silva et al., 2014; Shi et al., 2019; **Timmers et al., 2017**; Vaksmaa et al., 2016; van Grinsven et al., 2020; Ve et al., 2012; Vigneron et al., 2019; Winkel et al., 2018; Zhai et al., 2019; Zhang et al., 2019; Zhu et al., 2007).

Technologies relying on these **micro-organisms** are mainly related to biomethane production and CH₄ consumption:

- **biochar** (Reddy et al., 2020; Zhang et al., 2019);
- **biotrickling filtration** (Cassarini et al., 2019a,b; Yoon et al., 2009);
- **digesters** (Bracmort et al., 2014; Chianese et al., 2015; Vignali and Vitale, 2017; **Wentworth et al., 1979**; **Wise et al., 1981**; Xie et al., 2018);
- **microbial fuel cells** (McAnulty et al., 2017).

For separation/adsorption processes, **mineral-based technologies** predominate:

Membranes (Air Liquide targets landfill gas-to-energy, 2007; Ajayi et al., 2015; Bhattarai et al., 2018; Casey et al., 2004; Cashman et al., 2018; Cramer et al., 2009; **Garcia and Cervik, 1988**; Henares et al., 2016; Ito et al., 2018; Kampman et al., 2014; **Kapoor et al., 2019**; Luo et al., 2019; Lv et al., 2019; Modin et al., 2008; Moiseev et al., 2016b; Sanchis-Perucho et al., 2020; Shi et al., 2019; Sun et al., 2013; Vignali and Vitale, 2017; Xie et al., 2018):

- degasing membranes (Cramer et al., 2009; Henares et al., 2016; Moiseev et al., 2016b; Sanchis-Perucho et al., 2020);
- membrane-based reactors (Bhattarai et al., 2018; Casey et al., 2004; Cashman et al., 2018; Kampman et al., 2014; Luo et al., 2019; Lv et al., 2019; Modin et al., 2008; Shi et al., 2019; Sun et al., 2013; Xie et al., 2018);
- zeolite membranes (Ajayi et al., 2015).

Zeolites are nanomaterials that can be used as membranes for gas separation. Such **nanomaterials** are also being used for CH₄ adsorption. More generally **nanomaterials** also being used for CH₄ adsorption are divided into non-carbonaceous materials (e.g. zeolites, metal-organic frameworks and porous polymers) and carbonaceous materials (e.g. activated carbons, ordered porous carbons and activated carbon fibres) (Choi et al., 2016):

- biochar (Delgado et al., 2018);
- graphene (Alonso et al., 2017; Mahmoudian et al., 2016);

⁽¹⁰⁾ See NCBI, 'Taxonomy', n.d., available at: <https://www.ncbi.nlm.nih.gov/taxonomy>, accessed 20.10.2020.

- other carbonaceous materials (including activated carbon beads, ultra-microporous activated carbon cloth, coconut shell activated carbon, palm shell activated carbon and coal-based activated carbon) (Bar-Or et al., 2017; **Bennett et al., 2018**; Bhattarai et al., 2018; **Choi et al., 2016**; Kostoglou et al., 2017; Poomisitiporn et al., 2016; Rashidi et al., 2011; Ternet'ev et al., 1996; Yang et al., 2016, 2018);
- metal-organic frameworks (MOF) (Alonso et al., 2017; **Choi et al., 2016**; Delgado et al., 2018; Shi et al., 2020; Ursueguía et al., 2020);
- polymers (Abbott et al., 2011; **Choi et al., 2016**; Kostoglou et al., 2017; Cruaud et al., 2015; **Strong et al., 2015**);
- zeolites (Ajayi et al., 2015; **Choi et al., 2016**; Delgado et al., 2018).

Sorbents also form the basis of the separation process in **pressure swing adsorption** technologies (Cavenati et al., 2005; Utaki, 2010; Yang et al., 2018).


Stages of development

- Demonstration or commercial projects (Bhattarai et al., 2017; Fei et al., 2020; **He et al., 2018**; Hur et al., 2017; Kosmack et al., 2008; Liu et al., 2020b; Rice, 1980; Utaki, 2010; Yang et al., 2016).
- Patents (Air Liquide targets landfill gas-to-energy, 2007).
- Estimates of the potential for CH₄ emission reduction (Annachhatre and Khanna, 1987; Aromokeye et al., 2020; Hur et al., 2017; Industry news, 2004; Kosmack et al., 2008; Li et al., 2020; Lima et al., 2008; Liu et al., 2020b; Mei et al., 2019; Sanchis-Perucho et al., 2020; Serrano-Silva et al., 2014; Shi et al., 2020; Utaki, 2010; Vaksmaa et al., 2016; Valenzuela et al., 2017; Winkel et al., 2018; Yang et al., 2016; Yoon et al., 2009; Zhang et al., 2019).
- Quantified costs (Fei et al., 2020; Yoon et al., 2009).
- Other economic aspects (Bracmort et al., 2014; Chianese et al., 2015; **Choi et al., 2016**; Fei et al., 2020; **Garcia and Cervik, 1988**; **Kapoor et al., 2019**; Kosmack et al., 2008; **Lieven et al., 2018**; Mahmoudian et al., 2016; Rice, 1980; Sanchis-Perucho et al., 2020; **Wentworth et al., 1979**; Yoon et al., 2009).
- Programmes (Bracmort et al., 2014; Chianese et al., 2015).
- Partnerships (Industry news, 2004).

Annex 2: JRC activities on methane


The JRC has registered 94 outputs related to CH₄ since its reorganisation in mid-2016. Figure 2 shows the JRC directorates leading these deliverables and the count of outputs produced per type of deliverable, while Figure 3 shows the JRC units leading these deliverables within Directorates C (Energy, transport and climate) and D (Sustainable resources), which lead 74 and 18 deliverables, respectively.

Figure 2. JRC outputs related to CH₄ per JRC directorate and output type



Source: JRC, 2020.

Figure 3. JRC outputs related to CH₄ per unit in JRC Directorates C and D



Source: JRC, 2020.

The analysis of these reports shows a dichotomy within the JRC, with CH₄ considered as either a primary energy source and carrier or a GHG, linked to the missions of the respective units/directorates. These different views are expected to converge with decarbonisation of the energy sector, policy integration aimed at breaking silos and upcoming JRC projects.

GETTING IN TOUCH WITH THE EU

In person

All over the European Union there are hundreds of Europe Direct information centres. You can find the address of the centre nearest you at: https://europa.eu/european-union/contact_en

On the phone or by email

Europe Direct is a service that answers your questions about the European Union. You can contact this service:

- by freephone: 00 800 6 7 8 9 10 11 (certain operators may charge for these calls),
- at the following standard number: +32 22999696, or
- by electronic mail via: https://europa.eu/european-union/contact_en

FINDING INFORMATION ABOUT THE EU

Online

Information about the European Union in all the official languages of the EU is available on the Europa website at: https://europa.eu/european-union/index_en

EU publications

You can download or order free and priced EU publications from EU Bookshop at: <https://publications.europa.eu/en/publications>. Multiple copies of free publications may be obtained by contacting Europe Direct or your local information centre (see https://europa.eu/european-union/contact_en).

The European Commission's science and knowledge service

Joint Research Centre

JRC Mission

As the science and knowledge service of the European Commission, the Joint Research Centre's mission is to support EU policies with independent evidence throughout the whole policy cycle.



EU Science Hub
ec.europa.eu/jrc



@EU_ScienceHub



EU Science Hub - Joint Research Centre



EU Science, Research and Innovation



EU Science Hub



Publications Office
of the European Union

doi:10.2760/09370

ISBN 978-92-76-22969-8