



ÉCOLE POLYTECHNIQUE FÉDÉRALE DE LAUSANNE

Analyse III - Lebesgue

MATH-208

Note de

Anthony Gabino

Professeur : Krieger Joachim

Dernière mise à jour 7 septembre 2026
Bachelor Mathématique
Lausanne

Table des matières

Cours	5
1 Champs vectoriels	7
1.1 Définition de base	7
1.2 Intégrale le long d'un chemin	7
2 Measure theory	9
2.1 Motivation	9
2.2 Outer-measure	9

TABLE DES MATIÈRES

Cours

Cours n° 1– July 27th	9
---------------------------------	---

TABLE DES MATIÈRES

CHAPITRE 1

Champs vectoriels

1.1 Définition de base

Définition 1.1 (Domaine). Soit $U \subset \mathbb{R}^n$ un sous-ensemble, alors on dit que U est **domaine** s'il est ouvert et connexe.

Définition 1.2 (Champ scalaire). Un **champ scalaire** est une fonction $f : U \rightarrow \mathbb{R}$ de régularité C^k sur U .

Définition 1.3 (Champ vectoriel). Un **champ vectoriel** est une fonction $\mathbf{f} : U \rightarrow \mathbb{R}^n$ de régularité C^k sur U .

Définition 1.4 (chemin). Un **chemin** est une application continue

$$\gamma : I \rightarrow U$$

où $I = [s, t]$, $s < t \in \mathbb{R}$, et tel qu'il existe une partition de $I = \bigcup_{j=1}^N I_j$ où les I_j sont des intervalles fermés et d'intérieurs disjoints, tel que les restrictions

$$\gamma|_{I_j} \in C^1(I_j, U)$$

autrement dit γ est **continue par morceaux**.

1.2 Intégrale le long d'un chemin

Définition 1.5 (Intégrale curviligne). Soit $f \in C^0(U, \mathbb{R})$ un champ scalaire, et soit $\gamma : I \rightarrow U$ un chemin, alors on définit l'**intégrale curviligne** de f le long de γ par

$$\int_{\gamma} f \, ds = \int_I f(\gamma(t)) \cdot \|\gamma'(t)\| \, dt =: \sum_{j=1}^N \int_{I_j} f|_{I_j}(\gamma|_{I_j}(t)) \cdot \|\gamma'_{I_j}(t)\| \, dt$$

où

$$\|\gamma'(t)\| = \left(\sum_{i=1}^n \gamma'_i(t)^2 \right)^{\frac{1}{2}}$$

Définition 1.6 (Reparamétrisation d'un chemin). Soit $\gamma : [s, t] \rightarrow U$ un chemin et soit

$$\varphi : [\tilde{s}, \tilde{t}] \rightarrow [s, t]$$

un difféomorphisme de classe C^1 tel que $\varphi' > 0$. Alors la composition $\gamma \circ \varphi : [\tilde{s}, \tilde{t}] \rightarrow U$ est encore un chemin.

Lemme 1.1. Soit $U \subset \mathbb{R}^n$ un domaine, $f \in C^0(U)$ un champ scalaire, $\gamma : I \rightarrow U$ et $\gamma \circ \varphi : \tilde{I} \rightarrow U$ une

reparamétrisation. Alors

$$\int_{\gamma \circ \varphi} f \, ds = \int_{\gamma} f \, ds$$

DÉMONSTRATION :

Soit $\tilde{I} = [\tilde{s}, \tilde{t}]$, $I = [s, t]$. Alors

$$\int_{\gamma \circ \varphi} f \, ds = \int_{\tilde{I}} f(\gamma \circ \varphi) \cdot \|(\gamma \circ \varphi)'(t)\| \, dt = \int_{\tilde{s}}^{\tilde{t}} f(\gamma \circ \varphi(t)) \cdot \|\gamma' \circ \varphi(t)\varphi'(t)\| \, dt = \int_{\tilde{s}}^{\tilde{t}} f(\gamma \circ \varphi(t))\varphi'(t) \cdot \|\gamma' \circ \varphi(t)\| \, dt$$

et donc avec un changement de variable on trouve

$$\int_{\gamma \circ \varphi} f \, ds = \int_{\varphi(\tilde{s})}^{\varphi(\tilde{t})} f(\gamma(x)) \cdot \|\gamma'(x)\| \, dx = \int_s^t f(\gamma(x)) \cdot \|\gamma'(x)\| \, dx = \int_{\gamma} f \, ds$$

□

Measure theory

2.1 Motivation

Studying integration in several variables, the general question we wish to answer is this :

Given some subset $\Omega \subset \mathbb{R}^n$, and some real-valued function $f : \Omega \rightarrow \mathbb{R}$, is it possible to integrate f on Ω and therefore define $\int_{\Omega} f$?

In the case $n = 1$, we have already developed the notion of *Riemann integral*, which answers the above question when Ω is an interval $\Omega = [a, b] \subset \mathbb{R}$, and f is Riemann integrable. However, the class of Riemann integral functions is rather unsatisfactorily small and the extension of this notion to higher dimensions is possible but requires quite a bit of effort. For such reasons, we must look beyond the Riemann integral and introduce the notion of the *Lebesgue integral*.

Before we turn to the details, we begin with an informal discussion. In order to understand how to compute an integral $\int_{\Omega} f$, we must first understand a more basic and fundamental question :

How does one compute the length, area or volume of a subset $E \subset \mathbb{R}^n$?

This question is connected to that integration, because if one integrates the function 1 on the set E , then one should obtain, the length, the area or the volume of E depending on the dimension. To avoid splitting into cases depending on the dimension, we shall refer to the **measure** of E as either.

Ideally, to every subset $E \subset \mathbb{R}^n$ we would like to associate a non-negative number $m(E)$, which will be the measure of E such that,

1. $m(\emptyset) = 0$.
2. $m(E) \in [0, \infty]$.
3. $m(E) \leq m(F)$ whenever $E \subseteq F$.
4. $m(E \cup F) = m(E) + m(F)$ if E and F are disjoint.
5. $m(x + E) = m(E)$ for any $x \in \mathbb{R}^n$

Remarkably, it turns out that such a measure does not exist ; one cannot assign a non-negative number to every subset of $\mathbb{R}^n$ which has the above properties. However, we can rescue the situation by only measuring a certain class of sets in $\mathbb{R}^n$, that we will define **measurable set**.

2.2 Outer-measure

Before we construct Lebesgue measure, we first discuss a somewhat naive approach to finding the measure of a set - namely, we try to cover the set by boxes, and then add up the volume of each box. We begin by starting with the notion of an open box.

Définition 2.1 (Open box). An **open box** B in $\mathbb{R}^n$ is any set of the form

$$B = \prod_{i=1}^n]a_i, b_i[$$

where $b_i \geq a_i$ are real numbers. We define the **volume** $\text{vol}(B)$ of this box to be the number

$$\text{vol}(B) := \prod_{i=1}^n (b_i - a_i)$$

Définition 2.2 (Outer measure). If $E \subseteq \mathbb{R}^n$ is a set, we define the **outer measure** $m^*(E)$ of E to be the quantity

$$m^*(E) := \inf \left\{ \sum_{j \in J} \text{vol}(B_j) \mid (B_j)_{j \in J} \text{ covers } E; J \text{ countable} \right\}$$

where $(B_j)_{j \in J}$ is a collection of **boxes**.

Lemme 2.1 (Properties of outer measure). The outer measure has the following properties :

1. The empty set $\emptyset$ has outer measure $m^*(\emptyset) = 0$.
2. For every set E we have $m^*(E) \in [0, \infty]$.
3. If $A \subseteq B \subseteq \mathbb{R}^n$, then $m^*(A) \leq m^*(B)$.
4. If E is a subset of $\mathbb{R}^n$, and $x \in \mathbb{R}^n$, then $m^*(x + E) = m^*(E)$.
5. If $(A_j)_{j \in J}$ are countable collection of subsets of $\mathbb{R}^n$, then

$$m^*\left(\bigcup_{j \in J} A_j\right) \leq \sum_{j \in J} m^*(A_j).$$

DÉMONSTRATION :

1. We can cover $\emptyset$ with $]0, \varepsilon[^n$ for any $\varepsilon > 0$. It means

$$m^*(\emptyset) \leq \text{vol}(]0, \varepsilon[^n) = \varepsilon^n$$

We can conclude the $m^*(\emptyset) = 0$ by the fact that ε is arbitrary.

2. Since $\sum_{j \in J} \text{vol}(B_j)$ is non-negative, we know that $m^*(E) \geq 0$. However, it quite possible that $m^*(E)$ could equal ∞ .
3. Follow from the fact that any cover of B is a cover of A .
4. Similarly, $(B_j)_{j \in J}$ is a cover of E if and only if $(x + B_j)_{j \in J}$ is a cover of $x + E$. Moreover, $\text{vol}(x + B_j) = \text{vol}(B_j)$. We then deduce the claim, because

$$\begin{aligned} m^*(E) &= \inf \left\{ \sum_{j \in J} \text{vol}(B_j) \mid (B_j)_{j \in J} \text{ covers } E \right\} \\ &= \inf \left\{ \sum_{j \in J} \text{vol}(x + B_j) \mid (B_j)_{j \in J} \text{ covers } E \right\} \\ &= \inf \left\{ \sum_{j \in J} \text{vol}(B'_j) \mid (B'_j)_{j \in J} \text{ covers } x + E \right\} \\ &= m^*(x + E) \end{aligned}$$

5. For any $j \in J$, by definition of the outer measure as an infimum, it exists $(B_i^j)_{i \in I_j}$ a countable cover of A_j by boxes such that

$$\sum_{i \in I_j} \text{vol}(B_i^j) \leq m^*(A_j) + \frac{\varepsilon}{2^j}$$

Taking $(B_i^j)_{i \in I_j, j \in J}$ as a cover of $\bigcup_{j \in J} A_j$, it follows

$$m^* \left(\bigcup_{j \in J} A_j \right) \leq \sum_{j \in J} \sum_{i \in I_j} \text{vol}(B_i^j) \leq \sum_{j \in J} \left(m^*(A_j) + \frac{\varepsilon}{2^j} \right) = \varepsilon + \sum_{j \in J} m^*(A_j)$$

□

Théorème 2.1 (Théorème fondamental de l'analyse). *Soit f une fonction de classe C^1 , alors*

$$\forall x \in \mathbb{R}, \quad f(x) = \int_0^x f'(t) \, dt$$