

1 The Algebra of Least Squares

Notation: y is a scalar, and x is a $1 \times K$ vector. Y is an $n \times 1$ vector, and X is an $n \times K$ matrix.

1.1 OLS estimator

As we have learned from the previous lecture, the parameter β in the linear projection model

$$\begin{aligned} y &= x'\beta + e \\ E[xe] &= 0 \end{aligned}$$

can be written as $\beta = (E[xx'])^{-1}E[xy]$. In reality we possess a sample of n observations, not the population. We thus replace the population mean $E[\cdot]$ by the sample mean, and the resulting estimator is

$$\hat{\beta} = \left(\frac{1}{n} \sum_{i=1}^n x_i x_i' \right)^{-1} \frac{1}{n} \sum_{i=1}^n x_i y_i = (X'X)^{-1} X'y.$$

This is one way to motivate the OLS estimator.

Alternatively, we can derive the OLS estimator from minimizing the sum of squared residuals

$$\frac{1}{n} \sum_{i=1}^n (y_i - x_i' \beta)^2,$$

which is the sample counterpart of the MSE. By a similar optimization procedure, we derive exactly the same $\hat{\beta}$.

Some definitions.

- Fitted value: $\hat{Y} = X\hat{\beta}$.
- Residual: $\hat{e} = Y - \hat{Y}$.
- Projector: $P_X = X(X'X)^{-1}X'$; Annihilator: $M_X = I_n - P_X$.
- Idempotent matrix. $P_X P_X = P_X$, $M_X M_X = M_X$. $P_X M_X = M_X P_X = 0$.

Some properties of the residual

- $\hat{e} = Y - \hat{Y} = Y - X\hat{\beta} = M_X Y = M_X (X\beta + e) = M_X e$.
- $X'\hat{e} = X M_X e = 0$.
- $\frac{1}{n} \sum_{i=1}^n \hat{e}_i = 0$ if x_i contains a constant.

1.2 Goodness of Fit

The so-called R-square is the most popular measure of goodness-of-fit in the linear regression. R-square is defined when a constant is included in the regressors. Let $M_\iota = I_n - \frac{1}{n}\iota\iota'$, where ι is

an $n \times 1$ vector of 1's. M_l is the demeaner, that is, $M_l(z_1, \dots, z_n)' = (z_1 - \bar{z}, \dots, z_n - \bar{z})'$, where $\bar{z} = \frac{1}{n} \sum_{i=1}^n z_i$. For any X , we can decompose $Y = P_X Y + M_X Y = \hat{Y} + \hat{e}$. The squared sum is

$$\begin{aligned} Y' M_l Y &= (\hat{Y} + \hat{e})' M_l (\hat{Y} + \hat{e}) \\ &= \hat{Y}' M_l \hat{Y} + 2\hat{Y}' M_l \hat{e} + \hat{e}' M_l \hat{e} \\ &= \hat{Y}' M_l \hat{Y} + \hat{e}' \hat{e}. \end{aligned}$$

where the last equality follows as $\frac{1}{n} \sum_{i=1}^n \hat{e}_i = 0$ so that $M_l \hat{e} = \hat{e}$, and $\hat{Y}' \hat{e} = Y' P_X M_X e = 0$. R-square is defined as $1 - \hat{e}' \hat{e} / Y' M_l Y$.

1.3 Frish-Waugh-Lovell Theorem

If $y_i = x'_1 \beta_1 + x'_2 \beta_2 + e$, then $\hat{\beta}_1 = (X'_1 M_2 X_1)^{-1} X'_1 M_2 Y$.

2 Statistical Properties of Least Squares

2.1 Bias and Variance

When we discuss the statistical properties of the least squares in finite sample, we assume

$$\begin{aligned} y &= x' \beta + e \\ E[e|x] &= 0, \end{aligned}$$

which is equivalent to assume $E[y|x] = x' \beta$ so the linear projection model and the conditional mean model coincide. In other words, the following statistical properties hold only when the conditional mean is a linear function.

- Unbiasedness: $E[\hat{\beta}|X] = E[(X'X)^{-1} X'Y|X] = E[(X'X)^{-1} X'(X'\beta + e)|X] = \beta$.
- Variance $\text{var}(\hat{\beta}|X) = E[(\hat{\beta} - E\hat{\beta})(\hat{\beta} - E\hat{\beta})'|X] = E[(X'X)^{-1} X'ee'X(X'X)^{-1}|X] = (X'X)^{-1} X'E[ee'|X]X(X'X)^{-1}$. Under the *homoskedasticity* assumption,¹ we simplify it as

$$\text{var}(\hat{\beta}) = (X'X)^{-1} X'(\sigma^2 I_n)X(X'X)^{-1} = \sigma^2 (X'X)^{-1}.$$

2.2 Gauss-Markov Theorem

Gauss-Markov theorem justifies the OLS estimator as the efficient estimator among all linear unbiased ones. Efficient here means that it enjoys the smallest variance in a family of estimators.

¹The error term is homoskedastic if $E[e_i^2|X] = \sigma^2$ for all i . Homoskedasticity is a restrictive assumption. An example of *heteroskedasticity*: consumption = income + e. A rich family has more variation in consumption than a poor family.

There are numerous linearly unbiased estimators. For example, $(Z'X)^{-1}Z'y$ for $z_i = x_i^2$ is unbiased because $E[(Z'X)^{-1}Z'y] = E[(Z'X)^{-1}Z'(X\beta + e)] = \beta$.

Let $\tilde{\beta} = A'y$ be a generic linear estimator, where A is any $n \times K$ functions of X . As

$$E[A'y|X] = E[A'(X\beta + e)|X] = AX\beta.$$

So the unbiasedness of $\tilde{\beta}$ implies $AX = I_n$. Moreover, the variance

$$\text{var}(A'y|X) = E[(A'y - \beta)(A'y - \beta)'|X] = E[A'ee'A|X] = \sigma^2 A'A.$$

Let $A = C + X(X'X)^{-1}$.

$$\begin{aligned} & A'A - (X'X)^{-1} \\ &= (C + X(X'X)^{-1})'(C + X(X'X)^{-1}) - (X'X)^{-1} \\ &= C'C + (X'X)^{-1}X'C + C'X(X'X)^{-1} = C'C, \end{aligned}$$

where the last equality follows as

$$(X'X)^{-1}X'C = (X'X)^{-1}X'(A - X(X'X)^{-1}) = (X'X)^{-1} - (X'X)^{-1} = 0.$$

The variance of any $\tilde{\beta}$ is no smaller than the OLS estimator $\hat{\beta}$.

However, notice that Gauss-Markov theorem is derived only in the conditional mean model under homoskedasticity. These conditions are restrictive.

2.3 Variance Estimation

Under homoskedasticity, $\text{var}(\hat{\beta}) = \sigma^2(X'X)^{-1}$. Popular estimator of σ^2 is the sample mean of the residuals $\hat{\sigma}^2 = \frac{1}{n}\hat{e}'\hat{e}$ or the unbiased one $s^2 = \frac{1}{n-K}\hat{e}'\hat{e}$.

Under heteroskedasticity,

$$\text{var}(\hat{\beta}) = (X'X)^{-1}X'DX(X'X)^{-1}$$

where $D = \text{diag}(\sigma_1^2, \dots, \sigma_n^2)$. We propose to estimate $\hat{\sigma}_i^2 = \hat{e}_i^2$ so that $X'DX = \sum_{i=1}^n x_i x_i' \hat{e}_i^2$.

2.4 Normal Regression

If we assume $e_i \sim N(0, \sigma^2)$ is independent of x_i , then $y_i \sim N(x_i'\beta, \sigma^2)$. The likelihood of observing a sample $(y_i, x_i)_{i=1}^n$ is

$$\prod_{i=1}^n \frac{1}{\sqrt{2\pi}\sigma} \exp\left(-\frac{1}{2\sigma^2}(y_i - x_i'\beta)^2\right),$$

and the log-likelihood function is

$$L(\beta, \sigma) = -\frac{n}{2} \log 2\pi - n \log \sigma - \frac{1}{2\sigma^2} \sum_{i=1}^n (y_i - x_i' \beta)^2.$$

Therefore, the maximum likelihood estimator coincides with the OLS estimator.

Moreover, consider the statistic

$$T_k = \frac{\hat{\beta}_k - \beta_k}{\sqrt{s^2 (X'X)^{-1}_{kk}}} = \frac{\eta'_k (X'X)^{-1} X' e / \sigma}{\sqrt{\frac{s^2}{\sigma^2} \eta'_k (X'X)^{-1} \eta_k}} = \frac{a'_k \frac{e}{\sigma} / \sqrt{a'_k a_k}}{\sqrt{s^2 / \sigma^2}}$$

where $\eta_k = (1 \{l = k\})_{l=1, \dots, K}$ is the selector, and $a_k = \eta'_k (X'X)^{-1} X'$. It is easy to see that the numerator follows a standard normal, and the denominator follows a χ^2 with the degree of freedom $n - K$. Therefore $T_k \sim t(n - K)$.