



School of Computer Science
Faculty of Science
National University of Engineering

Test 5

Topics: Iterative Solution Algorithm for Unconstrained NLOP

Subject: Computational Mathematics

Period: 2020.1

Consider the following unconstrained NLOP:

$$\min_{(x_1, x_2) \in \mathbb{R}^2} f(x_1, x_2),$$

where $f(x_1, x_2)$ is defined by

$$14 \cdot \sqrt{(x_1 - 7)^2 + (x_2 - 2)^2} + 20 \cdot \sqrt{(x_1 - 5)^2 + (x_2 + 3)^2} + 30 \cdot \sqrt{(x_1 + 6)^2 + (x_2 - 4)^2}.$$

1. (3 pts.) Plot the graph of f over an appropriate domain.
2. First attempt.
 - (a) (2 pts.) Starting from the point $x^0 := (0, 0)$ conduct iterations of the Generic Algorithm for Unconstrained Nonlinear Optimization Problems using the steepest descent search direction and an exact line search to solve this problem.
 - (b) (4 pts.) Implement a program in Python for the previous task.
 - (c) (2 pts.) Is the point at which the algorithm terminates guaranteed to be a local or global optimum? Why?
3. Second attempt.

- (a) (2 pts.) Starting from the point $x^0 := (0, 0)$ conduct iterations of the Generic Algorithm for Unconstrained Nonlinear Optimization Problems using the Newton's method search direction and the Armijo rule with $\sigma = 10^{-1}$, $\beta = 1/2$ and $s = 1$.
- (b) (5 pts.) Implement a program in Python for the previous task.
- (c) (2 pts.) Is the point at which the algorithm terminates guaranteed to be a local or global optimum? Why?

August 26, 2020