OSHPD
Office of Statewide Health Planning and Development

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators (Conditions), 2015 (January – September) ‡

County	Hospital Name	Acute Myocardial Infarction	Acute Stroke				Gastro-Intestinal Hemorrhage	Heart Failure	Hip Fracture	Pneumonia
			Total	Hemorrhagic	Ischemic	Subarachnoid				
		NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.								
	STATEWIDE	5.9% (2,162;36,453)	8.7% (3,816;43,654)	20.9% (1,752;8,391)	5.0% (1,659;33,331)	21.0% (405;1,932)	2.3% (823;36,420)	2.9% (1,936;67,612)	2.0% (363;18,253)	3.3% (1,547;46,752)
Alameda	Alameda Hospital	0.0% (0;16)	5.8% (2;45)	0.0% (0;4)	5.1% (2;41)		2.4% (1;46)	1.1% (1;79)	0.0% (0;21)	1.9% (1;56)
	Alta Bates Summit Medical Center**	4.0% (15;314)	5.9% (15;277)	24.4% (8;15)	**2.3% (7;261) Better**		1.8% (7;246)	**1.0% (8;450) Better**	3.8% (2;55)	1.8% (6;171)
	Alta Bates Summit Medical Center – Alta Bates Campus**	3.8% (2;25)	**5.8% (18;222) Better**	16.4% (14;74)	2.3% (4;142)	0.0% (0;6)	1.2% (2;158)	2.2% (6;186)	2.2% (2;87)	2.4% (5;111)
	Eden Medical Center	2.4% (1;17)	**6.5% (38;367) Better**	16.7% (21;91)	3.0% (10;234)	16.5% (7;42)	0.6% (1;103)	1.9% (6;188)	2.5% (2;84)	**0.8% (2;123) Better**
	Highland Hospital	10.6% (6;81)	7.9% (12;130)	19.6% (8;36)	2.0% (1;90)	27.2% (3;4)	1.2% (1;86)	3.8% (8;314)	0.0% (0;23)	2.4% (2;121)
	Kaiser Foundation Hospital – Fremont	2.2% (1;42)	8.4% (10;70)	11.4% (2;14)	6.7% (7;54)		0.0% (0;68)	3.9% (9;109)	3.2% (1;50)	3.0% (8;113)
	Kaiser Foundation Hospital – Oakland/Richmond	3.5% (8;102)	**4.9% (22;334) Better**	**11.3% (9;52) Better**	**3.0% (12;277) Better**	8.0% (1;5)	1.6% (5;217)	**1.9% (20;519) Better**	0.9% (1;106)	1.4% (3;105)
	Kaiser Foundation Hospital – San Leandro	5.5% (6;46)	5.7% (12;99)	19.2% (6;13)	2.7% (6;85)		2.5% (4;92)	1.6% (6;181)	2.8% (2;69)	3.0% (12;179)
	Saint Rose Hospital	7.7% (16;164)	10.4% (3;45)		6.4% (3;43)		4.4% (4;109)	**0.5% (1;164) Better**	0.0% (0;22)	0.7% (1;125)
	San Leandro Hospital	3.2% (1;11)	16.5% (3;20)		13.1% (2;18)		0.0% (0;84)	**0.0% (0;127) Better**	0.0% (0;17)	1.3% (1;74)
	Valleycare Medical Center	7.0% (6;84)	13.4% (6;46)	39.3% (3;4)	6.5% (3;42)		1.5% (2;79)	**1.2% (5;196) Better**	4.9% (1;55)	**0.6% (2;216) Better**

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 5.0, and OSHPD hospital patient discharge data, 2015. See Technical Note for details about methodology.

‡ Data are reported for January – September 2015 due to coding changes from ICD-9-CM to ICD-10-CM/PCS, which began October 1, 2015. Comparisons of IMIs across years should be made with caution since previous years' results are based on 12 months of data, while this analysis is based on 9 months of data.

*****Hospital name changed from 2014 to 2015, see Technical Note for details.***

Note: Blank cells indicate that fewer than three conditions were treated.

Definition of Terms: Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. Total Deaths: number of patients that died in this hospital. Total Cases: number of patients with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.

OSHPD
Office of Statewide Health Planning and Development

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators (Conditions), 2015 (January – September) ‡

County	Hospital Name	Acute Myocardial Infarction	Acute Stroke				Gastro-Intestinal Hemorrhage	Heart Failure	Hip Fracture	Pneumonia
			Total	Hemorrhagic	Ischemic	Subarachnoid				
		NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.								
	STATEWIDE	5.9% (2,162;36,453)	8.7% (3,816;43,654)	20.9% (1,752;8,391)	5.0% (1,659;33,331)	21.0% (405;1,932)	2.3% (823;36,420)	2.9% (1,936;67,612)	2.0% (363;18,253)	3.3% (1,547;46,752)
Alameda (continued)	Washington Hospital – Fremont	5.3% (7;153)	9.6% (29;244)	16.3% (10;47)	7.5% (14;168)	23.5% (5;29)	1.5% (3;205)	2.8% (9;299)	**8.9% (5;89) Worse**	4.9% (19;296)
Amador	Sutter Amador Hospital	6.8% (1;6)	6.6% (2;70)	0.0% (0;3)	4.2% (2;67)		0.0% (0;57)	0.0% (0;70)	0.0% (0;32)	3.5% (4;87)
Butte	Enloe Medical Center – Esplanade	8.0% (24;308)	7.3% (21;229)	19.3% (11;45)	3.4% (8;182)		1.4% (6;250)	1.8% (9;324)	2.9% (3;121)	3.1% (15;306)
	Feather River Hospital	2.2% (1;26)	**3.4% (3;49) Better**	8.0% (1;5)	**1.0% (1;43) Better**		0.0% (0;67)	1.5% (4;104)	0.0% (0;62)	1.5% (3;100)
	Orchard Hospital**	0.0% (0;5)	0.0% (0;3)				0.0% (0;7)	0.0% (0;8)		0.0% (0;51)
	Oroville Hospital	2.3% (3;45)	**3.2% (3;126) Better**		**1.9% (3;124) Better**		0.5% (1;123)	**0.4% (2;240) Better**	0.6% (1;56)	0.0% (0;8)
Calaveras	Mark Twain Medical Center	0.0% (0;3)	48.2% (2;12)		27.3% (2;12)			0.0% (0;23)	0.0% (0;28)	0.0% (0;43)
Colusa	Colusa Regional Medical Center	0.0% (0;4)	0.0% (0;5)		0.0% (0;5)		5.7% (1;9)	2.8% (1;19)		0.0% (0;31)
Contra Costa	Contra Costa Regional Medical Center	0.0% (0;40)	4.5% (2;76)		2.5% (2;76)		0.0% (0;61)	1.2% (2;213)	0.0% (0;15)	1.9% (2;108)
	Doctors Medical Center – San Pablo	16.5% (3;22)	0.0% (0;8)		0.0% (0;8)			3.1% (2;56)		1.6% (1;29)
	John Muir Medical Center – Concord Campus	5.4% (14;251)	8.5% (10;143)	30.9% (4;6)	4.0% (6;136)		3.1% (6;172)	2.2% (16;405)	2.2% (1;56)	2.0% (4;175)
	John Muir Medical Center – Walnut Creek Campus	3.6% (6;84)	8.7% (60;360)	17.4% (31;106)	5.7% (21;234)	34.4% (8;20)	1.5% (6;279)	2.2% (15;317)	3.3% (4;139)	2.2% (9;216)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 5.0, and OSHPD hospital patient discharge data, 2015. See Technical Note for details about methodology.

‡ Data are reported for January – September 2015 due to coding changes from ICD-9-CM to ICD-10-CM/PCS, which began October 1, 2015. Comparisons of IMIs across years should be made with caution since previous years' results are based on 12 months of data, while this analysis is based on 9 months of data.

*****Hospital name changed from 2014 to 2015, see Technical Note for details.***

Note: Blank cells indicate that fewer than three conditions were treated.

Definition of Terms: Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. Total Deaths: number of patients that died in this hospital. Total Cases: number of patients with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.

OSHPD
Office of Statewide Health Planning and Development

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators (Conditions), 2015 (January – September) ‡

County	Hospital Name	Acute Myocardial Infarction	Acute Stroke				Gastro-Intestinal Hemorrhage	Heart Failure	Hip Fracture	Pneumonia
			Total	Hemorrhagic	Ischemic	Subarachnoid				
		NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.								
	STATEWIDE	5.9% (2,162;36,453)	8.7% (3,816;43,654)	20.9% (1,752;8,391)	5.0% (1,659;33,331)	21.0% (405;1,932)	2.3% (823;36,420)	2.9% (1,936;67,612)	2.0% (363;18,253)	3.3% (1,547;46,752)
Contra Costa (continued)	Kaiser Foundation Hospital – Antioch	**1.1% (1;46) Better**	7.2% (24;161)	18.6% (14;30)	4.1% (10;126)	0.0% (0;5)	1.3% (2;117)	1.3% (5;194)	4.4% (2;58)	3.7% (6;108)
	Kaiser Foundation Hospital – Walnut Creek	3.6% (5;174)	**4.5% (10;196) Better**	**4.8% (2;27) Better**	3.9% (8;167)		2.0% (6;216)	2.0% (14;370)	3.1% (3;146)	**1.6% (7;265) Better**
	San Ramon Regional Medical Center	5.5% (4;69)	10.3% (8;89)		5.5% (6;87)		2.6% (1;70)	3.1% (4;91)	0.0% (0;28)	1.3% (2;115)
	Sutter Delta Medical Center	5.9% (14;161)	3.9% (3;77)	36.0% (3;5)	**0.0% (0;71) Better**		0.0% (0;80)	2.1% (9;271)	1.8% (1;46)	**0.8% (2;165) Better**
Del Norte	Sutter Coast Hospital	2.5% (1;13)	12.4% (6;28)		7.1% (4;26)		0.0% (0;31)	4.0% (6;56)	9.0% (1;8)	**0.0% (0;73) Better**
El Dorado	Barton Memorial Hospital		0.0% (0;5)		0.0% (0;5)		0.0% (0;20)	3.2% (1;28)	0.0% (0;16)	1.8% (1;69)
	Marshall Medical Center	10.0% (6;20)	10.2% (7;75)	22.6% (3;4)	6.4% (4;71)		0.6% (1;72)	1.6% (4;143)	6.5% (3;23)	1.7% (6;210)
Fresno	Adventist Medical Center – Reedley	0.0% (0;5)	0.0% (0;14)		0.0% (0;14)			0.0% (0;10)		0.0% (0;40)
	Clovis Community Medical Center	3.4% (4;100)	**2.6% (2;124) Better**	13.0% (1;8)	**1.0% (1;116) Better**		1.9% (4;187)	1.6% (8;304)	5.8% (4;111)	1.8% (6;226)
	Coalinga Regional Medical Center		0.0% (0;3)		0.0% (0;3)			0.0% (0;9)		0.0% (0;14)
	Community Regional Medical Center – Fresno *(data for the month of September are incomplete)*	5.9% (23;472)	8.8% (90;761)	21.4% (52;217)	3.8% (21;468)	31.7% (17;76)	1.3% (6;428)	1.8% (13;738)	0.0% (0;93)	3.5% (18;486)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 5.0, and OSHPD hospital patient discharge data, 2015. See Technical Note for details about methodology.

‡ Data are reported for January – September 2015 due to coding changes from ICD-9-CM to ICD-10-CM/PCS, which began October 1, 2015. Comparisons of IMIs across years should be made with caution since previous years' results are based on 12 months of data, while this analysis is based on 9 months of data.

*****Hospital name changed from 2014 to 2015, see Technical Note for details.***

Note: Blank cells indicate that fewer than three conditions were treated.

Definition of Terms: Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. Total Deaths: number of patients that died in this hospital. Total Cases: number of patients with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.

OSHPD
Office of Statewide Health Planning and Development

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators (Conditions), 2015 (January – September) ‡

County	Hospital Name	Acute Myocardial Infarction	Acute Stroke				Gastro-Intestinal Hemorrhage	Heart Failure	Hip Fracture	Pneumonia
			Total	Hemorrhagic	Ischemic	Subarachnoid				
		NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.								
	STATEWIDE	5.9% (2,162;36,453)	8.7% (3,816;43,654)	20.9% (1,752;8,391)	5.0% (1,659;33,331)	21.0% (405;1,932)	2.3% (823;36,420)	2.9% (1,936;67,612)	2.0% (363;18,253)	3.3% (1,547;46,752)
Fresno (continued)	Fresno Heart and Surgical Hospital	0.0% (0;37)					0.0% (0;23)	1.7% (2;85)		0.0% (0;14)
	Kaiser Foundation Hospital – Fresno	2.1% (2;72)	5.9% (14;174)	12.1% (6;32)	3.6% (7;141)		2.1% (2;92)	2.1% (7;212)	0.0% (0;98)	5.2% (11;109)
	Saint Agnes Medical Center	5.1% (17;312)	9.9% (34;339)	23.7% (14;38)	5.8% (20;298)	0.0% (0;3)	2.7% (13;311)	2.2% (21;623)	**5.8% (9;191) Worse**	2.7% (16;473)
Glenn	Glenn Medical Center							0.0% (0;11)		0.0% (0;9)
Humboldt	Jerold Phelps Community Hospital							0.0% (0;4)		0.0% (0;6)
	Mad River Community Hospital		0.0% (0;14)		0.0% (0;13)		0.0% (0;12)	4.1% (2;40)	5.3% (1;28)	1.4% (1;68)
	Redwood Memorial Hospital		29.0% (1;7)		0.0% (0;6)		0.0% (0;11)	5.8% (2;17)	0.0% (0;5)	13.9% (3;29)
	Saint Joseph Hospital – Eureka	**2.5% (5;162) Better**	7.9% (4;72)	18.2% (1;5)	4.6% (3;67)		2.0% (4;87)	1.8% (6;167)	0.0% (0;41)	3.0% (3;85)
Imperial	El Centro Regional Medical Center	11.2% (7;34)	5.6% (5;91)	11.9% (1;5)	2.7% (3;85)		1.8% (2;98)	2.8% (5;127)	**10.1% (3;44) Worse**	**0.9% (2;134) Better**
	Pioneers Memorial Healthcare District	10.0% (4;17)	5.2% (2;56)	0.0% (0;4)	4.0% (2;52)		**0.0% (0;81) Better**	1.9% (2;103)	0.0% (0;19)	1.9% (3;130)
Inyo	Northern Inyo Hospital	0.0% (0;4)	0.0% (0;11)		0.0% (0;9)		13.2% (1;14)	10.1% (1;9)	0.0% (0;12)	2.6% (1;24)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 5.0, and OSHPD hospital patient discharge data, 2015. See Technical Note for details about methodology.

‡ Data are reported for January – September 2015 due to coding changes from ICD-9-CM to ICD-10-CM/PCS, which began October 1, 2015. Comparisons of IMIs across years should be made with caution since previous years' results are based on 12 months of data, while this analysis is based on 9 months of data.

*****Hospital name changed from 2014 to 2015, see Technical Note for details.***

Note: Blank cells indicate that fewer than three conditions were treated.

Definition of Terms: Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. Total Deaths: number of patients that died in this hospital. Total Cases: number of patients with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.

OSHPD
Office of Statewide Health Planning and Development

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators (Conditions), 2015 (January – September) ‡

County	Hospital Name	Acute Myocardial Infarction	Acute Stroke				Gastro-Intestinal Hemorrhage	Heart Failure	Hip Fracture	Pneumonia
			Total	Hemorrhagic	Ischemic	Subarachnoid				
		NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.								
	STATEWIDE	5.9% (2,162;36,453)	8.7% (3,816;43,654)	20.9% (1,752;8,391)	5.0% (1,659;33,331)	21.0% (405;1,932)	2.3% (823;36,420)	2.9% (1,936;67,612)	2.0% (363;18,253)	3.3% (1,547;46,752)
Kern	Bakersfield Heart Hospital	6.1% (8;134)	8.9% (4;37)	22.9% (1;3)	5.0% (3;34)		0.0% (0;60)	1.8% (7;251)		1.7% (2;109)
	Bakersfield Memorial Hospital	7.0% (22;304)	10.1% (19;209)	19.8% (5;33)	5.8% (11;163)	39.5% (3;13)	1.7% (3;152)	3.5% (14;345)	3.0% (3;100)	3.9% (9;217)
	Delano Regional Medical Center	0.0% (0;3)	0.0% (0;11)		0.0% (0;10)		7.4% (2;40)	0.0% (0;55)	54.1% (1;9)	0.0% (0;73)
	Good Samaritan Hospital – Bakersfield							5.4% (1;32)		0.0% (0;58)
	Kern Medical Center	16.0% (2;20)	22.2% (4;35)	**84.1% (3;8) Worse**	7.1% (1;25)		3.6% (1;44)	0.0% (0;86)	0.0% (0;19)	8.5% (3;50)
	Kern Valley Healthcare District		0.0% (0;3)		0.0% (0;3)		0.0% (0;9)	3.7% (1;22)		2.7% (1;76)
	Mercy Hospital – Bakersfield	2.1% (1;19)	7.4% (9;95)	19.7% (5;18)	3.7% (3;74)	16.9% (1;3)	3.8% (4;117)	**0.0% (0;159) Better**	0.0% (0;104)	1.5% (4;244)
	Ridgecrest Regional Hospital	0.0% (0;5)	20.3% (2;19)		6.9% (1;17)		4.2% (1;28)	2.4% (1;40)	0.0% (0;6)	5.5% (4;65)
	San Joaquin Community Hospital	6.3% (25;349)	7.2% (38;409)	16.7% (19;90)	4.1% (16;314)	20.1% (3;5)	1.4% (5;247)	1.7% (11;460)	4.3% (5;108)	2.6% (9;254)
	Tehachapi Hospital									0.0% (0;9)
Kings	Adventist Medical Center	5.1% (8;83)	5.9% (7;151)	16.3% (1;5)	3.3% (6;146)		2.1% (5;178)	1.7% (6;291)	6.9% (3;73)	2.6% (6;221)
Lake	Saint Helena Hospital – Clearlake		0.0% (0;6)		0.0% (0;5)		0.0% (0;22)	3.3% (1;35)		4.0% (3;70)
	Sutter Lakeside Hospital		15.6% (2;25)		8.9% (2;25)		0.0% (0;19)	3.6% (2;40)	20.9% (1;22)	5.2% (3;66)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 5.0, and OSHPD hospital patient discharge data, 2015. See Technical Note for details about methodology.

‡ Data are reported for January – September 2015 due to coding changes from ICD-9-CM to ICD-10-CM/PCS, which began October 1, 2015. Comparisons of IMIs across years should be made with caution since previous years' results are based on 12 months of data, while this analysis is based on 9 months of data.

*****Hospital name changed from 2014 to 2015, see Technical Note for details.***

Note: Blank cells indicate that fewer than three conditions were treated.

Definition of Terms: Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. Total Deaths: number of patients that died in this hospital. Total Cases: number of patients with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.

OSHPD
Office of Statewide Health Planning and Development

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators (Conditions), 2015 (January – September) ‡

County	Hospital Name	Acute Myocardial Infarction	Acute Stroke				Gastro-Intestinal Hemorrhage	Heart Failure	Hip Fracture	Pneumonia
			Total	Hemorrhagic	Ischemic	Subarachnoid				
		NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.								
	STATEWIDE	5.9% (2,162;36,453)	8.7% (3,816;43,654)	20.9% (1,752;8,391)	5.0% (1,659;33,331)	21.0% (405;1,932)	2.3% (823;36,420)	2.9% (1,936;67,612)	2.0% (363;18,253)	3.3% (1,547;46,752)
Lassen	Banner Lassen Medical Center		66.5% (1;4)		37.1% (1;4)		0.0% (0;7)	2.5% (1;13)		0.0% (0;43)
Los Angeles	Alhambra Hospital Medical Center	0.0% (0;12)	3.5% (2;66)	7.1% (1;16)	2.4% (1;50)		1.5% (1;84)	**0.0% (0;128) Better**	0.0% (0;31)	2.5% (10;190)
	Antelope Valley Hospital	8.4% (22;309)	11.7% (29;271)	23.6% (11;45)	7.4% (15;220)	30.9% (3;6)	2.1% (7;244)	5.0% (13;243)	4.0% (3;117)	**5.9% (19;312) Worse**
	Beverly Hospital	5.0% (12;155)	9.1% (7;96)	12.4% (1;18)	6.1% (6;77)		1.6% (2;128)	1.2% (3;195)	0.0% (0;43)	**1.3% (4;186) Better**
	California Hospital Medical Center – Los Angeles	4.3% (4;83)	6.7% (19;224)	16.8% (10;45)	3.3% (7;174)	24.2% (2;5)	1.1% (2;151)	1.2% (4;336)	4.3% (1;22)	1.1% (1;132)
	Cedars Sinai Medical Center	5.2% (27;445)	**5.1% (48;573) Better**	**13.4% (23;118) Better**	**2.8% (18;388) Better**	**9.3% (7;67) Better**	**1.1% (9;499) Better**	2.3% (47;1,070)	3.7% (7;219)	**1.8% (11;424) Better**
	Centinela Hospital Medical Center	**1.7% (7;217) Better**	**3.8% (13;231) Better**	**11.5% (7;42) Better**	**1.5% (5;183) Better**	23.9% (1;6)	1.1% (5;246)	**0.4% (7;976) Better**	0.0% (0;35)	**0.3% (1;191) Better**
	Citrus Valley Medical Center – Inter Community Campus	8.8% (19;311)	6.0% (5;56)	6.1% (1;16)	4.1% (2;38)		1.4% (2;107)	1.3% (4;221)	0.0% (0;38)	1.6% (3;113)
	Citrus Valley Medical Center – Queen of the Valley Campus	4.1% (3;31)	8.3% (21;244)	19.4% (11;46)	4.6% (9;193)	52.7% (1;5)	1.5% (2;171)	1.4% (5;261)	2.4% (1;51)	1.6% (4;218)
	City of Hope Helford Clinical Research Hospital		14.9% (1;7)	37.0% (1;4)	0.0% (0;3)		0.0% (0;19)	15.0% (1;7)		**7.8% (8;93) Worse**
	Coast Plaza Hospital	2.9% (1;14)	8.7% (2;24)	44.3% (2;5)	0.0% (0;19)		4.1% (3;45)	4.2% (5;102)	0.0% (0;10)	4.2% (3;85)
	College Medical Center	4.7% (2;9)	0.0% (0;18)		0.0% (0;17)		2.6% (2;36)	0.0% (0;53)	0.0% (0;11)	5.1% (3;41)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 5.0, and OSHPD hospital patient discharge data, 2015. See Technical Note for details about methodology.

‡ Data are reported for January – September 2015 due to coding changes from ICD-9-CM to ICD-10-CM/PCS, which began October 1, 2015. Comparisons of IMIs across years should be made with caution since previous years' results are based on 12 months of data, while this analysis is based on 9 months of data.

*****Hospital name changed from 2014 to 2015, see Technical Note for details.***

Note: Blank cells indicate that fewer than three conditions were treated.

Definition of Terms: Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. Total Deaths: number of patients that died in this hospital. Total Cases: number of patients with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.

OSHPD
Office of Statewide Health Planning and Development

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators (Conditions), 2015 (January – September) ‡

County	Hospital Name	Acute Myocardial Infarction	Acute Stroke				Gastro-Intestinal Hemorrhage	Heart Failure	Hip Fracture	Pneumonia
			Total	Hemorrhagic	Ischemic	Subarachnoid				
		NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.								
	STATEWIDE	5.9% (2,162;36,453)	8.7% (3,816;43,654)	20.9% (1,752;8,391)	5.0% (1,659;33,331)	21.0% (405;1,932)	2.3% (823;36,420)	2.9% (1,936;67,612)	2.0% (363;18,253)	3.3% (1,547;46,752)
Los Angeles (continued)	Community Hospital Long Beach	6.3% (4;25)	9.4% (1;23)		5.4% (1;23)		1.3% (1;51)	4.6% (5;76)	0.0% (0;44)	3.0% (7;109)
	Community and Mission Hospital of Huntington Park – Slauson	3.3% (3;42)	0.0% (0;15)		0.0% (0;15)		0.0% (0;50)	1.1% (1;129)	0.0% (0;8)	3.1% (2;68)
	Doctors Hospital of West Covina, Inc.							0.0% (0;7)		0.0% (0;14)
	East Los Angeles Doctors Hospital	3.1% (3;42)	8.5% (1;14)		5.9% (1;12)		4.5% (3;40)	4.7% (6;83)	0.0% (0;8)	5.2% (4;76)
	Encino Hospital Medical Center	0.0% (0;12)	4.1% (1;12)		2.3% (1;12)		2.1% (1;29)	**0.0% (0;67) Better**	0.0% (0;6)	0.0% (0;32)
	Foothill Presbyterian Hospital – Johnston Memorial	8.0% (2;21)	5.1% (2;57)	22.9% (2;11)	0.0% (0;46)		4.4% (4;97)	1.9% (3;125)	11.5% (2;41)	1.1% (2;145)
	Gardens Regional Hospital and Medical Center	2.1% (1;19)	26.9% (1;11)		0.0% (0;9)		1.8% (1;37)	2.9% (3;84)	0.0% (0;6)	1.5% (2;75)
	Garfield Medical Center	6.7% (16;223)	**5.1% (12;247) Better**	11.6% (5;58)	2.5% (5;184)	25.6% (2;5)	2.3% (5;151)	**1.1% (5;323) Better**	0.0% (0;51)	4.2% (19;282)
	Glendale Adventist Medical Center	4.1% (12;261)	**6.6% (31;298) Better**	**13.5% (12;62) Better**	4.2% (13;204)	17.0% (6;32)	**0.3% (1;192) Better**	**0.9% (9;468) Better**	1.4% (2;104)	**2.1% (17;403) Better**
	Glendale Memorial Hospital and Medical Center	5.4% (12;163)	10.2% (7;74)	32.9% (5;8)	3.6% (2;66)		1.7% (3;126)	2.4% (10;319)	0.0% (0;67)	**1.4% (5;224) Better**
	Glendora Community Hospital	0.0% (0;3)	0.0% (0;4)		0.0% (0;4)		3.4% (1;21)	7.1% (3;19)	0.0% (0;4)	1.9% (2;59)
	Good Samaritan Hospital – Los Angeles	8.0% (16;208)	9.5% (15;143)	27.1% (11;38)	3.4% (3;103)		2.3% (6;163)	2.6% (10;318)	2.6% (1;47)	2.4% (6;143)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 5.0, and OSHPD hospital patient discharge data, 2015. See Technical Note for details about methodology.

‡ Data are reported for January – September 2015 due to coding changes from ICD-9-CM to ICD-10-CM/PCS, which began October 1, 2015. Comparisons of IMIs across years should be made with caution since previous years' results are based on 12 months of data, while this analysis is based on 9 months of data.

*****Hospital name changed from 2014 to 2015, see Technical Note for details.***

Note: Blank cells indicate that fewer than three conditions were treated.

Definition of Terms: Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. Total Deaths: number of patients that died in this hospital. Total Cases: number of patients with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.

OSHPD
Office of Statewide Health Planning and Development

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators (Conditions), 2015 (January – September) ‡

County	Hospital Name	Acute Myocardial Infarction	Acute Stroke				Gastro-Intestinal Hemorrhage	Heart Failure	Hip Fracture	Pneumonia
			Total	Hemorrhagic	Ischemic	Subarachnoid				
		NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.								
	STATEWIDE	5.9% (2,162;36,453)	8.7% (3,816;43,654)	20.9% (1,752;8,391)	5.0% (1,659;33,331)	21.0% (405;1,932)	2.3% (823;36,420)	2.9% (1,936;67,612)	2.0% (363;18,253)	3.3% (1,547;46,752)
Los Angeles (continued)	Greater El Monte Community Hospital	11.2% (1;5)	9.0% (1;19)		5.2% (1;19)		1.5% (1;35)	0.0% (0;40)		**0.0% (0;96) Better**
	Henry Mayo Newhall Hospital	**0.0% (0;145) Better**	**0.0% (0;106) Better**	0.0% (0;13)	**0.0% (0;88) Better**	0.0% (0;5)	**0.0% (0;168) Better**	**0.0% (0;256) Better**	0.0% (0;67)	**0.0% (0;256) Better**
	Hollywood Presbyterian Medical Center	3.7% (6;158)	10.9% (23;160)	24.0% (10;30)	6.3% (10;124)	35.9% (3;6)	1.5% (4;143)	**0.9% (3;232) Better**	3.8% (2;46)	2.6% (5;154)
	Huntington Memorial Hospital	5.4% (14;224)	6.5% (25;344)	22.6% (16;71)	**2.7% (8;247) Better**	5.3% (1;26)	1.8% (5;247)	1.6% (9;374)	0.0% (0;162)	2.5% (12;314)
	Kaiser Foundation Hospital – Baldwin Park	4.7% (6;76)	**4.3% (6;192) Better**	**0.0% (0;20) Better**	3.3% (6;171)		1.6% (4;186)	1.3% (5;349)	1.8% (1;96)	1.7% (6;223)
	Kaiser Foundation Hospital – Downey	**2.9% (6;128) Better**	5.8% (12;254)	15.1% (3;20)	3.3% (9;233)		2.0% (4;198)	**0.8% (3;306) Better**	3.7% (2;73)	3.8% (15;303)
	Kaiser Foundation Hospital – Panorama City	**1.8% (2;83) Better**	4.5% (5;116)	11.3% (2;17)	2.6% (3;98)		1.8% (3;95)	1.7% (6;182)	1.4% (1;101)	3.6% (11;167)
	Kaiser Foundation Hospital – South Bay	**2.5% (4;80) Better**	6.1% (9;160)	13.3% (3;27)	3.7% (6;133)		2.3% (5;162)	**1.0% (6;294) Better**	8.7% (2;46)	2.1% (3;100)
	Kaiser Foundation Hospital – Sunset	**3.6% (32;806) Better**	7.7% (63;496)	16.6% (32;176)	4.7% (19;267)	23.2% (12;53)	**0.3% (1;182) Better**	2.6% (21;484)	0.0% (0;40)	3.0% (5;92)
	Kaiser Foundation Hospital – West Los Angeles	3.4% (5;84)	5.7% (13;301)	13.1% (3;24)	3.3% (10;277)		2.8% (7;212)	2.1% (9;271)	0.0% (0;44)	2.2% (4;108)
	Kaiser Foundation Hospital – Woodland Hills	4.4% (6;98)	**2.1% (3;194) Better**	**0.0% (0;19) Better**	**1.7% (3;175) Better**		**0.4% (1;167) Better**	**1.4% (14;404) Better**	2.2% (3;105)	**1.4% (7;231) Better**

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 5.0, and OSHPD hospital patient discharge data, 2015. See Technical Note for details about methodology.

‡ Data are reported for January – September 2015 due to coding changes from ICD-9-CM to ICD-10-CM/PCS, which began October 1, 2015. Comparisons of IMIs across years should be made with caution since previous years' results are based on 12 months of data, while this analysis is based on 9 months of data.

*****Hospital name changed from 2014 to 2015, see Technical Note for details.***

Note: Blank cells indicate that fewer than three conditions were treated.

Definition of Terms: Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. Total Deaths: number of patients that died in this hospital. Total Cases: number of patients with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.

OSHPD
Office of Statewide Health Planning and Development

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators (Conditions), 2015 (January – September) ‡

County	Hospital Name	Acute Myocardial Infarction	Acute Stroke				Gastro-Intestinal Hemorrhage	Heart Failure	Hip Fracture	Pneumonia
			Total	Hemorrhagic	Ischemic	Subarachnoid				
		NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.								
	STATEWIDE	5.9% (2,162;36,453)	8.7% (3,816;43,654)	20.9% (1,752;8,391)	5.0% (1,659;33,331)	21.0% (405;1,932)	2.3% (823;36,420)	2.9% (1,936;67,612)	2.0% (363;18,253)	3.3% (1,547;46,752)
Los Angeles (continued)	Keck Hospital of University of Southern California	7.5% (5;63)	**6.2% (43;298) Better**	**13.3% (17;98) Better**	5.2% (12;114)	**12.1% (14;86) Better**	1.0% (1;69)	4.2% (11;147)	0.0% (0;8)	6.8% (5;40)
	Lakewood Regional Medical Center	7.2% (20;222)	9.9% (26;195)	26.8% (18;56)	4.7% (7;135)	14.5% (1;4)	2.8% (4;132)	2.2% (8;320)	5.5% (4;80)	2.5% (7;202)
	Long Beach Memorial Medical Center	6.4% (22;298)	7.1% (37;430)	19.5% (17;77)	3.4% (13;316)	16.8% (7;37)	**1.0% (5;270) Better**	**1.4% (7;477) Better**	5.8% (3;72)	3.2% (9;266)
	Los Angeles Community Hospital	0.0% (0;10)	8.9% (1;25)		6.9% (1;23)		0.0% (0;95)	**0.0% (0;143) Better**	0.0% (0;9)	1.9% (3;95)
	Los Angeles County/Harbor – UCLA Medical Center	5.8% (10;165)	6.9% (13;172)	18.2% (7;39)	3.0% (3;120)	16.8% (3;13)	1.3% (2;148)	2.5% (5;257)	7.7% (2;22)	2.0% (2;96)
	Los Angeles County/Olive View – UCLA Medical Center	5.2% (3;68)	5.5% (2;54)	12.7% (1;6)	3.5% (1;47)		0.0% (0;117)	**0.0% (0;262) Better**		0.0% (0;98)
	Los Angeles County/University of Southern California Medical Center	4.1% (6;131)	8.7% (22;215)	24.2% (12;51)	3.6% (7;149)	28.6% (3;15)	2.2% (7;224)	3.1% (13;519)	4.2% (2;44)	3.3% (9;251)
	Marina Del Rey Hospital	4.2% (3;23)	4.3% (1;37)		4.2% (1;35)		0.0% (0;55)	**9.4% (10;107) Worse**	4.2% (1;48)	1.8% (2;81)
	Memorial Hospital of Gardena	4.7% (3;25)	4.8% (2;51)	44.6% (2;4)	0.0% (0;47)		2.7% (4;114)	2.0% (4;204)	5.8% (1;15)	3.0% (4;93)
	Methodist Hospital of Southern California	4.7% (15;193)	6.9% (24;285)	18.7% (11;47)	3.5% (11;232)	19.2% (2;6)	2.0% (6;193)	**0.8% (6;368) Better**	3.2% (4;150)	**1.9% (13;277) Better**
	Mission Community Hospital – Panorama Campus	2.2% (1;14)	0.0% (0;30)	0.0% (0;4)	0.0% (0;25)		1.2% (1;49)	1.7% (5;153)	2.1% (1;26)	2.8% (6;99)
	Monterey Park Hospital	6.4% (2;15)	10.0% (2;40)	0.0% (0;4)	7.1% (2;36)		4.1% (2;52)	2.9% (2;94)	0.0% (0;20)	0.0% (0;92)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 5.0, and OSHPD hospital patient discharge data, 2015. See Technical Note for details about methodology.

‡ Data are reported for January – September 2015 due to coding changes from ICD-9-CM to ICD-10-CM/PCS, which began October 1, 2015. Comparisons of IMIs across years should be made with caution since previous years' results are based on 12 months of data, while this analysis is based on 9 months of data.

*****Hospital name changed from 2014 to 2015, see Technical Note for details.***

Note: Blank cells indicate that fewer than three conditions were treated.

Definition of Terms: Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. Total Deaths: number of patients that died in this hospital. Total Cases: number of patients with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.

OSHPD
Office of Statewide Health Planning and Development

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators (Conditions), 2015 (January – September) ‡

County	Hospital Name	Acute Myocardial Infarction	Acute Stroke				Gastro-Intestinal Hemorrhage	Heart Failure	Hip Fracture	Pneumonia
			Total	Hemorrhagic	Ischemic	Subarachnoid				
		NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.								
	STATEWIDE	5.9% (2,162;36,453)	8.7% (3,816;43,654)	20.9% (1,752;8,391)	5.0% (1,659;33,331)	21.0% (405;1,932)	2.3% (823;36,420)	2.9% (1,936;67,612)	2.0% (363;18,253)	3.3% (1,547;46,752)
Los Angeles (continued)	Northridge Hospital Medical Center	4.2% (15;281)	7.6% (18;167)	18.9% (9;36)	4.2% (8;125)	18.0% (1;6)	**0.0% (0;156) Better**	**1.0% (5;268) Better**	3.0% (3;72)	2.3% (7;188)
	Norwalk Community Hospital	0.0% (0;13)	3.0% (1;20)		0.0% (0;18)		0.0% (0;65)	1.3% (2;64)	0.0% (0;8)	0.9% (1;47)
	Olympia Medical Center	**0.0% (0;24) Better**	2.5% (1;38)		**0.0% (0;36) Better**		0.6% (1;103)	**0.3% (1;195) Better**	0.0% (0;21)	2.3% (5;104)
	PIH Hospital – Downey	5.7% (8;141)	7.0% (6;118)	17.5% (1;19)	2.9% (3;94)	40.3% (2;5)	1.7% (2;120)	**0.6% (3;254) Better**	0.0% (0;65)	**0.6% (2;159) Better**
	Pacific Alliance Medical Center	5.4% (1;11)	**0.0% (0;67) Better**	0.0% (0;9)	0.0% (0;58)		2.7% (2;68)	**0.5% (1;182) Better**	0.0% (0;23)	1.8% (2;95)
	Pacifica Hospital of the Valley		21.4% (1;10)		12.1% (1;10)		0.0% (0;40)	2.2% (1;49)	0.0% (0;8)	3.6% (3;68)
	Palmdale Regional Medical Center	3.1% (7;242)	7.1% (8;99)	19.4% (6;21)	3.1% (2;77)		2.4% (5;155)	2.7% (11;324)	0.0% (0;68)	5.7% (14;222)
	Pomona Valley Hospital Medical Center	4.4% (16;328)	7.9% (47;420)	14.7% (15;78)	5.9% (27;318)	12.6% (5;24)	1.3% (5;188)	3.3% (23;408)	0.0% (0;124)	2.7% (10;199)
	Presbyterian Intercommunity Hospital	5.2% (32;458)	**6.5% (29;346) Better**	16.3% (14;67)	3.5% (13;272)	15.6% (2;7)	0.7% (2;223)	**1.3% (10;461) Better**	0.0% (0;149)	**1.5% (7;293) Better**
	Providence Holy Cross Medical Center	2.8% (5;165)	6.5% (19;219)	16.7% (10;44)	3.7% (9;171)	0.0% (0;4)	2.0% (8;246)	**1.1% (10;452) Better**	2.0% (2;107)	**0.9% (5;297) Better**
	Providence Little Company of Mary Medical Center – San Pedro	9.5% (1;15)	6.3% (7;91)	**4.6% (1;15) Better**	6.4% (5;74)		3.9% (7;113)	**0.5% (1;130) Better**	3.7% (2;40)	**0.0% (0;103) Better**

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 5.0, and OSHPD hospital patient discharge data, 2015. See Technical Note for details about methodology.

‡ Data are reported for January – September 2015 due to coding changes from ICD-9-CM to ICD-10-CM/PCS, which began October 1, 2015. Comparisons of IMIs across years should be made with caution since previous years' results are based on 12 months of data, while this analysis is based on 9 months of data.

*****Hospital name changed from 2014 to 2015, see Technical Note for details.***

Note: Blank cells indicate that fewer than three conditions were treated.

Definition of Terms: Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. Total Deaths: number of patients that died in this hospital. Total Cases: number of patients with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.

OSHPD
Office of Statewide Health Planning and Development

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators (Conditions), 2015 (January – September) ‡

County	Hospital Name	Acute Myocardial Infarction	Acute Stroke				Gastro-Intestinal Hemorrhage	Heart Failure	Hip Fracture	Pneumonia
			Total	Hemorrhagic	Ischemic	Subarachnoid				
		NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.								
	STATEWIDE	5.9% (2,162;36,453)	8.7% (3,816;43,654)	20.9% (1,752;8,391)	5.0% (1,659;33,331)	21.0% (405;1,932)	2.3% (823;36,420)	2.9% (1,936;67,612)	2.0% (363;18,253)	3.3% (1,547;46,752)
Los Angeles (continued)	Providence Little Company of Mary Medical Center – Torrance	4.0% (15;344)	**4.3% (23;380) Better**	**8.7% (10;79) Better**	**2.3% (9;291) Better**	32.0% (4;10)	1.3% (6;281)	**1.0% (9;496) Better**	2.2% (3;111)	**1.8% (11;367) Better**
	Providence Saint John's Health Center**	4.1% (6;123)	5.8% (7;94)	9.9% (2;16)	4.0% (4;74)	14.4% (1;4)	2.8% (6;150)	2.3% (7;162)	3.0% (3;96)	4.2% (14;191)
	Providence Saint Joseph Medical Center	8.2% (15;230)	9.2% (41;374)	24.7% (17;65)	4.7% (18;280)	20.6% (6;29)	1.8% (5;230)	2.2% (17;406)	3.2% (2;130)	2.2% (11;260)
	Providence Tarzana Medical Center	**1.9% (3;122) Better**	8.1% (18;164)	14.3% (7;25)	5.8% (10;138)		0.8% (2;167)	**1.3% (8;273) Better**	**7.2% (4;84) Worse**	**1.2% (4;181) Better**
	Ronald Reagan UCLA Medical Center	7.3% (20;192)	**3.9% (31;397) Better**	**9.8% (16;106) Better**	**1.6% (7;243) Better**	**12.3% (8;48) Better**	1.9% (4;134)	2.7% (14;320)	3.5% (1;12)	4.6% (9;139)
	Saint Francis Medical Center	6.7% (14;158)	9.0% (36;295)	21.7% (18;54)	5.3% (18;238)	0.0% (0;3)	1.3% (3;192)	**1.3% (7;447) Better**	1.6% (1;45)	1.4% (3;162)
	Saint Mary Medical Center – Long Beach	4.7% (8;158)	7.1% (11;151)	22.1% (9;39)	1.2% (1;110)		4.7% (6;88)	1.3% (3;213)	0.0% (0;34)	1.8% (3;123)
	Saint Vincent Medical Center	**2.8% (7;150) Better**	**3.2% (3;99) Better**	13.2% (2;15)	1.0% (1;81)	0.0% (0;3)	0.0% (0;127)	2.4% (10;269)	1.8% (1;43)	1.8% (7;241)
	San Dimas Community Hospital	0.0% (0;11)	0.0% (0;24)		0.0% (0;24)		1.7% (1;34)	0.8% (1;58)	0.0% (0;28)	0.0% (0;31)
	San Gabriel Valley Medical Center	**1.0% (1;38) Better**	7.1% (8;112)	18.2% (4;21)	4.2% (4;87)	0.0% (0;4)	1.7% (3;119)	**0.9% (4;224) Better**	**9.1% (4;60) Worse**	3.3% (12;231)
	Santa Monica – UCLA Medical Center and Orthopedic Hospital	3.8% (8;128)	2.9% (2;90)	36.8% (2;4)	**0.0% (0;84) Better**		0.9% (2;143)	**1.0% (5;266) Better**	2.4% (2;129)	**0.5% (2;214) Better**

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 5.0, and OSHPD hospital patient discharge data, 2015. See Technical Note for details about methodology.

‡ Data are reported for January – September 2015 due to coding changes from ICD-9-CM to ICD-10-CM/PCS, which began October 1, 2015. Comparisons of IMIs across years should be made with caution since previous years' results are based on 12 months of data, while this analysis is based on 9 months of data.

*****Hospital name changed from 2014 to 2015, see Technical Note for details.***

Note: Blank cells indicate that fewer than three conditions were treated.

Definition of Terms: Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. Total Deaths: number of patients that died in this hospital. Total Cases: number of patients with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.

OSHPD
Office of Statewide Health Planning and Development

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators (Conditions), 2015 (January – September) ‡

County	Hospital Name	Acute Myocardial Infarction	Acute Stroke				Gastro-Intestinal Hemorrhage	Heart Failure	Hip Fracture	Pneumonia
			Total	Hemorrhagic	Ischemic	Subarachnoid				
		NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.								
	STATEWIDE	5.9% (2,162;36,453)	8.7% (3,816;43,654)	20.9% (1,752;8,391)	5.0% (1,659;33,331)	21.0% (405;1,932)	2.3% (823;36,420)	2.9% (1,936;67,612)	2.0% (363;18,253)	3.3% (1,547;46,752)
Los Angeles (continued)	Sherman Oaks Hospital	**1.2% (4;127) Better**	**0.0% (0;39) Better**	0.0% (0;3)	0.0% (0;36)		1.4% (3;97)	**0.2% (1;163) Better**	0.0% (0;27)	**0.6% (2;101) Better**
	Silver Lake Medical Center – Downtown Campus	0.0% (0;4)					0.0% (0;27)	0.0% (0;46)	0.0% (0;9)	3.8% (6;88)
	Southern California Hospital at Culver City	2.4% (2;64)	3.1% (2;75)	0.0% (0;3)	2.0% (2;72)		1.3% (3;113)	1.9% (8;243)	0.0% (0;32)	**1.1% (3;143) Better**
	Southern California Hospital at Hollywood	0.0% (0;12)	0.0% (0;7)		0.0% (0;7)		1.1% (1;53)	0.0% (0;76)	0.0% (0;4)	1.0% (1;82)
	Torrance Memorial Medical Center	5.7% (31;318)	8.0% (34;363)	16.5% (13;59)	4.9% (20;299)	83.4% (1;5)	1.5% (7;322)	1.9% (16;523)	1.7% (1;156)	2.4% (16;367)
	University of Southern California Verdugo Hills Hospital	3.1% (1;23)	7.1% (3;44)	0.0% (0;3)	4.3% (3;41)		0.0% (0;45)	0.7% (1;114)	2.5% (1;78)	2.0% (3;113)
	Valley Presbyterian Hospital	**0.0% (0;174) Better**	**0.0% (0;94) Better**	0.0% (0;15)	**0.0% (0;79) Better**		**0.0% (0;123) Better**	**0.0% (0;270) Better**	0.0% (0;28)	**0.0% (0;128) Better**
	West Hills Hospital and Medical Center	6.5% (13;195)	9.9% (12;125)	21.4% (5;17)	6.2% (7;107)		3.7% (8;158)	2.2% (7;181)	2.9% (2;90)	1.5% (4;144)
	White Memorial Medical Center	4.1% (11;189)	6.3% (11;190)	16.3% (4;39)	3.7% (7;146)	0.0% (0;5)	1.3% (5;176)	2.0% (7;264)	0.0% (0;30)	2.4% (5;151)
	Whittier Hospital Medical Center	6.5% (6;34)	4.3% (1;59)	24.3% (1;3)	0.0% (0;56)		**0.0% (0;119) Better**	3.5% (5;114)	6.9% (1;36)	4.4% (8;143)
Madera	Madera Community Hospital	0.0% (0;64)	6.7% (1;24)		4.3% (1;22)		2.4% (2;44)	3.3% (2;78)	0.0% (0;8)	1.5% (1;82)
Marin	Kaiser Foundation Hospital – San Rafael	3.7% (4;65)	9.5% (15;122)	13.8% (5;23)	8.7% (10;96)	0.0% (0;3)	2.9% (2;68)	1.8% (5;133)	1.1% (1;73)	1.8% (3;97)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 5.0, and OSHPD hospital patient discharge data, 2015. See Technical Note for details about methodology.

‡ Data are reported for January – September 2015 due to coding changes from ICD-9-CM to ICD-10-CM/PCS, which began October 1, 2015. Comparisons of IMIs across years should be made with caution since previous years' results are based on 12 months of data, while this analysis is based on 9 months of data.

*****Hospital name changed from 2014 to 2015, see Technical Note for details.***

Note: Blank cells indicate that fewer than three conditions were treated.

Definition of Terms: Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. Total Deaths: number of patients that died in this hospital. Total Cases: number of patients with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.

OSHPD
Office of Statewide Health Planning and Development

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators (Conditions), 2015 (January – September) ‡

County	Hospital Name	Acute Myocardial Infarction	Acute Stroke				Gastro-Intestinal Hemorrhage	Heart Failure	Hip Fracture	Pneumonia
			Total	Hemorrhagic	Ischemic	Subarachnoid				
		NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.								
	STATEWIDE	5.9% (2,162;36,453)	8.7% (3,816;43,654)	20.9% (1,752;8,391)	5.0% (1,659;33,331)	21.0% (405;1,932)	2.3% (823;36,420)	2.9% (1,936;67,612)	2.0% (363;18,253)	3.3% (1,547;46,752)
Marin (continued)	Marin General Hospital	4.9% (8;174)	7.6% (14;138)	19.2% (9;34)	4.1% (5;102)		**0.4% (1;159) Better**	**1.0% (3;163) Better**	2.3% (2;84)	1.5% (4;167)
	Novato Community Hospital	0.0% (0;6)	0.0% (0;34)		0.0% (0;32)		0.0% (0;27)	0.9% (1;50)	0.0% (0;31)	2.0% (1;39)
Mariposa	John C. Fremont Healthcare District							9.6% (1;24)		0.0% (0;11)
Mendocino	Frank R. Howard Memorial Hospital		0.0% (0;10)		0.0% (0;10)		0.0% (0;10)	0.0% (0;23)	6.1% (2;50)	0.0% (0;36)
	Mendocino Coast District Hospital	0.0% (0;3)	12.7% (2;15)		6.5% (1;14)		0.0% (0;15)	0.0% (0;27)	22.7% (1;9)	6.0% (2;35)
	Ukiah Valley Medical Center	8.9% (3;11)	12.0% (6;46)	0.0% (0;6)	10.2% (5;39)		0.0% (0;65)	2.8% (5;87)	2.9% (1;21)	1.3% (2;74)
Merced	Memorial Hospital Los Banos						0.0% (0;5)	0.7% (1;58)		1.3% (1;36)
	Mercy Medical Center – Merced	2.5% (1;23)	7.5% (4;101)		2.7% (2;99)		1.9% (2;138)	1.5% (7;263)	2.9% (2;68)	**1.7% (10;426) Better**
Modoc	Modoc Medical Center							0.0% (0;6)		30.6% (1;8)
Mono	Mammoth Hospital						0.0% (0;3)	0.0% (0;3)	0.0% (0;5)	0.0% (0;10)
Monterey	Community Hospital of The Monterey Peninsula	9.0% (12;129)	7.6% (19;176)	16.0% (9;33)	4.8% (7;136)	21.4% (3;7)	2.6% (5;158)	2.8% (9;199)	2.7% (2;109)	2.7% (8;232)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 5.0, and OSHPD hospital patient discharge data, 2015. See Technical Note for details about methodology.

‡ Data are reported for January – September 2015 due to coding changes from ICD-9-CM to ICD-10-CM/PCS, which began October 1, 2015. Comparisons of IMIs across years should be made with caution since previous years' results are based on 12 months of data, while this analysis is based on 9 months of data.

*****Hospital name changed from 2014 to 2015, see Technical Note for details.***

Note: Blank cells indicate that fewer than three conditions were treated.

Definition of Terms: Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. Total Deaths: number of patients that died in this hospital. Total Cases: number of patients with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.

OSHPD
Office of Statewide Health Planning and Development

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators (Conditions), 2015 (January – September) ‡

County	Hospital Name	Acute Myocardial Infarction	Acute Stroke				Gastro-Intestinal Hemorrhage	Heart Failure	Hip Fracture	Pneumonia
			Total	Hemorrhagic	Ischemic	Subarachnoid				
		NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.								
	STATEWIDE	5.9% (2,162;36,453)	8.7% (3,816;43,654)	20.9% (1,752;8,391)	5.0% (1,659;33,331)	21.0% (405;1,932)	2.3% (823;36,420)	2.9% (1,936;67,612)	2.0% (363;18,253)	3.3% (1,547;46,752)
Monterey (continued)	George L. Mee Memorial Hospital							0.0% (0;12)	0.0% (0;3)	0.0% (0;22)
	Natividad Medical Center	0.0% (0;4)	0.0% (0;31)	0.0% (0;6)	0.0% (0;25)		0.0% (0;51)	0.0% (0;59)	0.0% (0;28)	3.3% (1;71)
	Salinas Valley Memorial Hospital	4.8% (14;227)	6.3% (17;213)	16.1% (10;52)	3.1% (6;159)		0.5% (1;137)	1.9% (8;245)	2.5% (1;59)	3.9% (7;121)
Napa	Queen of the Valley Medical Center**	5.6% (7;91)	9.9% (10;97)	21.2% (5;19)	7.8% (5;77)		0.0% (0;54)	2.5% (8;167)	3.8% (3;59)	1.6% (2;105)
	Saint Helena Hospital	7.8% (10;125)	0.0% (0;17)		0.0% (0;17)		0.0% (0;42)	**0.0% (0;113) Better**	0.0% (0;31)	3.1% (2;39)
Nevada	Sierra Nevada Memorial Hospital	7.2% (4;25)	13.5% (5;58)	19.4% (1;4)	9.2% (4;54)		2.4% (2;89)	4.2% (9;132)	0.0% (0;63)	1.6% (3;159)
	Tahoe Forest Hospital						0.0% (0;12)	0.0% (0;18)	0.0% (0;24)	4.2% (1;54)
Orange	AHMC Anaheim Regional Medical Center	6.3% (13;181)	7.5% (7;100)	21.3% (4;18)	3.9% (3;79)	0.0% (0;3)	2.0% (3;143)	1.6% (6;310)	0.0% (0;44)	**0.3% (1;221) Better**
	Anaheim Global Medical Center**	3.1% (1;39)	0.0% (0;10)		0.0% (0;9)		6.1% (1;22)	0.0% (0;22)	0.0% (0;8)	6.2% (3;40)
	Chapman Global Medical Center**	7.2% (1;3)	0.0% (0;10)		0.0% (0;8)		0.0% (0;14)	0.0% (0;20)	0.0% (0;7)	0.0% (0;32)
	Fountain Valley Regional Hospital and Medical Center – Euclid	6.8% (26;265)	6.7% (27;267)	19.0% (17;57)	2.8% (8;204)	18.3% (2;6)	1.5% (5;197)	**1.1% (6;338) Better**	1.8% (1;71)	2.7% (13;303)
	Garden Grove Hospital and Medical Center	**0.9% (1;39) Better**	0.0% (0;38)	0.0% (0;8)	0.0% (0;30)		1.7% (1;69)	1.0% (2;93)	0.0% (0;30)	**0.0% (0;69) Better**

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 5.0, and OSHPD hospital patient discharge data, 2015. See Technical Note for details about methodology.

‡ Data are reported for January – September 2015 due to coding changes from ICD-9-CM to ICD-10-CM/PCS, which began October 1, 2015. Comparisons of IMIs across years should be made with caution since previous years' results are based on 12 months of data, while this analysis is based on 9 months of data.

*****Hospital name changed from 2014 to 2015, see Technical Note for details.***

Note: Blank cells indicate that fewer than three conditions were treated.

Definition of Terms: Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. Total Deaths: number of patients that died in this hospital. Total Cases: number of patients with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.

OSHPD
Office of Statewide Health Planning and Development

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators (Conditions), 2015 (January – September) ‡

County	Hospital Name	Acute Myocardial Infarction	Acute Stroke				Gastro-Intestinal Hemorrhage	Heart Failure	Hip Fracture	Pneumonia
			Total	Hemorrhagic	Ischemic	Subarachnoid				
		NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.								
	STATEWIDE	5.9% (2,162;36,453)	8.7% (3,816;43,654)	20.9% (1,752;8,391)	5.0% (1,659;33,331)	21.0% (405;1,932)	2.3% (823;36,420)	2.9% (1,936;67,612)	2.0% (363;18,253)	3.3% (1,547;46,752)
Orange (continued)	Hoag Memorial Hospital Presbyterian	5.3% (22;308)	**5.6% (36;499) Better**	15.2% (20;92)	**2.3% (10;379) Better**	18.2% (6;28)	**1.1% (7;413) Better**	**1.6% (15;373) Better**	1.1% (2;61)	2.1% (9;288)
	Hoag Orthopedic Institute								0.7% (1;193)	
	Huntington Beach Hospital	1.7% (1;32)	0.0% (0;20)		0.0% (0;20)		1.5% (1;32)	1.9% (4;83)	0.0% (0;15)	2.0% (2;63)
	Kaiser Foundation Hospital – Orange County – Anaheim	**2.9% (8;176) Better**	**5.2% (33;494) Better**	**12.5% (16;105) Better**	**2.4% (11;357) Better**	18.0% (6;32)	1.4% (6;299)	**1.3% (11;485) Better**	2.0% (2;148)	**0.8% (3;276) Better**
	La Palma Intercommunity Hospital	1.5% (1;30)	5.9% (1;25)		3.9% (1;24)		0.0% (0;42)	1.5% (3;94)	0.0% (0;21)	1.2% (2;89)
	Los Alamitos Medical Center	3.5% (5;102)	6.1% (17;241)	9.6% (4;45)	4.5% (12;183)	11.1% (1;13)	2.6% (6;143)	4.3% (11;211)	2.3% (1;111)	5.1% (18;219)
	Mission Hospital Laguna Beach	0.0% (0;3)	6.9% (1;4)		0.0% (0;3)		0.0% (0;31)	0.0% (0;23)	5.3% (1;30)	3.0% (1;34)
	Mission Hospital Regional Medical Center	4.9% (15;249)	**6.2% (29;286) Better**	**10.6% (8;60) Better**	4.3% (15;202)	16.3% (6;24)	1.4% (4;184)	**1.7% (13;279) Better**	1.2% (2;132)	**1.1% (4;205) Better**
	Orange Coast Memorial Medical Center	7.9% (13;132)	**0.0% (0;96) Better**	0.0% (0;23)	0.0% (0;70)	0.0% (0;3)	1.6% (4;160)	2.1% (9;260)	0.0% (0;74)	1.4% (4;186)
	Orange County Global Medical Center**	6.4% (6;122)	9.2% (14;146)	19.8% (7;44)	7.1% (7;96)	0.0% (0;6)	0.0% (0;58)	3.6% (4;111)	0.0% (0;21)	4.2% (5;77)
	Placentia Linda Hospital	5.9% (2;21)	7.6% (1;42)		5.3% (1;41)		1.1% (1;66)	0.7% (1;113)	7.9% (1;51)	1.6% (2;97)
	Saddleback Memorial Medical Center	4.1% (14;264)	7.3% (21;260)	25.5% (13;42)	2.8% (7;205)	10.4% (1;13)	**0.5% (2;190) Better**	1.9% (14;325)	1.4% (3;174)	**0.5% (2;232) Better**

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 5.0, and OSHPD hospital patient discharge data, 2015. See Technical Note for details about methodology.

‡ Data are reported for January – September 2015 due to coding changes from ICD-9-CM to ICD-10-CM/PCS, which began October 1, 2015. Comparisons of IMIs across years should be made with caution since previous years' results are based on 12 months of data, while this analysis is based on 9 months of data.

*****Hospital name changed from 2014 to 2015, see Technical Note for details.***

Note: Blank cells indicate that fewer than three conditions were treated.

Definition of Terms: Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. Total Deaths: number of patients that died in this hospital. Total Cases: number of patients with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.

OSHPD
Office of Statewide Health Planning and Development

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators (Conditions), 2015 (January – September) ‡

County	Hospital Name	Acute Myocardial Infarction	Acute Stroke				Gastro-Intestinal Hemorrhage	Heart Failure	Hip Fracture	Pneumonia
			Total	Hemorrhagic	Ischemic	Subarachnoid				
		NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.								
	STATEWIDE	5.9% (2,162;36,453)	8.7% (3,816;43,654)	20.9% (1,752;8,391)	5.0% (1,659;33,331)	21.0% (405;1,932)	2.3% (823;36,420)	2.9% (1,936;67,612)	2.0% (363;18,253)	3.3% (1,547;46,752)
Orange (continued)	Saint Joseph Hospital – Orange	**3.7% (13;253) Better**	**3.5% (13;309) Better**	**11.1% (9;66) Better**	**1.5% (4;232) Better**	0.0% (0;11)	2.3% (10;247)	2.5% (16;377)	0.0% (0;141)	2.8% (16;324)
	Saint Jude Medical Center	**2.9% (9;259) Better**	**3.9% (29;340) Better**	**9.4% (16;92) Better**	**2.0% (9;226) Better**	12.8% (4;22)	1.4% (3;190)	**1.6% (13;367) Better**	0.5% (1;164)	2.4% (9;237)
	South Coast Global Medical Center**	12.6% (2;10)	0.0% (0;7)		0.0% (0;7)		5.3% (1;28)	1.9% (1;33)	0.0% (0;12)	0.0% (0;45)
	UC Irvine Medical Center	6.4% (7;109)	**6.6% (52;409) Better**	**10.6% (20;139) Better**	5.7% (21;223)	20.0% (11;47)	1.1% (3;162)	2.5% (8;213)	2.8% (2;40)	1.4% (2;68)
	West Anaheim Medical Center	4.6% (8;107)	**3.6% (3;57) Better**	**0.0% (0;4) Better**	2.3% (2;52)		1.2% (2;81)	**0.9% (4;200) Better**	0.0% (0;33)	1.3% (3;105)
Placer	Kaiser Foundation Hospital – Roseville	5.5% (14;230)	10.8% (23;237)	29.9% (6;18)	5.9% (17;218)		2.5% (5;179)	1.8% (12;360)	3.2% (8;212)	**0.8% (2;149) Better**
	Sutter Auburn Faith Hospital	12.1% (5;14)	8.7% (3;62)		5.8% (3;58)		1.7% (2;78)	1.1% (2;85)	0.0% (0;73)	2.2% (4;104)
	Sutter Roseville Medical Center	**2.6% (7;274) Better**	7.1% (23;266)	16.5% (10;54)	4.1% (11;206)	18.5% (2;6)	2.9% (11;240)	2.2% (13;319)	3.2% (6;165)	2.1% (9;308)
Plumas	Eastern Plumas Hospital – Portola Campus									**43.3% (2;18) Worse**
	Plumas District Hospital						0.0% (0;9)	12.2% (1;13)		0.0% (0;15)
	Seneca District Hospital**							10.1% (1;17)		0.0% (0;19)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 5.0, and OSHPD hospital patient discharge data, 2015. See Technical Note for details about methodology.

‡ Data are reported for January – September 2015 due to coding changes from ICD-9-CM to ICD-10-CM/PCS, which began October 1, 2015. Comparisons of IMIs across years should be made with caution since previous years' results are based on 12 months of data, while this analysis is based on 9 months of data.

*****Hospital name changed from 2014 to 2015, see Technical Note for details.***

Note: Blank cells indicate that fewer than three conditions were treated.

Definition of Terms: Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. Total Deaths: number of patients that died in this hospital. Total Cases: number of patients with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.

OSHPD
Office of Statewide Health Planning and Development

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators (Conditions), 2015 (January – September) ‡

County	Hospital Name	Acute Myocardial Infarction	Acute Stroke				Gastro-Intestinal Hemorrhage	Heart Failure	Hip Fracture	Pneumonia
			Total	Hemorrhagic	Ischemic	Subarachnoid				
		NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.								
	STATEWIDE	5.9% (2,162;36,453)	8.7% (3,816;43,654)	20.9% (1,752;8,391)	5.0% (1,659;33,331)	21.0% (405;1,932)	2.3% (823;36,420)	2.9% (1,936;67,612)	2.0% (363;18,253)	3.3% (1,547;46,752)
Riverside	Corona Regional Medical Center – Main	3.4% (2;28)	4.0% (3;66)	15.3% (1;3)	1.9% (2;63)		4.3% (6;119)	**1.0% (5;233) Better**	0.0% (0;60)	2.6% (13;213)
	Desert Regional Medical Center	**9.0% (30;376) Worse**	9.2% (53;442)	20.8% (25;108)	5.1% (18;296)	27.2% (10;38)	2.4% (5;263)	3.0% (9;292)	2.1% (2;156)	2.6% (9;287)
	Eisenhower Medical Center	6.6% (18;342)	**5.7% (27;494) Better**	14.2% (13;76)	3.1% (12;408)	14.7% (2;10)	1.6% (7;367)	**1.0% (8;490) Better**	1.8% (4;239)	2.7% (19;458)
	Hemet Valley Medical Center	3.9% (7;70)	5.8% (3;76)	0.0% (0;4)	4.4% (3;72)		2.6% (8;201)	**1.4% (7;350) Better**	2.2% (1;105)	2.0% (9;269)
	John F. Kennedy Memorial Hospital	**10.7% (11;90) Worse**	9.5% (4;40)		5.4% (4;40)		**26.3% (3;26) Worse**	**0.5% (1;113) Better**	0.0% (0;55)	**0.6% (1;97) Better**
	Kaiser Foundation Hospital – Moreno Valley	**1.2% (1;41) Better**	6.0% (2;54)	40.3% (2;5)	0.0% (0;48)		0.0% (0;46)	2.0% (4;117)	0.0% (0;16)	2.8% (3;60)
	Kaiser Foundation Hospital – Riverside	5.3% (8;81)	7.2% (12;165)	9.9% (2;18)	4.6% (9;146)		2.0% (3;109)	2.3% (8;214)	3.8% (2;56)	2.0% (4;122)
	Loma Linda University Medical Center – Murrieta	7.1% (16;275)	8.3% (5;90)	32.0% (4;8)	1.9% (1;82)		1.2% (2;132)	1.8% (5;267)	0.0% (0;64)	2.1% (5;169)
	Menifee Valley Medical Center	6.5% (5;29)	0.0% (0;21)		0.0% (0;21)		1.5% (1;56)	**0.0% (0;114) Better**	8.5% (3;50)	3.1% (3;90)
	Palo Verde Hospital	0.0% (0;5)	0.0% (0;3)		0.0% (0;3)		7.0% (1;12)	7.8% (1;13)		3.4% (1;30)
	Parkview Community Hospital Medical Center	9.1% (4;16)	10.0% (2;52)		3.1% (1;50)		1.9% (2;103)	2.6% (5;161)	3.6% (1;38)	**0.9% (2;186) Better**

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 5.0, and OSHPD hospital patient discharge data, 2015. See Technical Note for details about methodology.

‡ Data are reported for January – September 2015 due to coding changes from ICD-9-CM to ICD-10-CM/PCS, which began October 1, 2015. Comparisons of IMIs across years should be made with caution since previous years' results are based on 12 months of data, while this analysis is based on 9 months of data.

*****Hospital name changed from 2014 to 2015, see Technical Note for details.***

Note: Blank cells indicate that fewer than three conditions were treated.

Definition of Terms: Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. Total Deaths: number of patients that died in this hospital. Total Cases: number of patients with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.

OSHPD
Office of Statewide Health Planning and Development

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators (Conditions), 2015 (January – September) ‡

County	Hospital Name	Acute Myocardial Infarction	Acute Stroke				Gastro-Intestinal Hemorrhage	Heart Failure	Hip Fracture	Pneumonia
			Total	Hemorrhagic	Ischemic	Subarachnoid				
		NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.								
	STATEWIDE	5.9% (2,162;36,453)	8.7% (3,816;43,654)	20.9% (1,752;8,391)	5.0% (1,659;33,331)	21.0% (405;1,932)	2.3% (823;36,420)	2.9% (1,936;67,612)	2.0% (363;18,253)	3.3% (1,547;46,752)
Riverside (continued)	Riverside Community Hospital	7.5% (33;509)	7.3% (27;326)	20.7% (16;60)	3.1% (9;263)	25.1% (2;3)	1.6% (7;316)	2.0% (13;534)	1.8% (1;104)	2.9% (15;376)
	Riverside County Regional Medical Center	4.8% (2;31)	7.2% (19;216)	19.6% (10;42)	3.0% (6;170)	24.3% (3;4)	3.6% (4;112)	2.1% (5;254)	0.0% (0;61)	2.3% (2;109)
	San Gorgonio Memorial Hospital	5.6% (6;31)	12.0% (1;12)		8.3% (1;11)		0.0% (0;80)	1.0% (2;151)	4.0% (1;30)	4.9% (7;175)
	Southwest Healthcare System – Murrieta	2.6% (2;42)	6.6% (15;224)	19.2% (8;37)	3.7% (7;181)	0.0% (0;6)	3.5% (5;149)	1.9% (7;231)	**9.6% (4;102) Worse**	2.0% (9;301)
	Temecula Valley Hospital	5.3% (19;331)	8.2% (13;149)	24.0% (7;34)	4.5% (6;111)	0.0% (0;4)	0.9% (1;101)	1.6% (6;200)	5.6% (4;74)	1.8% (5;169)
Sacramento	Kaiser Foundation Hospital – Sacramento	5.8% (12;115)	**4.3% (63;441) Better**	**8.0% (32;166) Better**	4.0% (20;224)	**12.2% (11;51) Better**	1.0% (3;165)	**1.0% (3;177) Better**	0.0% (0;104)	2.8% (5;108)
	Kaiser Foundation Hospital – South Sacramento	5.3% (11;179)	6.4% (15;204)	13.9% (6;35)	3.6% (8;167)		2.5% (4;115)	2.1% (14;398)	3.2% (3;98)	1.8% (3;107)
	Mercy General Hospital	4.7% (41;1,056)	**6.1% (22;280) Better**	14.0% (11;82)	3.3% (8;189)	23.4% (3;9)	1.5% (5;185)	2.4% (17;431)	1.7% (1;58)	2.1% (7;188)
	Mercy Hospital of Folsom**	1.7% (1;23)	5.1% (2;67)		1.9% (1;66)		2.9% (4;120)	1.3% (5;184)	0.0% (0;66)	2.4% (3;105)
	Mercy San Juan Hospital	4.8% (18;267)	8.3% (100;674)	18.0% (48;169)	5.6% (39;454)	19.1% (13;51)	**0.5% (2;226) Better**	**0.8% (10;673) Better**	1.9% (3;127)	**1.3% (5;256) Better**
	Methodist Hospital of Sacramento	3.8% (7;60)	7.8% (10;162)	40.9% (3;9)	3.6% (7;152)		2.0% (5;192)	**1.0% (5;260) Better**	6.9% (2;53)	1.6% (3;135)
	Sutter Medical Center, Sacramento**	6.1% (12;136)	8.5% (29;251)	18.3% (11;39)	5.0% (13;193)	24.3% (5;19)	2.0% (4;155)	3.0% (11;264)	3.7% (4;128)	2.0% (5;213)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 5.0, and OSHPD hospital patient discharge data, 2015. See Technical Note for details about methodology.

‡ Data are reported for January – September 2015 due to coding changes from ICD-9-CM to ICD-10-CM/PCS, which began October 1, 2015. Comparisons of IMIs across years should be made with caution since previous years' results are based on 12 months of data, while this analysis is based on 9 months of data.

*****Hospital name changed from 2014 to 2015, see Technical Note for details.***

Note: Blank cells indicate that fewer than three conditions were treated.

Definition of Terms: Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. Total Deaths: number of patients that died in this hospital. Total Cases: number of patients with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.

OSHPD
Office of Statewide Health Planning and Development

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators (Conditions), 2015 (January – September) ‡

County	Hospital Name	Acute Myocardial Infarction	Acute Stroke				Gastro-Intestinal Hemorrhage	Heart Failure	Hip Fracture	Pneumonia
			Total	Hemorrhagic	Ischemic	Subarachnoid				
		NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.								
	STATEWIDE	5.9% (2,162;36,453)	8.7% (3,816;43,654)	20.9% (1,752;8,391)	5.0% (1,659;33,331)	21.0% (405;1,932)	2.3% (823;36,420)	2.9% (1,936;67,612)	2.0% (363;18,253)	3.3% (1,547;46,752)
Sacramento (continued)	Sutter Memorial Hospital	7.0% (24;327)	5.6% (5;68)	19.1% (4;12)	1.5% (1;56)		0.0% (0;42)	2.9% (10;241)	0.0% (0;3)	1.5% (2;90)
	UC Davis Medical Center	**2.4% (7;278) Better**	7.1% (34;365)	**14.1% (16;78) Better**	5.4% (14;253)	15.2% (4;34)	1.8% (7;267)	**1.5% (12;561) Better**	0.0% (0;78)	1.6% (4;189)
San Benito	Hazel Hawkins Memorial Hospital	0.0% (0;15)	0.0% (0;19)		0.0% (0;19)		0.0% (0;34)	0.0% (0;55)	0.0% (0;21)	0.0% (0;75)
San Bernardino	Arrowhead Regional Medical Center	**0.0% (0;61) Better**	**0.0% (0;194) Better**	**0.0% (0;44) Better**	**0.0% (0;141) Better**	0.0% (0;9)	0.0% (0;92)	**0.0% (0;330) Better**	0.0% (0;32)	**0.0% (0;181) Better**
	Barstow Community Hospital	2.2% (1;36)	0.0% (0;35)		0.0% (0;35)		0.0% (0;32)	1.3% (2;102)	0.0% (0;13)	1.3% (1;61)
	Bear Valley Community Hospital							0.0% (0;7)		25.3% (1;10)
	Chino Valley Medical Center	1.9% (1;21)	**0.0% (0;24) Better**		**0.0% (0;23) Better**		0.7% (1;59)	0.6% (1;79)	0.0% (0;19)	0.0% (0;32)
	Colorado River Medical Center						0.0% (0;15)	0.0% (0;14)		0.0% (0;22)
	Community Hospital of San Bernardino	0.0% (0;15)	13.6% (3;37)		5.4% (2;36)		3.1% (3;53)	1.3% (1;95)	0.0% (0;13)	3.0% (3;107)
	Desert Valley Hospital	**2.6% (8;198) Better**	**1.2% (1;103) Better**	0.0% (0;3)	**0.8% (1;99) Better**		1.5% (3;131)	**0.8% (4;312) Better**	0.0% (0;34)	**0.6% (2;171) Better**
	Hi-Desert Medical Center	0.0% (0;5)	15.8% (2;18)		9.1% (2;18)		19.4% (1;12)	5.1% (3;106)	0.0% (0;22)	4.1% (3;93)
	Kaiser Foundation Hospital – Fontana	4.0% (20;382)	7.2% (46;515)	**13.9% (22;175) Better**	5.9% (17;300)	18.3% (7;40)	1.2% (4;281)	**1.8% (22;552) Better**	2.7% (3;125)	**1.4% (14;453) Better**

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 5.0, and OSHPD hospital patient discharge data, 2015. See Technical Note for details about methodology.

‡ Data are reported for January – September 2015 due to coding changes from ICD-9-CM to ICD-10-CM/PCS, which began October 1, 2015. Comparisons of IMIs across years should be made with caution since previous years' results are based on 12 months of data, while this analysis is based on 9 months of data.

*****Hospital name changed from 2014 to 2015, see Technical Note for details.***

Note: Blank cells indicate that fewer than three conditions were treated.

Definition of Terms: Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. Total Deaths: number of patients that died in this hospital. Total Cases: number of patients with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.

OSHPD
Office of Statewide Health Planning and Development

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators (Conditions), 2015 (January – September) ‡

County	Hospital Name	Acute Myocardial Infarction	Acute Stroke				Gastro-Intestinal Hemorrhage	Heart Failure	Hip Fracture	Pneumonia
			Total	Hemorrhagic	Ischemic	Subarachnoid				
		NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.								
	STATEWIDE	5.9% (2,162;36,453)	8.7% (3,816;43,654)	20.9% (1,752;8,391)	5.0% (1,659;33,331)	21.0% (405;1,932)	2.3% (823;36,420)	2.9% (1,936;67,612)	2.0% (363;18,253)	3.3% (1,547;46,752)
San Bernardino (continued)	Loma Linda University Medical Center	7.5% (26;379)	**6.6% (53;466) Better**	**14.1% (24;129) Better**	4.1% (22;306)	19.7% (7;31)	1.1% (3;186)	4.0% (26;495)	0.0% (0;55)	2.4% (6;179)
	Montclair Hospital Medical Center	3.1% (1;14)	0.0% (0;5)		0.0% (0;5)		3.4% (1;24)	3.4% (4;51)		0.0% (0;28)
	Mountains Community Hospital							0.0% (0;8)		0.0% (0;33)
	Redlands Community Hospital	1.5% (1;36)	5.2% (8;141)	16.6% (4;29)	2.6% (4;111)		1.2% (2;117)	0.9% (2;119)	1.2% (1;119)	1.9% (6;184)
	Saint Bernardine Medical Center	**9.1% (27;435) Worse**	8.9% (15;165)	23.9% (9;33)	2.6% (2;126)	25.5% (4;6)	2.0% (3;160)	1.3% (5;325)	2.9% (2;67)	2.5% (11;267)
	Saint Mary Medical Center – Apple Valley	**3.4% (12;336) Better**	4.5% (4;134)		2.6% (4;133)		1.7% (4;171)	**1.2% (7;482) Better**	0.0% (0;76)	1.9% (6;226)
	San Antonio Community Hospital	6.7% (24;395)	7.1% (27;301)	20.9% (17;64)	**2.4% (7;230) Better**	34.2% (3;7)	1.9% (5;226)	1.9% (14;431)	1.5% (2;132)	**1.9% (10;368) Better**
	Victor Valley Global Medical Center	4.7% (1;18)	0.0% (0;45)		0.0% (0;44)		2.1% (2;48)	2.3% (4;176)	0.0% (0;25)	1.2% (1;96)
San Diego	Alvarado Hospital	6.3% (10;96)	4.3% (4;63)	26.3% (4;12)	**0.0% (0;51) Better**		2.3% (3;53)	1.2% (2;118)	0.0% (0;32)	2.0% (2;75)
	Grossmont Hospital	4.3% (10;267)	9.5% (48;427)	20.0% (16;51)	5.4% (24;362)	28.1% (8;14)	1.3% (5;299)	**1.0% (8;625) Better**	3.3% (4;140)	2.5% (15;417)
	Kaiser Foundation Hospital – San Diego	**3.2% (8;147) Better**	6.5% (27;352)	**12.4% (10;78) Better**	4.5% (14;251)	15.3% (3;23)	2.1% (8;274)	2.6% (28;649)	1.8% (5;274)	3.4% (16;319)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 5.0, and OSHPD hospital patient discharge data, 2015. See Technical Note for details about methodology.

‡ Data are reported for January – September 2015 due to coding changes from ICD-9-CM to ICD-10-CM/PCS, which began October 1, 2015. Comparisons of IMIs across years should be made with caution since previous years' results are based on 12 months of data, while this analysis is based on 9 months of data.

*****Hospital name changed from 2014 to 2015, see Technical Note for details.***

Note: Blank cells indicate that fewer than three conditions were treated.

Definition of Terms: Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. Total Deaths: number of patients that died in this hospital. Total Cases: number of patients with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.

OSHPD
Office of Statewide Health Planning and Development

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators (Conditions), 2015 (January – September) ‡

County	Hospital Name	Acute Myocardial Infarction	Acute Stroke				Gastro-Intestinal Hemorrhage	Heart Failure	Hip Fracture	Pneumonia
			Total	Hemorrhagic	Ischemic	Subarachnoid				
		NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.								
	STATEWIDE	5.9% (2,162;36,453)	8.7% (3,816;43,654)	20.9% (1,752;8,391)	5.0% (1,659;33,331)	21.0% (405;1,932)	2.3% (823;36,420)	2.9% (1,936;67,612)	2.0% (363;18,253)	3.3% (1,547;46,752)
San Diego (continued)	Palomar Health Downtown Campus	3.5% (11;291)	9.9% (51;493)	21.5% (22;83)	6.5% (27;393)	14.5% (2;17)	1.9% (5;273)	3.0% (28;549)	4.8% (5;182)	2.6% (15;469)
	Paradise Valley Hospital	**2.2% (3;60) Better**	6.6% (4;53)	10.8% (1;7)	3.2% (2;45)		**0.0% (0;74) Better**	**0.3% (1;204) Better**	0.0% (0;17)	0.9% (1;68)
	Pomerado Hospital	5.7% (3;20)	9.3% (6;68)	36.3% (4;4)	3.1% (2;64)		2.0% (3;107)	2.2% (6;127)	2.8% (2;76)	**1.1% (3;135) Better**
	Scripps Green Hospital	4.5% (5;119)	**4.6% (12;133) Better**	**12.4% (9;49) Better**	**0.9% (1;76) Better**	18.1% (2;8)	**0.0% (0;100) Better**	1.3% (5;227)	0.0% (0;52)	1.5% (2;65)
	Scripps Memorial Hospital – Encinitas	**1.6% (3;139) Better**	**4.5% (11;173) Better**	**10.7% (5;31) Better**	2.7% (6;141)		2.0% (4;128)	**1.1% (5;249) Better**	1.6% (2;130)	1.8% (5;133)
	Scripps Memorial Hospital – La Jolla	7.6% (22;254)	8.8% (38;315)	19.3% (13;45)	5.8% (23;244)	10.6% (2;26)	2.2% (3;118)	2.7% (13;220)	1.5% (2;129)	1.8% (7;157)
	Scripps Mercy Hospital	4.9% (18;278)	**5.3% (27;383) Better**	16.3% (17;80)	**2.1% (9;298) Better**	14.2% (1;5)	1.1% (5;310)	**1.3% (16;696) Better**	1.7% (3;156)	**0.8% (6;460) Better**
	Sharp Chula Vista Medical Center	5.4% (13;186)	8.1% (23;250)	23.4% (9;36)	3.0% (8;202)	35.1% (6;12)	2.7% (8;176)	**1.4% (12;538) Better**	2.2% (2;76)	2.2% (10;242)
	Sharp Coronado Hospital and Healthcare Center	0.0% (0;3)	6.9% (1;16)		4.3% (1;15)		0.0% (0;31)	1.8% (1;30)	0.0% (0;17)	0.0% (0;24)
	Sharp Memorial Hospital	8.1% (19;253)	8.7% (34;334)	16.9% (12;65)	5.6% (18;251)	24.8% (4;18)	3.3% (10;259)	2.1% (18;514)	2.0% (3;183)	3.4% (23;405)
	Tri-City Medical Center – Oceanside	**8.8% (22;198) Worse**	7.2% (26;270)	15.7% (13;68)	4.2% (10;190)	28.6% (3;12)	3.5% (9;187)	2.1% (7;274)	1.6% (2;124)	4.0% (10;211)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 5.0, and OSHPD hospital patient discharge data, 2015. See Technical Note for details about methodology.

‡ Data are reported for January – September 2015 due to coding changes from ICD-9-CM to ICD-10-CM/PCS, which began October 1, 2015. Comparisons of IMIs across years should be made with caution since previous years' results are based on 12 months of data, while this analysis is based on 9 months of data.

*****Hospital name changed from 2014 to 2015, see Technical Note for details.***

Note: Blank cells indicate that fewer than three conditions were treated.

Definition of Terms: Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. Total Deaths: number of patients that died in this hospital. Total Cases: number of patients with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.

OSHPD
Office of Statewide Health Planning and Development

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators (Conditions), 2015 (January – September) ‡

County	Hospital Name	Acute Myocardial Infarction	Acute Stroke				Gastro-Intestinal Hemorrhage	Heart Failure	Hip Fracture	Pneumonia
			Total	Hemorrhagic	Ischemic	Subarachnoid				
		NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.								
	STATEWIDE	5.9% (2,162;36,453)	8.7% (3,816;43,654)	20.9% (1,752;8,391)	5.0% (1,659;33,331)	21.0% (405;1,932)	2.3% (823;36,420)	2.9% (1,936;67,612)	2.0% (363;18,253)	3.3% (1,547;46,752)
San Diego (continued)	UC San Diego Health System – Hillcrest Medical Center	6.9% (13;233)	**6.0% (31;278) Better**	**13.7% (16;71) Better**	4.0% (11;182)	12.3% (4;25)	1.0% (4;230)	1.9% (14;659)	1.2% (1;79)	2.1% (6;236)
San Francisco	California Pacific Medical Center – Davies Campus	6.4% (1;5)	7.5% (38;333)	**10.6% (9;70) Better**	5.8% (24;235)	17.2% (5;28)	7.6% (1;30)	2.2% (2;59)	0.0% (0;12)	2.0% (1;51)
	California Pacific Medical Center – Pacific Campus	4.4% (11;197)	7.8% (27;228)	14.1% (9;49)	5.4% (17;175)	21.5% (1;4)	**0.2% (1;236) Better**	**1.6% (11;349) Better**	3.2% (5;87)	2.4% (8;211)
	California Pacific Medical Center – St. Luke's Campus	0.0% (0;10)	5.1% (1;23)		2.9% (1;23)		2.1% (1;45)	2.4% (5;117)	0.0% (0;11)	0.0% (0;78)
	Chinese Hospital	4.7% (1;11)	10.6% (1;18)		0.0% (0;16)		1.9% (1;49)	1.1% (3;103)	0.0% (0;17)	1.8% (3;133)
	Kaiser Foundation Hospital – San Francisco	**3.4% (12;395) Better**	**4.5% (13;171) Better**	**9.3% (5;33) Better**	3.2% (7;134)	7.1% (1;4)	2.2% (3;106)	**1.0% (5;262) Better**	2.4% (1;54)	2.4% (3;56)
	Laguna Honda Hospital and Rehabilitation Center									4.9% (1;10)
	Saint Francis Memorial Hospital	4.9% (2;17)	7.4% (12;100)	23.4% (10;22)	1.9% (2;78)		1.4% (2;95)	0.9% (1;122)	0.0% (0;26)	2.2% (3;126)
	Saint Mary's Medical Center, San Francisco	6.3% (7;88)	8.1% (6;69)	23.9% (3;11)	4.0% (3;58)		1.2% (3;90)	2.5% (9;175)	5.2% (1;31)	2.3% (3;73)
	San Francisco General Hospital	7.3% (14;116)	9.7% (27;189)	23.5% (16;56)	5.7% (8;124)	19.6% (3;9)	1.4% (2;150)	4.1% (11;337)	0.0% (0;54)	**0.6% (1;186) Better**
	UC San Francisco Medical Center	6.7% (8;87)	8.5% (50;293)	15.0% (18;79)	5.4% (14;147)	25.1% (18;67)	1.4% (5;177)	2.6% (14;309)	0.0% (0;58)	1.9% (5;186)
San Joaquin	Dameron Hospital	4.8% (9;240)	7.0% (5;72)	38.2% (3;3)	2.2% (2;69)		2.4% (4;117)	2.8% (9;213)	14.4% (2;37)	5.7% (11;143)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 5.0, and OSHPD hospital patient discharge data, 2015. See Technical Note for details about methodology.

‡ Data are reported for January – September 2015 due to coding changes from ICD-9-CM to ICD-10-CM/PCS, which began October 1, 2015. Comparisons of IMIs across years should be made with caution since previous years' results are based on 12 months of data, while this analysis is based on 9 months of data.

*****Hospital name changed from 2014 to 2015, see Technical Note for details.***

Note: Blank cells indicate that fewer than three conditions were treated.

Definition of Terms: Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. Total Deaths: number of patients that died in this hospital. Total Cases: number of patients with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.

OSHPD
Office of Statewide Health Planning and Development

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators (Conditions), 2015 (January – September) ‡

County	Hospital Name	Acute Myocardial Infarction	Acute Stroke				Gastro-Intestinal Hemorrhage	Heart Failure	Hip Fracture	Pneumonia
			Total	Hemorrhagic	Ischemic	Subarachnoid				
		NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.								
	STATEWIDE	5.9% (2,162;36,453)	8.7% (3,816;43,654)	20.9% (1,752;8,391)	5.0% (1,659;33,331)	21.0% (405;1,932)	2.3% (823;36,420)	2.9% (1,936;67,612)	2.0% (363;18,253)	3.3% (1,547;46,752)
San Joaquin (continued)	Doctors Hospital of Manteca	0.0% (0;21)	4.4% (1;39)		2.5% (1;39)		2.4% (1;77)	2.3% (3;119)	7.2% (1;32)	3.0% (3;115)
	Kaiser Foundation Hospital – Manteca	**0.6% (1;116) Better**	6.2% (6;75)	15.4% (2;9)	3.5% (3;64)		1.0% (2;104)	**1.2% (6;220) Better**	0.0% (0;93)	2.0% (3;93)
	Lodi Memorial Hospital	3.0% (3;49)	10.3% (7;81)	22.6% (3;7)	6.4% (4;74)		3.1% (3;109)	2.8% (8;251)	3.0% (1;51)	1.8% (4;256)
	Saint Joseph's Medical Center of Stockton	**0.0% (0;310) Better**	**0.0% (0;180) Better**	**0.0% (0;28) Better**	**0.0% (0;151) Better**		**0.0% (0;202) Better**	**0.0% (0;358) Better**	0.0% (0;78)	**0.0% (0;182) Better**
	San Joaquin General Hospital	4.9% (1;11)	6.4% (4;84)	11.8% (2;10)	5.4% (2;74)		3.5% (3;72)	2.3% (6;179)	0.0% (0;15)	2.4% (3;66)
	Sutter Tracy Community Hospital	2.1% (1;14)	5.6% (1;34)		3.5% (1;33)		1.3% (1;67)	**0.5% (1;106) Better**	0.0% (0;13)	0.0% (0;82)
San Luis Obispo	French Hospital Medical Center	5.1% (6;184)	9.8% (2;40)		6.3% (2;39)		1.4% (1;68)	1.9% (4;119)	2.8% (1;43)	0.8% (1;71)
	Marian Regional Medical Center – Arroyo Grande	0.0% (0;9)	5.8% (2;44)		2.3% (1;42)		4.5% (2;55)	1.2% (1;54)	3.7% (1;32)	3.5% (4;68)
	Sierra Vista Regional Medical Center	**24.4% (4;5) Worse**	9.8% (10;98)	23.1% (6;39)	4.9% (3;58)		0.0% (0;47)	2.8% (1;34)	4.4% (2;41)	0.0% (0;78)
	Twin Cities Community Hospital	5.0% (2;12)	14.9% (8;77)		8.7% (7;75)		0.0% (0;64)	2.0% (2;86)	**19.0% (4;64) Worse**	5.9% (9;131)
San Mateo	Kaiser Foundation Hospital – Redwood City	10.0% (7;60)	**5.1% (47;349) Better**	**10.3% (27;135) Better**	5.5% (13;150)	**10.6% (7;64) Better**	2.0% (2;66)	3.5% (11;120)	2.4% (2;56)	2.2% (4;88)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 5.0, and OSHPD hospital patient discharge data, 2015. See Technical Note for details about methodology.

‡ Data are reported for January – September 2015 due to coding changes from ICD-9-CM to ICD-10-CM/PCS, which began October 1, 2015. Comparisons of IMIs across years should be made with caution since previous years' results are based on 12 months of data, while this analysis is based on 9 months of data.

*****Hospital name changed from 2014 to 2015, see Technical Note for details.***

Note: Blank cells indicate that fewer than three conditions were treated.

Definition of Terms: Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. Total Deaths: number of patients that died in this hospital. Total Cases: number of patients with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.

OSHPD
Office of Statewide Health Planning and Development

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators (Conditions), 2015 (January – September) ‡

County	Hospital Name	Acute Myocardial Infarction	Acute Stroke				Gastro-Intestinal Hemorrhage	Heart Failure	Hip Fracture	Pneumonia
			Total	Hemorrhagic	Ischemic	Subarachnoid				
		NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.								
	STATEWIDE	5.9% (2,162;36,453)	8.7% (3,816;43,654)	20.9% (1,752;8,391)	5.0% (1,659;33,331)	21.0% (405;1,932)	2.3% (823;36,420)	2.9% (1,936;67,612)	2.0% (363;18,253)	3.3% (1,547;46,752)
San Mateo (continued)	Kaiser Foundation Hospital – South San Francisco	3.0% (3;40)	6.1% (11;104)	**10.4% (4;21) Better**	4.6% (7;82)		1.7% (3;91)	1.4% (6;210)	0.0% (0;64)	1.1% (2;83)
	Peninsula Medical Center	6.3% (12;198)	**5.7% (14;188) Better**	20.2% (9;36)	**2.2% (5;149) Better**	0.0% (0;3)	1.1% (5;205)	**1.0% (7;271) Better**	2.0% (3;130)	**0.6% (2;181) Better**
	San Mateo Medical Center	0.0% (0;5)	11.8% (2;24)	19.3% (1;4)	20.2% (1;19)		0.0% (0;60)	1.4% (1;96)	56.4% (1;5)	2.8% (2;105)
	Sequoia Hospital	0.0% (0;41)	3.6% (1;42)	23.7% (1;4)	0.0% (0;38)		0.0% (0;36)	0.8% (1;107)	0.0% (0;48)	**0.0% (0;82) Better**
	Seton Medical Center	4.8% (8;133)	8.2% (13;122)	31.4% (11;28)	**1.6% (2;93) Better**		2.1% (3;138)	**0.8% (5;276) Better**	7.7% (2;47)	2.6% (9;167)
Santa Barbara	Goleta Valley Cottage Hospital							0.0% (0;8)	0.0% (0;55)	0.0% (0;23)
	Lompoc Valley Medical Center	0.0% (0;9)	5.0% (1;19)		2.8% (1;19)		8.5% (3;41)	3.5% (2;72)	0.0% (0;16)	5.9% (5;78)
	Marian Regional Medical Center	5.2% (6;108)	**6.0% (19;165) Better**	13.5% (10;48)	4.3% (9;114)	0.0% (0;3)	4.0% (9;155)	3.0% (12;184)	2.1% (2;76)	2.5% (3;83)
	Santa Barbara Cottage Hospital	4.2% (6;193)	8.5% (50;422)	19.0% (17;81)	5.4% (23;281)	18.7% (10;60)	1.8% (5;220)	2.8% (16;249)	3.8% (3;123)	2.6% (8;212)
	Santa Ynez Valley Cottage Hospital							8.8% (1;10)		5.5% (1;18)
Santa Clara	El Camino Hospital	7.3% (12;174)	7.7% (26;249)	15.4% (12;46)	5.6% (14;200)	0.0% (0;3)	2.2% (5;212)	4.2% (16;265)	1.3% (2;160)	4.1% (16;299)
	Good Samaritan Hospital – San Jose	7.7% (13;173)	7.2% (21;282)	15.6% (8;48)	4.8% (10;211)	13.5% (3;23)	1.0% (3;198)	2.4% (12;282)	3.4% (3;97)	2.5% (8;200)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 5.0, and OSHPD hospital patient discharge data, 2015. See Technical Note for details about methodology.

‡ Data are reported for January – September 2015 due to coding changes from ICD-9-CM to ICD-10-CM/PCS, which began October 1, 2015. Comparisons of IMIs across years should be made with caution since previous years' results are based on 12 months of data, while this analysis is based on 9 months of data.

*****Hospital name changed from 2014 to 2015, see Technical Note for details.***

Note: Blank cells indicate that fewer than three conditions were treated.

Definition of Terms: Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. Total Deaths: number of patients that died in this hospital. Total Cases: number of patients with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.

OSHPD
Office of Statewide Health Planning and Development

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators (Conditions), 2015 (January – September) ‡

County	Hospital Name	Acute Myocardial Infarction	Acute Stroke				Gastro-Intestinal Hemorrhage	Heart Failure	Hip Fracture	Pneumonia
			Total	Hemorrhagic	Ischemic	Subarachnoid				
		NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.								
	STATEWIDE	5.9% (2,162;36,453)	8.7% (3,816;43,654)	20.9% (1,752;8,391)	5.0% (1,659;33,331)	21.0% (405;1,932)	2.3% (823;36,420)	2.9% (1,936;67,612)	2.0% (363;18,253)	3.3% (1,547;46,752)
Santa Clara (continued)	Kaiser Foundation Hospital – San Jose	3.4% (7;111)	5.8% (12;179)	**6.8% (2;19) Better**	4.0% (9;158)		1.5% (4;149)	2.3% (16;295)	3.7% (3;76)	1.1% (2;100)
	Kaiser Foundation Hospital – Santa Clara	**3.1% (16;443) Better**	7.1% (20;190)	13.3% (8;39)	5.1% (12;150)		1.7% (5;185)	2.0% (21;562)	2.0% (3;108)	3.0% (7;111)
	O'Connor Hospital	5.1% (10;151)	5.2% (8;125)	12.6% (4;29)	3.0% (4;96)		3.3% (9;158)	**1.3% (6;256) Better**	5.0% (2;80)	2.9% (6;166)
	Regional Medical of San Jose	5.4% (13;191)	9.1% (56;398)	17.4% (22;105)	5.7% (24;268)	31.8% (10;25)	1.9% (5;226)	1.9% (11;426)	0.0% (0;76)	2.3% (13;429)
	Saint Louise Regional Hospital	18.1% (2;5)	10.5% (3;31)	39.9% (2;4)	3.8% (1;26)		0.0% (0;21)	1.4% (2;83)	0.0% (0;22)	4.4% (5;103)
	Santa Clara Valley Medical Center	4.7% (5;124)	6.5% (13;231)	12.7% (6;50)	4.8% (7;179)		1.2% (3;252)	1.2% (4;527)	6.5% (2;29)	2.8% (5;174)
	Stanford Hospital	6.4% (13;169)	**5.8% (37;360) Better**	**12.8% (16;84) Better**	3.7% (17;231)	12.4% (4;45)	1.6% (5;230)	2.2% (13;366)	3.4% (4;105)	1.5% (4;210)
Santa Cruz	Dominican Hospital	6.2% (10;193)	8.3% (14;175)	21.0% (8;29)	3.6% (4;142)	32.2% (2;4)	**6.7% (10;170) Worse**	3.0% (10;299)	4.3% (2;85)	4.0% (7;190)
	Watsonville Community Hospital	15.9% (1;8)	0.0% (0;38)	0.0% (0;3)	0.0% (0;35)		0.0% (0;57)	4.8% (6;101)	5.4% (1;32)	4.8% (6;91)
Shasta	Mayers Memorial Hospital		13.1% (1;5)		8.2% (1;4)		0.0% (0;3)	23.3% (1;14)		5.6% (1;24)
	Mercy Medical Center – Redding	6.4% (20;247)	7.7% (31;225)	18.4% (21;69)	4.6% (10;155)		1.4% (4;156)	2.9% (18;324)	4.0% (5;128)	**1.5% (8;248) Better**

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 5.0, and OSHPD hospital patient discharge data, 2015. See Technical Note for details about methodology.

‡ Data are reported for January – September 2015 due to coding changes from ICD-9-CM to ICD-10-CM/PCS, which began October 1, 2015. Comparisons of IMIs across years should be made with caution since previous years' results are based on 12 months of data, while this analysis is based on 9 months of data.

*****Hospital name changed from 2014 to 2015, see Technical Note for details.***

Note: Blank cells indicate that fewer than three conditions were treated.

Definition of Terms: Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. Total Deaths: number of patients that died in this hospital. Total Cases: number of patients with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.

OSHPD
Office of Statewide Health Planning and Development

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators (Conditions), 2015 (January – September) ‡

County	Hospital Name	Acute Myocardial Infarction	Acute Stroke				Gastro-Intestinal Hemorrhage	Heart Failure	Hip Fracture	Pneumonia
			Total	Hemorrhagic	Ischemic	Subarachnoid				
		NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.								
	STATEWIDE	5.9% (2,162;36,453)	8.7% (3,816;43,654)	20.9% (1,752;8,391)	5.0% (1,659;33,331)	21.0% (405;1,932)	2.3% (823;36,420)	2.9% (1,936;67,612)	2.0% (363;18,253)	3.3% (1,547;46,752)
Shasta (continued)	Shasta Regional Medical Center	4.2% (9;158)	12.5% (11;123)	**96.5% (3;5) Worse**	5.7% (8;118)		1.9% (5;146)	**0.5% (3;254) Better**	0.0% (0;48)	1.9% (5;143)
Siskiyou	Fairchild Medical Center	0.0% (0;6)	7.6% (1;24)		4.9% (1;23)		10.5% (1;24)	5.4% (1;21)	6.6% (1;20)	2.0% (2;75)
	Mercy Medical Center Mt. Shasta		26.2% (2;13)		14.8% (2;13)		7.4% (1;18)	0.0% (0;16)	**15.9% (2;19) Worse**	0.0% (0;36)
Solano	Kaiser Foundation Hospital – Rehabilitation Center Vallejo	7.8% (15;165)	**4.4% (12;203) Better**	**11.1% (7;38) Better**	2.4% (5;164)		2.1% (5;162)	1.7% (9;259)	1.9% (2;85)	1.6% (3;95)
	Kaiser Foundation Hospital – Vacaville	4.7% (2;41)	**5.5% (15;182) Better**	**7.4% (5;47) Better**	5.0% (9;129)	41.0% (1;6)	1.6% (3;92)	3.0% (10;145)	2.2% (1;53)	1.4% (3;108)
	North Bay Medical Center	5.8% (16;181)	11.2% (20;130)	**33.4% (15;27) Worse**	4.1% (5;103)		**0.7% (2;128) Better**	2.3% (9;273)	1.4% (1;51)	3.5% (11;194)
	Sutter Solano Medical Center	3.7% (1;13)	5.8% (2;44)	0.0% (0;3)	5.0% (2;41)		1.4% (1;61)	**0.0% (0;135) Better**	5.6% (1;22)	2.2% (2;102)
Sonoma	Healdsburg District Hospital	13.2% (1;3)	14.1% (1;11)		7.9% (1;11)		0.0% (0;4)	9.0% (2;22)	0.0% (0;17)	4.8% (1;36)
	Kaiser Foundation Hospital – Santa Rosa	2.6% (2;47)	9.0% (16;124)	17.4% (4;21)	5.9% (11;100)	12.5% (1;3)	1.4% (3;130)	2.0% (8;180)	3.8% (4;101)	**0.9% (2;94) Better**
	Petaluma Valley Hospital	3.7% (2;25)	9.6% (2;31)		4.4% (1;30)		5.9% (4;43)	0.0% (0;71)	0.0% (0;23)	1.8% (1;62)
	Santa Rosa Memorial Hospital – Montgomery	4.4% (13;262)	8.9% (24;192)	15.2% (11;54)	8.5% (13;136)		1.8% (5;204)	2.4% (9;220)	1.3% (1;116)	2.9% (7;191)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 5.0, and OSHPD hospital patient discharge data, 2015. See Technical Note for details about methodology.

‡ Data are reported for January – September 2015 due to coding changes from ICD-9-CM to ICD-10-CM/PCS, which began October 1, 2015. Comparisons of IMIs across years should be made with caution since previous years' results are based on 12 months of data, while this analysis is based on 9 months of data.

*****Hospital name changed from 2014 to 2015, see Technical Note for details.***

Note: Blank cells indicate that fewer than three conditions were treated.

Definition of Terms: Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. Total Deaths: number of patients that died in this hospital. Total Cases: number of patients with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.

OSHPD
Office of Statewide Health Planning and Development

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators (Conditions), 2015 (January – September) ‡

County	Hospital Name	Acute Myocardial Infarction	Acute Stroke				Gastro-Intestinal Hemorrhage	Heart Failure	Hip Fracture	Pneumonia
			Total	Hemorrhagic	Ischemic	Subarachnoid				
		NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.								
	STATEWIDE	5.9% (2,162;36,453)	8.7% (3,816;43,654)	20.9% (1,752;8,391)	5.0% (1,659;33,331)	21.0% (405;1,932)	2.3% (823;36,420)	2.9% (1,936;67,612)	2.0% (363;18,253)	3.3% (1,547;46,752)
Sonoma (continued)	Sonoma Valley Hospital	0.0% (0;4)	0.0% (0;15)	0.0% (0;3)	0.0% (0;12)		5.2% (2;19)	2.9% (2;33)	0.0% (0;23)	0.0% (0;14)
	Sutter Santa Rosa Regional Hospital	4.0% (3;97)	7.3% (5;55)	13.4% (2;5)	4.6% (2;49)		1.6% (2;73)	3.4% (7;108)	1.7% (1;43)	1.9% (2;63)
Stanislaus	Doctors Medical Center	**9.2% (24;308) Worse**	8.5% (40;392)	20.6% (26;100)	4.8% (12;286)	17.2% (2;6)	3.2% (10;265)	2.1% (8;445)	0.7% (1;138)	4.5% (15;397)
	Emanuel Medical Center	5.5% (14;274)	6.3% (4;58)	20.1% (2;3)	2.9% (2;55)		1.3% (1;84)	1.3% (4;300)	0.0% (0;41)	3.6% (15;335)
	Memorial Hospital Medical Center – Modesto	4.4% (15;294)	**3.7% (15;338) Better**	**11.1% (9;69) Better**	**1.7% (6;264) Better**	0.0% (0;5)	1.1% (5;307)	1.9% (14;429)	1.4% (2;180)	2.0% (9;267)
	Oak Valley District Hospital	0.0% (0;4)	12.0% (1;13)		6.8% (1;13)		0.0% (0;19)	4.6% (1;36)	0.0% (0;7)	7.4% (2;44)
Tehama	Saint Elizabeth Community Hospital	6.6% (1;5)	2.7% (1;37)	0.0% (0;3)	0.0% (0;33)		0.0% (0;59)	2.6% (3;86)	8.2% (1;32)	4.4% (5;84)
Trinity	Trinity Hospital	0.0% (0;3)					0.0% (0;4)	0.0% (0;10)		0.0% (0;30)
Tulare	Kaweah Delta Medical Center	5.6% (23;416)	9.8% (33;318)	20.3% (16;51)	6.8% (15;262)	23.1% (2;5)	3.4% (12;206)	2.3% (24;564)	2.5% (7;159)	**1.9% (16;510) Better**
	Sierra View Medical Center	9.8% (8;60)	4.7% (3;82)		2.7% (3;81)		0.0% (0;77)	3.7% (7;130)	2.4% (1;56)	**7.0% (20;227) Worse**
	Tulare Regional Medical Center	2.8% (1;18)	16.2% (3;14)		7.7% (1;11)		1.6% (4;151)	1.7% (2;64)	13.2% (1;22)	1.8% (4;142)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 5.0, and OSHPD hospital patient discharge data, 2015. See Technical Note for details about methodology.

‡ Data are reported for January – September 2015 due to coding changes from ICD-9-CM to ICD-10-CM/PCS, which began October 1, 2015. Comparisons of IMIs across years should be made with caution since previous years' results are based on 12 months of data, while this analysis is based on 9 months of data.

*****Hospital name changed from 2014 to 2015, see Technical Note for details.***

Note: Blank cells indicate that fewer than three conditions were treated.

Definition of Terms: Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. Total Deaths: number of patients that died in this hospital. Total Cases: number of patients with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.

OSHPD
Office of Statewide Health Planning and Development

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators (Conditions), 2015 (January – September) ‡

County	Hospital Name	Acute Myocardial Infarction	Acute Stroke				Gastro-Intestinal Hemorrhage	Heart Failure	Hip Fracture	Pneumonia
			Total	Hemorrhagic	Ischemic	Subarachnoid				
		NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.								
	STATEWIDE	5.9% (2,162;36,453)	8.7% (3,816;43,654)	20.9% (1,752;8,391)	5.0% (1,659;33,331)	21.0% (405;1,932)	2.3% (823;36,420)	2.9% (1,936;67,612)	2.0% (363;18,253)	3.3% (1,547;46,752)
Tuolumne	Sonora Regional Medical Center – Greenley	0.0% (0;16)	6.6% (5;70)		3.8% (5;69)		3.4% (4;80)	1.6% (3;114)	5.7% (3;40)	2.0% (4;127)
Ventura	Community Memorial Hospital – San Buenaventura	5.6% (5;143)	**4.1% (5;124) Better**	5.7% (1;24)	2.4% (3;98)		0.4% (1;148)	2.8% (7;147)	0.0% (0;103)	**0.5% (1;126) Better**
	Los Robles Hospital and Medical Center	5.4% (17;231)	8.0% (20;209)	24.2% (9;38)	2.9% (6;158)	27.7% (5;13)	1.6% (4;174)	2.1% (14;290)	0.0% (0;117)	1.9% (6;183)
	Ojai Valley Community Hospital	9.9% (1;3)	7.2% (1;20)		0.0% (0;18)		0.0% (0;23)	4.0% (1;9)	**10.5% (3;20) Worse**	0.0% (0;21)
	Saint John's Pleasant Valley Hospital	20.9% (3;8)	8.1% (3;62)	21.4% (1;4)	4.4% (2;58)		2.8% (2;65)	2.4% (6;135)	8.0% (3;45)	3.7% (6;129)
	Saint John's Regional Medical Center	4.7% (13;217)	**5.3% (14;150) Better**	**11.1% (7;39) Better**	3.2% (4;105)	17.3% (3;6)	1.0% (2;146)	1.6% (9;306)	0.0% (0;55)	3.0% (14;252)
	Simi Valley Hospital and Healthcare Services – Sycamore	6.0% (4;50)	10.1% (10;99)	**42.1% (7;15) Worse**	3.4% (3;83)		0.0% (0;102)	1.6% (5;117)	0.0% (0;52)	2.3% (5;122)
	Ventura County Medical Center	0.0% (0;29)	9.2% (10;78)	30.2% (8;22)	1.5% (1;55)		0.9% (1;87)	3.3% (4;109)	0.0% (0;43)	4.7% (6;91)
Yolo	Sutter Davis Hospital	9.2% (1;5)	3.5% (2;36)	9.8% (1;4)	2.4% (1;31)		0.0% (0;55)	1.4% (1;49)	0.0% (0;44)	0.0% (0;50)
	Woodland Memorial Hospital	10.6% (2;12)	7.0% (2;40)	0.0% (0;3)	6.5% (2;37)		1.8% (1;41)	3.3% (4;57)	9.7% (1;26)	**9.2% (6;68) Worse**
Yuba	Rideout Memorial Hospital	5.9% (13;209)	9.0% (16;185)	19.9% (4;25)	5.2% (10;150)	27.0% (2;10)	1.7% (4;176)	3.8% (22;371)	3.6% (2;98)	4.1% (17;346)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 5.0, and OSHPD hospital patient discharge data, 2015. See Technical Note for details about methodology.

‡ Data are reported for January – September 2015 due to coding changes from ICD-9-CM to ICD-10-CM/PCS, which began October 1, 2015. Comparisons of IMIs across years should be made with caution since previous years' results are based on 12 months of data, while this analysis is based on 9 months of data.

*****Hospital name changed from 2014 to 2015, see Technical Note for details.***

Note: Blank cells indicate that fewer than three conditions were treated.

Definition of Terms: Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. Total Deaths: number of patients that died in this hospital. Total Cases: number of patients with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.