# *California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators (Conditions), 2020*

HCAI Department of Health Care Access and Information

| County | Hospital Name | Acute Myocardial Infarction | Acute Stroke | | | | Gastro-Intestinal Hemorrhage | Heart Failure | Hip Fracture | Pneumonia |
|---|---|---|---|---|---|---|---|---|---|---|
| | | | Total | Hemorrhagic | Ischemic | Subarachnoid | | | | |
| | | Note: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating. | | | | | | | | |
| | STATEWIDE | 5.9%<br>(2,660;44,767) | 8.5%<br>(5,089;59,573) | 19.9%<br>(2,341;11,790) | 4.8%<br>(2,170;45,165) | 22.1%<br>(578;2,618) | 2.8%<br>(1,415;51,109) | 2.9%<br>(2,795;97,448) | 1.5%<br>(385;25,657) | 6.3%<br>(6,055;95,457) |
| Alameda | Alameda Hospital | 0.0%<br>(0;4) | 3.3%<br>(1;84) | 0.0%<br>(0;3) | 2.0%<br>(1;81) | | 5.2%<br>(6;70) | 2.4%<br>(3;144) | 5.3%<br>(1;28) | 6.2%<br>(12;170) |
| | Alta Bates Summit Medical Center | 5.3%<br>(20;382) | 7.2%<br>(15;360) | 25.1%<br>(4;12) | 3.6%<br>(11;348) | | 2.8%<br>(8;278) | 2.3%<br>(17;713) | 0.0%<br>(0;102) | 5.5%<br>(13;271) |
| | Alta Bates Summit Medical Center – Alta Bates Campus | 10.0%<br>(4;16) | 7.2%<br>(19;197) | 19.5%<br>(17;102) | 2.7%<br>(2;89) | **0.0%<br>(0;6)<br>Better** | 1.8%<br>(3;154) | 3.3%<br>(11;261) | 2.4%<br>(2;79) | 7.2%<br>(10;135) |
| | Eden Medical Center | 2.8%<br>(1;27) | 8.4%<br>(47;444) | 15.4%<br>(16;89) | 6.2%<br>(18;313) | 22.1%<br>(13;42) | 1.8%<br>(3;162) | 3.3%<br>(12;346) | 0.7%<br>(1;127) | 6.0%<br>(16;217) |
| | Highland Hospital | 5.3%<br>(7;110) | 9.7%<br>(16;155) | 21.0%<br>(10;39) | 6.9%<br>(5;113) | 20.0%<br>(1;3) | 2.7%<br>(7;209) | **1.2%<br>(5;509)<br>Better** | 0.0%<br>(0;75) | 6.6%<br>(22;396) |
| | Kaiser Foundation Hospital – Fremont | **0.0%<br>(0;70)<br>Better** | **15.7%<br>(14;115)<br>Worse** | 21.6%<br>(3;19) | **11.0%<br>(11;95)<br>Worse** | | **0.0%<br>(0;118)<br>Better** | 4.4%<br>(8;156) | 0.0%<br>(0;61) | 5.2%<br>(14;227) |
| | Kaiser Foundation Hospital – Oakland/Richmond | 4.7%<br>(13;302) | 11.3%<br>(35;474) | **33.7%<br>(22;94)<br>Worse** | 4.7%<br>(13;376) | 0.0%<br>(0;4) | 1.6%<br>(8;334) | 2.4%<br>(20;756) | 1.1%<br>(2;174) | 5.0%<br>(29;603) |
| | Kaiser Foundation Hospital – San Leandro | 5.8%<br>(3;42) | **3.7%<br>(8;216)<br>Better** | **7.1%<br>(3;47)<br>Better** | 2.5%<br>(5;164) | 0.0%<br>(0;5) | 4.7%<br>(9;158) | 3.5%<br>(15;348) | 2.0%<br>(2;88) | 7.3%<br>(26;364) |
| | Saint Rose Hospital | **10.3%<br>(20;179)<br>Worse** | 8.3%<br>(2;32) | 28.0%<br>(2;4) | 0.0%<br>(0;28) | | 5.5%<br>(5;136) | 3.1%<br>(5;203) | 0.0%<br>(0;19) | 7.0%<br>(22;279) |
| | Stanford Health Care – ValleyCare | 8.1%<br>(13;140) | 9.7%<br>(9;113) | 22.0%<br>(3;13) | 5.3%<br>(4;97) | 27.8%<br>(2;3) | 1.5%<br>(3;122) | 2.2%<br>(9;314) | 2.1%<br>(2;81) | 5.2%<br>(15;248) |
| | Washington Hospital – Fremont | 7.7%<br>(18;247) | 11.0%<br>(29;271) | 27.8%<br>(15;55) | 6.3%<br>(10;204) | 20.9%<br>(4;12) | 4.4%<br>(9;230) | 4.2%<br>(17;429) | 0.0%<br>(0;77) | **9.0%<br>(31;423)<br>Worse** |
| Amador | Sutter Amador Hospital | | 10.6%<br>(2;65) | | 6.8%<br>(2;63) | | 0.0%<br>(0;47) | 6.1%<br>(3;61) | 5.6%<br>(2;38) | 7.2%<br>(6;123) |

**Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 2021, and HCAI hospital patient discharge data, 2020. See Technical Note for details about methodology.**
****Hospital name changed from 2019 to 2020, see Technical Note for details.**
**Note: Blank cells indicate that fewer than three conditions were treated**
**Definition of Terms: Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. Total Deaths: number of patients that died in this hospital. Total Cases: number of patients with this medical condition in this hospital.**
**Quality Ratings are risk-adjusted using age, gender, APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.**

# *California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators (Conditions), 2020*

HCAI Department of Health Care Access and Information

| County | Hospital Name | Acute Myocardial Infarction | Acute Stroke | | | | Gastro-Intestinal Hemorrhage | Heart Failure | Hip Fracture | Pneumonia |
|---|---|---|---|---|---|---|---|---|---|---|
| | | | Total | Hemorrhagic | Ischemic | Subarachnoid | | | | |
| | | Note: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating. | | | | | | | | |
| | STATEWIDE | 5.9%<br>(2,660;44,767) | 8.5%<br>(5,089;59,573) | 19.9%<br>(2,341;11,790) | 4.8%<br>(2,170;45,165) | 22.1%<br>(578;2,618) | 2.8%<br>(1,415;51,109) | 2.9%<br>(2,795;97,448) | 1.5%<br>(385;25,657) | 6.3%<br>(6,055;95,457) |
| Butte | Enloe Medical Center – Esplanade | 5.7%<br>(22;376) | **13.6%<br>(42;383)<br>Worse** | **37.4%<br>(21;66)<br>Worse** | 6.3%<br>(17;311) | 38.2%<br>(4;6) | 3.5%<br>(10;413) | 3.3%<br>(27;843) | 0.0%<br>(0;212) | 7.5%<br>(49;745) |
| | Orchard Hospital | | | | | | 0.0%<br>(0;4) | 0.0%<br>(0;18) | | 7.6%<br>(1;36) |
| | Oroville Hospital | 3.9%<br>(2;54) | 10.4%<br>(13;183) | 21.6%<br>(2;7) | 5.5%<br>(9;173) | 45.6%<br>(2;3) | 1.2%<br>(3;213) | **0.9%<br>(3;445)<br>Better** | 1.1%<br>(1;65) | **4.4%<br>(27;942)<br>Better** |
| Calaveras | Mark Twain Medical Center | | 0.0%<br>(0;14) | | 0.0%<br>(0;13) | | 0.0%<br>(0;4) | 0.0%<br>(0;52) | 5.6%<br>(1;24) | 4.3%<br>(3;94) |
| Colusa | Colusa Medical Center | 0.0%<br>(0;3) | | | | | 0.0%<br>(0;5) | 5.7%<br>(2;51) | | 5.7%<br>(2;55) |
| Contra Costa | Contra Costa Regional Medical Center | 0.0%<br>(0;4) | 10.9%<br>(2;55) | 73.8%<br>(2;4) | 0.0%<br>(0;51) | | 4.7%<br>(2;61) | 5.5%<br>(6;275) | 0.0%<br>(0;9) | **19.5%<br>(7;129)<br>Worse** |
| | John Muir Medical Center – Concord Campus | 5.1%<br>(17;340) | 12.9%<br>(11;153) | 38.0%<br>(3;5) | 6.7%<br>(8;148) | | 1.9%<br>(6;275) | 2.1%<br>(13;659) | 0.0%<br>(0;74) | 7.0%<br>(31;409) |
| | John Muir Medical Center – Walnut Creek Campus | 9.1%<br>(14;97) | 10.1%<br>(85;536) | 24.4%<br>(35;133) | 5.8%<br>(39;359) | 21.5%<br>(11;44) | 2.9%<br>(11;366) | 3.1%<br>(18;481) | 2.5%<br>(5;212) | 6.8%<br>(43;539) |
| | Kaiser Foundation Hospital – Antioch | 6.0%<br>(4;51) | 8.9%<br>(16;184) | 29.4%<br>(7;31) | 3.8%<br>(7;149) | 25.9%<br>(2;4) | 1.9%<br>(2;122) | 2.9%<br>(8;337) | 4.0%<br>(2;68) | 4.4%<br>(8;259) |
| | Kaiser Foundation Hospital – Walnut Creek | 4.5%<br>(9;256) | 7.0%<br>(10;234) | 7.1%<br>(2;57) | 6.0%<br>(8;174) | 0.0%<br>(0;3) | 2.0%<br>(5;242) | 3.4%<br>(13;404) | 2.6%<br>(5;225) | 8.1%<br>(30;437) |
| | San Ramon Regional Medical Center | 6.3%<br>(5;77) | 5.4%<br>(4;96) | 26.8%<br>(3;5) | 1.2%<br>(1;91) | | 4.1%<br>(3;108) | 3.3%<br>(5;132) | 1.9%<br>(1;50) | 7.6%<br>(11;195) |
| | Sutter Delta Medical Center | 7.6%<br>(14;229) | 2.2%<br>(1;95) | 0.0%<br>(0;4) | 1.4%<br>(1;90) | | 3.6%<br>(3;126) | 2.6%<br>(10;537) | 2.8%<br>(2;70) | 8.5%<br>(15;245) |
| Del Norte | Sutter Coast Hospital | 0.0%<br>(0;9) | 21.1%<br>(3;17) | | 11.2%<br>(2;15) | | 7.8%<br>(4;38) | 2.3%<br>(2;74) | 0.0%<br>(0;19) | 6.0%<br>(5;97) |

**Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 2021, and HCAI hospital patient discharge data, 2020. See Technical Note for details about methodology.**

****Hospital name changed from 2019 to 2020, see Technical Note for details.**

**Note: Blank cells indicate that fewer than three conditions were treated**

**Definition of Terms: Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. Total Deaths: number of patients that died in this hospital. Total Cases: number of patients with this medical condition in this hospital.**

**Quality Ratings are risk-adjusted using age, gender, APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.**

# *California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators (Conditions), 2020*

HCAI Department of Health Care Access and Information

| County | Hospital Name | Acute Myocardial Infarction | Acute Stroke | | | | Gastro-Intestinal Hemorrhage | Heart Failure | Hip Fracture | Pneumonia |
|---|---|---|---|---|---|---|---|---|---|---|
| | | | Total | Hemorrhagic | Ischemic | Subarachnoid | | | | |
| | | Note: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating. | | | | | | | | |
| | STATEWIDE | 5.9%<br>(2,660;44,767) | 8.5%<br>(5,089;59,573) | 19.9%<br>(2,341;11,790) | 4.8%<br>(2,170;45,165) | 22.1%<br>(578;2,618) | 2.8%<br>(1,415;51,109) | 2.9%<br>(2,795;97,448) | 1.5%<br>(385;25,657) | 6.3%<br>(6,055;95,457) |
| El Dorado | Barton Memorial Hospital | | 0.0%<br>(0;10) | | 0.0%<br>(0;10) | | 0.0%<br>(0;19) | 10.2%<br>(3;47) | 0.0%<br>(0;22) | 9.8%<br>(2;68) |
| | Marshall Medical Center | 13.3%<br>(2;19) | 5.9%<br>(6;105) | 33.6%<br>(4;5) | **0.8%<br>(1;99)<br>Better** | | 1.0%<br>(1;130) | 3.5%<br>(7;230) | 1.2%<br>(1;94) | 3.0%<br>(6;240) |
| Fresno | Adventist Health Reedley | 0.0%<br>(0;3) | 33.9%<br>(2;17) | | 19.1%<br>(2;17) | | | 0.0%<br>(0;16) | | 0.0%<br>(0;25) |
| | Clovis Community Medical Center | 7.3%<br>(14;162) | 5.8%<br>(5;251) | 20.4%<br>(1;4) | 3.1%<br>(4;245) | | 3.6%<br>(10;339) | **5.9%<br>(23;481)<br>Worse** | 1.1%<br>(2;176) | **9.7%<br>(40;560)<br>Worse** |
| | Coalinga Regional Medical Center | | | | | | | | | 0.0%<br>(0;3) |
| | Community Regional Medical Center – Fresno | **8.9%<br>(34;512)<br>Worse** | **10.3%<br>(123;859)<br>Worse** | 20.8%<br>(74;307) | **8.2%<br>(41;510)<br>Worse** | 23.6%<br>(8;42) | 2.4%<br>(9;539) | **4.7%<br>(34;1,242)<br>Worse** | 1.6%<br>(2;147) | 5.7%<br>(25;834) |
| | Kaiser Foundation Hospital – Fresno | 2.6%<br>(3;109) | 7.9%<br>(10;222) | 23.4%<br>(5;41) | 4.0%<br>(5;178) | 0.0%<br>(0;3) | 3.8%<br>(6;155) | 3.9%<br>(12;349) | 0.7%<br>(1;162) | 5.2%<br>(15;384) |
| | Saint Agnes Medical Center | **8.6%<br>(32;433)<br>Worse** | **12.7%<br>(23;342)<br>Worse** | 33.0%<br>(9;28) | 6.6%<br>(13;313) | | **4.7%<br>(24;582)<br>Worse** | **4.7%<br>(35;1,022)<br>Worse** | 3.6%<br>(6;207) | **9.9%<br>(57;902)<br>Worse** |
| Glenn | Glenn Medical Center | | | | | | | 0.0%<br>(0;6) | | 0.0%<br>(0;7) |
| Humboldt | Jerold Phelps Community Hospital | | | | | | | 0.0%<br>(0;3) | | 0.0%<br>(0;5) |
| | Mad River Community Hospital | | 0.0%<br>(0;21) | | 0.0%<br>(0;21) | | 6.2%<br>(1;20) | 13.3%<br>(2;23) | 0.0%<br>(0;14) | 13.1%<br>(5;90) |
| | Redwood Memorial Hospital | | 0.0%<br>(0;5) | | 0.0%<br>(0;5) | | 0.0%<br>(0;28) | 4.4%<br>(1;28) | | 11.0%<br>(4;53) |
| | Saint Joseph Hospital – Eureka | 5.3%<br>(6;174) | 8.0%<br>(5;95) | 13.6%<br>(2;14) | 4.0%<br>(2;80) | | 1.1%<br>(1;101) | 3.2%<br>(8;264) | 0.0%<br>(0;118) | 6.2%<br>(20;319) |

**Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 2021, and HCAI hospital patient discharge data, 2020. See Technical Note for details about methodology.**

****Hospital name changed from 2019 to 2020, see Technical Note for details.**

**Note: Blank cells indicate that fewer than three conditions were treated**

**Definition of Terms: Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. Total Deaths: number of patients that died in this hospital. Total Cases: number of patients with this medical condition in this hospital.**

**Quality Ratings are risk-adjusted using age, gender, APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.**

# *California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators (Conditions), 2020*

HCAI Department of Health Care Access and Information

| County | Hospital Name | Acute Myocardial Infarction | Acute Stroke | | | | Gastro-Intestinal Hemorrhage | Heart Failure | Hip Fracture | Pneumonia |
|---|---|---|---|---|---|---|---|---|---|---|
| | | | Total | Hemorrhagic | Ischemic | Subarachnoid | | | | |
| | | Note: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating. | | | | | | | | |
| | STATEWIDE | 5.9%<br>(2,660;44,767) | 8.5%<br>(5,089;59,573) | 19.9%<br>(2,341;11,790) | 4.8%<br>(2,170;45,165) | 22.1%<br>(578;2,618) | 2.8%<br>(1,415;51,109) | 2.9%<br>(2,795;97,448) | 1.5%<br>(385;25,657) | 6.3%<br>(6,055;95,457) |
| Imperial | El Centro Regional Medical Center | 12.8%<br>(2;15) | 15.0%<br>(6;101) | | 7.0%<br>(4;99) | | 4.7%<br>(7;147) | 4.5%<br>(9;240) | 0.0%<br>(0;45) | **11.9%<br>(31;288)<br>Worse** |
| | Pioneers Memorial Healthcare District | 0.0%<br>(0;11) | 12.1%<br>(6;54) | 23.9%<br>(3;10) | 8.3%<br>(3;44) | | 5.1%<br>(2;25) | 1.5%<br>(1;94) | 5.1%<br>(1;23) | 7.5%<br>(10;160) |
| Inyo | Northern Inyo Hospital | | 0.0%<br>(0;9) | | 0.0%<br>(0;7) | | 5.2%<br>(1;15) | 3.2%<br>(1;36) | 6.8%<br>(1;17) | 0.0%<br>(0;42) |
| | Southern Inyo Hospital | | | | | | | | | 0.0%<br>(0;4) |
| Kern | Adventist Health Bakersfield | 6.5%<br>(30;433) | 9.3%<br>(36;452) | 20.3%<br>(15;85) | 5.3%<br>(19;362) | 45.1%<br>(2;5) | 2.5%<br>(10;393) | 2.8%<br>(15;541) | 2.2%<br>(4;207) | 6.2%<br>(37;638) |
| | Adventist Health Delano | | 30.6%<br>(1;14) | | 17.2%<br>(1;14) | | 0.0%<br>(0;28) | 3.5%<br>(2;75) | 0.0%<br>(0;10) | 14.8%<br>(7;83) |
| | Adventist Health Tehachapi Valley | 0.0%<br>(0;4) | 0.0%<br>(0;6) | | 0.0%<br>(0;4) | | 0.0%<br>(0;5) | 0.0%<br>(0;33) | | 3.3%<br>(1;40) |
| | Bakersfield Heart Hospital | 3.3%<br>(6;244) | 5.6%<br>(1;34) | | 0.0%<br>(0;32) | | 4.7%<br>(2;69) | 3.2%<br>(11;317) | | 4.7%<br>(8;191) |
| | Bakersfield Memorial Hospital | 8.0%<br>(27;377) | 9.6%<br>(44;467) | 19.1%<br>(17;97) | 6.2%<br>(17;338) | 26.7%<br>(10;32) | **5.6%<br>(13;315)<br>Worse** | 3.3%<br>(16;686) | 2.0%<br>(2;127) | **11.4%<br>(35;484)<br>Worse** |
| | Good Samaritan Hospital – Bakersfield | 8.0%<br>(1;21) | | | | | 0.0%<br>(0;6) | 0.0%<br>(0;45) | | 4.9%<br>(2;96) |
| | Kern Medical Center | 5.1%<br>(2;34) | 11.7%<br>(7;90) | 32.7%<br>(2;13) | 5.8%<br>(3;73) | 30.9%<br>(2;4) | **6.9%<br>(8;103)<br>Worse** | 4.4%<br>(4;195) | 0.0%<br>(0;49) | **17.0%<br>(9;108)<br>Worse** |
| | Kern Valley Healthcare District | | | | | | 0.0%<br>(0;3) | 0.0%<br>(0;9) | | 8.3%<br>(1;37) |
| | Mercy Hospital – Bakersfield | 10.2%<br>(8;38) | 6.4%<br>(10;193) | 15.8%<br>(6;35) | 3.5%<br>(4;155) | 0.0%<br>(0;3) | 3.1%<br>(4;200) | 2.9%<br>(7;285) | 2.1%<br>(2;134) | 8.5%<br>(25;439) |
| | Ridgecrest Regional Hospital | | 24.1%<br>(2;21) | | 13.6%<br>(2;21) | | 3.6%<br>(1;31) | 4.6%<br>(3;62) | 12.0%<br>(1;14) | 11.1%<br>(7;78) |

**Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 2021, and HCAI hospital patient discharge data, 2020. See Technical Note for details about methodology.**

****Hospital name changed from 2019 to 2020, see Technical Note for details.**

**Note: Blank cells indicate that fewer than three conditions were treated**

**Definition of Terms: Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. Total Deaths: number of patients that died in this hospital. Total Cases: number of patients with this medical condition in this hospital.**

**Quality Ratings are risk-adjusted using age, gender, APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.**

# *California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators (Conditions), 2020*

HCAI Department of Health Care Access and Information

| County | Hospital Name | Acute Myocardial Infarction | Acute Stroke | | | | Gastro-Intestinal Hemorrhage | Heart Failure | Hip Fracture | Pneumonia |
|---|---|---|---|---|---|---|---|---|---|---|
| | | | Total | Hemorrhagic | Ischemic | Subarachnoid | | | | |
| | | Note: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating. | | | | | | | | |
| | STATEWIDE | 5.9%<br>(2,660;44,767) | 8.5%<br>(5,089;59,573) | 19.9%<br>(2,341;11,790) | 4.8%<br>(2,170;45,165) | 22.1%<br>(578;2,618) | 2.8%<br>(1,415;51,109) | 2.9%<br>(2,795;97,448) | 1.5%<br>(385;25,657) | 6.3%<br>(6,055;95,457) |
| Kings | Adventist Health Hanford | 11.4%<br>(8;89) | 11.7%<br>(10;185) | 47.3%<br>(3;6) | 5.6%<br>(7;179) | | 3.1%<br>(4;115) | 4.0%<br>(17;407) | **5.8%<br>(4;83)<br>Worse** | 8.2%<br>(22;394) |
| Lake | Adventist Health Clearlake | | 17.7%<br>(1;18) | | 10.0%<br>(1;18) | | 8.0%<br>(1;16) | 2.1%<br>(1;78) | | 6.1%<br>(4;110) |
| | Sutter Lakeside Hospital | 0.0%<br>(0;3) | 0.0%<br>(0;5) | | 0.0%<br>(0;5) | | 0.0%<br>(0;35) | 0.0%<br>(0;82) | 2.8%<br>(1;34) | 1.7%<br>(1;76) |
| Lassen | Banner Lassen Medical Center | 14.8%<br>(1;3) | 0.0%<br>(0;4) | | 0.0%<br>(0;4) | | 0.0%<br>(0;12) | 0.0%<br>(0;24) | 0.0%<br>(0;5) | 15.4%<br>(3;52) |
| Los Angeles | Adventist Health Glendale | 4.6%<br>(20;382) | 7.1%<br>(40;443) | 15.2%<br>(16;85) | 3.8%<br>(16;318) | 26.5%<br>(8;40) | 1.7%<br>(5;250) | **1.5%<br>(11;576)<br>Better** | 1.7%<br>(4;182) | 5.8%<br>(68;904) |
| | Adventist Health White Memorial | 7.4%<br>(15;206) | 8.0%<br>(8;200) | 34.9%<br>(6;31) | 2.0%<br>(2;168) | | 4.4%<br>(9;253) | 1.8%<br>(7;493) | 2.6%<br>(1;51) | 5.7%<br>(14;304) |
| | Alhambra Hospital Medical Center | 4.4%<br>(2;18) | **1.6%<br>(1;52)<br>Better** | 0.0%<br>(0;9) | **1.1%<br>(1;43)<br>Better** | | 4.5%<br>(3;73) | 2.6%<br>(3;129) | 0.0%<br>(0;26) | 6.2%<br>(23;325) |
| | Antelope Valley Hospital | **12.8%<br>(28;312)<br>Worse** | 9.3%<br>(33;366) | 19.1%<br>(17;86) | 6.3%<br>(15;275) | 21.4%<br>(1;5) | **7.2%<br>(18;400)<br>Worse** | 4.4%<br>(12;455) | 1.5%<br>(2;173) | **10.4%<br>(60;748)<br>Worse** |
| | Beverly Hospital | 5.6%<br>(9;157) | 9.7%<br>(10;170) | 24.0%<br>(5;28) | 5.3%<br>(5;141) | | 3.7%<br>(7;229) | 2.1%<br>(4;265) | 2.3%<br>(1;58) | 8.7%<br>(26;339) |
| | California Hospital Medical Center – Los Angeles | 4.2%<br>(6;143) | 5.1%<br>(8;215) | 14.9%<br>(5;46) | 2.2%<br>(3;168) | | 2.4%<br>(4;236) | **0.5%<br>(3;703)<br>Better** | 0.0%<br>(0;30) | 4.8%<br>(15;347) |
| | Casa Colina Hospital | | 0.0%<br>(0;26) | 0.0%<br>(0;5) | 0.0%<br>(0;21) | | 0.0%<br>(0;27) | 0.0%<br>(0;52) | 0.0%<br>(0;19) | **0.0%<br>(0;56)<br>Better** |
| | Cedars-Sinai Marina Del Rey Hospital | 0.0%<br>(0;10) | 6.1%<br>(1;40) | | 3.4%<br>(1;40) | | 1.3%<br>(3;112) | 3.8%<br>(6;133) | 1.7%<br>(1;55) | 4.3%<br>(14;235) |

**Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 2021, and HCAI hospital patient discharge data, 2020. See Technical Note for details about methodology.**
****Hospital name changed from 2019 to 2020, see Technical Note for details.**
**Note: Blank cells indicate that fewer than three conditions were treated**
**Definition of Terms: Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. Total Deaths: number of patients that died in this hospital. Total Cases: number of patients with this medical condition in this hospital.**
**Quality Ratings are risk-adjusted using age, gender, APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.**

# *California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators (Conditions), 2020*

HCAI Department of Health Care Access and Information

| County | Hospital Name | Acute Myocardial Infarction | Acute Stroke | | | | Gastro-Intestinal Hemorrhage | Heart Failure | Hip Fracture | Pneumonia |
|---|---|---|---|---|---|---|---|---|---|---|
| | | | Total | Hemorrhagic | Ischemic | Subarachnoid | | | | |
| | | Note: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating. | | | | | | | | |
| | STATEWIDE | 5.9%<br>(2,660;44,767) | 8.5%<br>(5,089;59,573) | 19.9%<br>(2,341;11,790) | 4.8%<br>(2,170;45,165) | 22.1%<br>(578;2,618) | 2.8%<br>(1,415;51,109) | 2.9%<br>(2,795;97,448) | 1.5%<br>(385;25,657) | 6.3%<br>(6,055;95,457) |
| Los Angeles (continued) | Cedars-Sinai Medical Center | 6.3%<br>(33;546) | 8.8%<br>(62;650) | 21.0%<br>(32;167) | 4.9%<br>(20;425) | 22.0%<br>(10;58) | 3.0%<br>(21;811) | **4.0%<br>(78;1,578)<br>Worse** | 0.6%<br>(2;235) | 6.8%<br>(99;1,022) |
| | Centinela Hospital Medical Center | 3.9%<br>(8;236) | **4.9%<br>(13;303)<br>Better** | 15.2%<br>(8;50) | **2.0%<br>(5;251)<br>Better** | | 1.6%<br>(4;286) | **0.8%<br>(5;717)<br>Better** | 0.0%<br>(0;50) | **4.0%<br>(57;802)<br>Better** |
| | City of Hope Helford Clinical Research Hospital | | 9.8%<br>(1;15) | 0.0%<br>(0;8) | 11.5%<br>(1;6) | | 2.3%<br>(1;41) | 9.7%<br>(1;14) | | 8.6%<br>(18;179) |
| | Coast Plaza Hospital | 3.5%<br>(1;22) | 4.5%<br>(1;31) | 17.5%<br>(1;4) | 0.0%<br>(0;27) | | **8.6%<br>(5;57)<br>Worse** | 2.2%<br>(3;136) | 0.0%<br>(0;21) | 4.9%<br>(15;204) |
| | College Medical Center | 7.9%<br>(1;5) | 45.8%<br>(1;5) | | 34.1%<br>(1;4) | | 0.0%<br>(0;74) | 2.4%<br>(2;82) | 16.1%<br>(1;10) | **11.6%<br>(14;119)<br>Worse** |
| | Community and Mission Hospital of Huntington Park – Slauson | 6.1%<br>(3;45) | 6.0%<br>(1;28) | | 3.4%<br>(1;28) | | 4.7%<br>(2;73) | 3.6%<br>(4;184) | 0.0%<br>(0;19) | 9.0%<br>(23;189) |
| | East Los Angeles Doctors Hospital | 0.0%<br>(0;17) | 0.0%<br>(0;12) | | 0.0%<br>(0;10) | | 2.1%<br>(1;45) | 5.5%<br>(3;104) | 0.0%<br>(0;3) | 9.3%<br>(20;153) |
| | Emanate Health Foothill Presbyterian Hospital | 9.6%<br>(3;24) | 3.8%<br>(1;79) | 0.0%<br>(0;4) | 2.3%<br>(1;75) | | 3.7%<br>(3;140) | 1.2%<br>(2;163) | 3.8%<br>(2;59) | 6.5%<br>(19;327) |
| | Emanate Health Inter-Community Hospital | 6.2%<br>(25;467) | 8.8%<br>(7;70) | 26.5%<br>(4;12) | 0.0%<br>(0;55) | 28.4%<br>(3;3) | 3.8%<br>(4;182) | **5.2%<br>(14;334)<br>Worse** | 0.0%<br>(0;72) | **9.0%<br>(40;396)<br>Worse** |
| | Emanate Health Queen of the Valley Hospital | 10.9%<br>(5;31) | 6.0%<br>(14;281) | 12.8%<br>(6;52) | 2.9%<br>(5;216) | 30.3%<br>(3;13) | 3.3%<br>(6;222) | 4.3%<br>(8;281) | 1.9%<br>(1;77) | 8.4%<br>(26;456) |
| | Encino Hospital Medical Center | 12.8%<br>(2;8) | 8.1%<br>(1;13) | | 0.0%<br>(0;12) | | 2.7%<br>(1;27) | 0.7%<br>(1;62) | 0.0%<br>(0;11) | **1.6%<br>(3;90)<br>Better** |
| | Garfield Medical Center | 7.1%<br>(13;227) | 8.0%<br>(24;287) | 18.0%<br>(12;67) | 3.7%<br>(8;206) | 43.7%<br>(4;14) | 4.0%<br>(7;152) | 4.1%<br>(10;281) | 2.7%<br>(1;47) | **13.9%<br>(40;309)<br>Worse** |
| | Glendale Memorial Hospital and Medical Center | 6.9%<br>(13;187) | 9.3%<br>(7;98) | 17.2%<br>(2;20) | 5.8%<br>(5;78) | | 5.5%<br>(3;103) | 1.6%<br>(5;361) | 1.5%<br>(1;73) | 7.0%<br>(29;424) |

**Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 2021, and HCAI hospital patient discharge data, 2020. See Technical Note for details about methodology.**

****Hospital name changed from 2019 to 2020, see Technical Note for details.**

**Note: Blank cells indicate that fewer than three conditions were treated**

**Definition of Terms: Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. Total Deaths: number of patients that died in this hospital. Total Cases: number of patients with this medical condition in this hospital.**

**Quality Ratings are risk-adjusted using age, gender, APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.**

# *California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators (Conditions), 2020*

HCAI Department of Health Care Access and Information

| County | Hospital Name | Acute Myocardial Infarction | Acute Stroke | | | | Gastro-Intestinal Hemorrhage | Heart Failure | Hip Fracture | Pneumonia |
|---|---|---|---|---|---|---|---|---|---|---|
| | | | Total | Hemorrhagic | Ischemic | Subarachnoid | | | | |
| | | Note: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating. | | | | | | | | |
| | STATEWIDE | 5.9%<br>(2,660;44,767) | 8.5%<br>(5,089;59,573) | 19.9%<br>(2,341;11,790) | 4.8%<br>(2,170;45,165) | 22.1%<br>(578;2,618) | 2.8%<br>(1,415;51,109) | 2.9%<br>(2,795;97,448) | 1.5%<br>(385;25,657) | 6.3%<br>(6,055;95,457) |
| Los Angeles (continued) | Good Samaritan Hospital – Los Angeles | 6.4%<br>(29;250) | 8.8%<br>(75;516) | 20.2%<br>(38;125) | 4.7%<br>(28;363) | 29.5%<br>(9;28) | 2.3%<br>(9;229) | 2.4%<br>(11;470) | 1.8%<br>(1;65) | 7.9%<br>(44;363) |
| | Greater El Monte Community Hospital | **47.1%<br>(4;14)<br>Worse** | 0.0%<br>(0;21) | | 0.0%<br>(0;21) | | 3.7%<br>(2;50) | **8.9%<br>(5;93)<br>Worse** | | 8.2%<br>(20;223) |
| | Henry Mayo Newhall Hospital | 5.7%<br>(14;210) | 9.4%<br>(11;164) | 18.2%<br>(4;26) | 5.9%<br>(7;138) | | 2.7%<br>(6;199) | 4.6%<br>(19;369) | 1.6%<br>(2;126) | 6.1%<br>(27;403) |
| | Hollywood Presbyterian Medical Center | 8.1%<br>(17;263) | 6.6%<br>(11;165) | 11.1%<br>(4;23) | 4.8%<br>(6;139) | 19.7%<br>(1;3) | 2.9%<br>(6;214) | 1.9%<br>(7;420) | 2.8%<br>(3;86) | 5.1%<br>(43;562) |
| | Huntington Memorial Hospital | 4.9%<br>(13;255) | 9.5%<br>(40;470) | 20.5%<br>(16;73) | 4.8%<br>(16;374) | 41.2%<br>(8;23) | 2.3%<br>(9;381) | 2.8%<br>(17;565) | 0.5%<br>(1;201) | 5.8%<br>(47;683) |
| | Kaiser Foundation Hospital – Baldwin Park | 10.2%<br>(7;62) | 7.7%<br>(10;147) | 14.4%<br>(3;28) | 4.4%<br>(6;116) | 67.4%<br>(1;3) | 1.7%<br>(4;166) | 4.0%<br>(13;400) | 1.5%<br>(2;117) | 5.5%<br>(10;228) |
| | Kaiser Foundation Hospital – Downey | 2.5%<br>(2;86) | 5.7%<br>(8;240) | 6.9%<br>(1;26) | 3.5%<br>(6;211) | 24.0%<br>(1;3) | 2.1%<br>(4;243) | **1.0%<br>(4;439)<br>Better** | 1.4%<br>(1;96) | 8.4%<br>(28;351) |
| | Kaiser Foundation Hospital – Panorama City | 1.4%<br>(1;91) | 5.1%<br>(5;117) | 7.4%<br>(1;20) | 3.8%<br>(4;95) | | 3.7%<br>(5;137) | 2.4%<br>(5;160) | 0.9%<br>(1;142) | 7.5%<br>(17;201) |
| | Kaiser Foundation Hospital – South Bay | 7.5%<br>(4;51) | 11.1%<br>(17;189) | 17.0%<br>(3;32) | 6.6%<br>(12;153) | 53.2%<br>(2;4) | 2.6%<br>(4;125) | **6.7%<br>(20;294)<br>Worse** | 0.0%<br>(0;103) | 7.6%<br>(17;206) |
| | Kaiser Foundation Hospital – Sunset | 5.1%<br>(42;1,168) | 9.1%<br>(73;705) | 22.8%<br>(44;244) | 5.0%<br>(24;402) | 16.6%<br>(5;59) | 3.0%<br>(9;255) | **4.4%<br>(31;550)<br>Worse** | 3.7%<br>(4;105) | 5.7%<br>(19;275) |
| | Kaiser Foundation Hospital – West Los Angeles | 5.9%<br>(4;86) | 7.3%<br>(13;244) | 7.1%<br>(1;27) | 4.0%<br>(10;213) | **100.0%<br>(2;4)<br>Worse** | 2.2%<br>(3;159) | 3.8%<br>(15;254) | 5.1%<br>(4;100) | 5.0%<br>(13;210) |
| | Kaiser Foundation Hospital – Woodland Hills | 3.4%<br>(4;80) | 7.3%<br>(9;169) | 0.0%<br>(0;25) | 5.2%<br>(9;141) | 0.0%<br>(0;3) | 4.2%<br>(7;133) | 2.5%<br>(11;242) | 1.0%<br>(2;188) | 4.7%<br>(17;247) |
| | Keck Hospital of University of Southern California | 6.0%<br>(13;85) | **6.4%<br>(72;406)<br>Better** | **14.1%<br>(33;158)<br>Better** | 4.4%<br>(19;149) | **15.0%<br>(20;99)<br>Better** | 2.8%<br>(7;104) | 3.5%<br>(17;206) | 0.0%<br>(0;11) | 5.5%<br>(6;70) |

**Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 2021, and HCAI hospital patient discharge data, 2020. See Technical Note for details about methodology.**

****Hospital name changed from 2019 to 2020, see Technical Note for details.**

**Note: Blank cells indicate that fewer than three conditions were treated**

**Definition of Terms: Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. Total Deaths: number of patients that died in this hospital. Total Cases: number of patients with this medical condition in this hospital.**

**Quality Ratings are risk-adjusted using age, gender, APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.**

# *California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators (Conditions), 2020*

HCAI Department of Health Care Access and Information

| County | Hospital Name | Acute Myocardial Infarction | Acute Stroke | | | | Gastro-Intestinal Hemorrhage | Heart Failure | Hip Fracture | Pneumonia |
|---|---|---|---|---|---|---|---|---|---|---|
| | | | Total | Hemorrhagic | Ischemic | Subarachnoid | | | | |
| | | Note: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating. | | | | | | | | |
| | STATEWIDE | 5.9%<br>(2,660;44,767) | 8.5%<br>(5,089;59,573) | 19.9%<br>(2,341;11,790) | 4.8%<br>(2,170;45,165) | 22.1%<br>(578;2,618) | 2.8%<br>(1,415;51,109) | 2.9%<br>(2,795;97,448) | 1.5%<br>(385;25,657) | 6.3%<br>(6,055;95,457) |
| Los Angeles (continued) | L.A. Downtown Medical Center | 0.0%<br>(0;3) | | | | | 2.5%<br>(2;62) | 4.0%<br>(2;45) | 0.0%<br>(0;10) | 7.0%<br>(19;272) |
| | Lakewood Regional Medical Center | **8.9%<br>(24;229)<br>Worse** | 8.8%<br>(20;200) | 21.4%<br>(13;51) | 3.5%<br>(4;139) | 34.3%<br>(3;10) | 3.9%<br>(6;215) | 2.0%<br>(9;385) | 3.1%<br>(2;76) | **9.0%<br>(47;388)<br>Worse** |
| | Los Angeles Community Hospital | 0.0%<br>(0;15) | 0.0%<br>(0;39) | | 0.0%<br>(0;36) | | 2.5%<br>(3;139) | 1.2%<br>(3;250) | 4.0%<br>(1;15) | **4.8%<br>(41;515)<br>Better** |
| | Los Angeles County/Harbor – UCLA Medical Center | 8.4%<br>(23;262) | **14.0%<br>(22;197)<br>Worse** | 25.4%<br>(8;40) | 9.4%<br>(9;148) | **43.8%<br>(5;9)<br>Worse** | 2.4%<br>(5;210) | 1.7%<br>(7;570) | 1.6%<br>(1;54) | 7.9%<br>(14;210) |
| | Los Angeles County/Olive View – UCLA Medical Center | 7.3%<br>(4;84) | 8.9%<br>(3;101) | 0.0%<br>(0;10) | 6.2%<br>(3;90) | | 1.6%<br>(2;127) | 1.5%<br>(5;447) | | 5.1%<br>(12;210) |
| | Los Angeles County/Rancho Los Amigos National Rehabilitation Center | | 3.6%<br>(1;85) | 0.0%<br>(0;5) | 2.4%<br>(1;79) | | | 5.9%<br>(1;33) | | 5.8%<br>(1;30) |
| | Los Angeles County/University of Southern California Medical Center | 6.2%<br>(13;204) | 8.9%<br>(44;309) | 21.1%<br>(22;78) | 5.3%<br>(17;204) | 18.3%<br>(5;27) | 3.2%<br>(16;383) | 2.2%<br>(22;1,031) | 0.0%<br>(0;62) | 7.3%<br>(35;516) |
| | Memorial Hospital of Gardena | 7.5%<br>(3;44) | 3.9%<br>(2;102) | 0.0%<br>(0;4) | 2.4%<br>(2;98) | | 0.6%<br>(1;149) | 3.6%<br>(8;340) | 0.0%<br>(0;21) | 4.8%<br>(32;419) |
| | MemorialCare Long Beach Medical Center | 5.5%<br>(20;312) | 10.1%<br>(56;663) | 21.9%<br>(22;134) | 6.6%<br>(28;499) | 18.8%<br>(6;30) | 3.3%<br>(13;345) | 2.8%<br>(14;574) | 1.9%<br>(3;135) | 7.1%<br>(33;474) |
| | Methodist Hospital of Southern California | 4.3%<br>(9;169) | **12.1%<br>(66;468)<br>Worse** | 24.3%<br>(25;93) | **8.0%<br>(34;357)<br>Worse** | 27.1%<br>(7;18) | 3.0%<br>(9;249) | 3.7%<br>(18;395) | 0.6%<br>(1;145) | 8.1%<br>(55;589) |
| | Mission Community Hospital – Panorama Campus | 0.0%<br>(0;12) | 0.0%<br>(0;27) | 0.0%<br>(0;7) | 0.0%<br>(0;20) | | 2.9%<br>(3;84) | 1.7%<br>(3;148) | 0.0%<br>(0;22) | **3.8%<br>(27;418)<br>Better** |
| | Monterey Park Hospital | 12.7%<br>(1;11) | 5.3%<br>(1;41) | 0.0%<br>(0;9) | 5.3%<br>(1;32) | | 9.7%<br>(2;52) | 2.4%<br>(2;98) | 0.0%<br>(0;12) | 11.8%<br>(9;124) |
| | Northridge Hospital Medical Center | **4.0%<br>(20;405)<br>Better** | 8.2%<br>(39;372) | 15.7%<br>(14;82) | 5.3%<br>(21;267) | 21.7%<br>(4;23) | 1.5%<br>(4;230) | 1.6%<br>(8;357) | 0.8%<br>(1;114) | **4.3%<br>(28;436)<br>Better** |

**Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 2021, and HCAI hospital patient discharge data, 2020. See Technical Note for details about methodology.**
****Hospital name changed from 2019 to 2020, see Technical Note for details.**
**Note: Blank cells indicate that fewer than three conditions were treated**
**Definition of Terms: Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. Total Deaths: number of patients that died in this hospital. Total Cases: number of patients with this medical condition in this hospital.**
**Quality Ratings are risk-adjusted using age, gender, APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.**

# *California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators (Conditions), 2020*

HCAI Department of Health Care Access and Information

| County | Hospital Name | Acute Myocardial Infarction | Acute Stroke | | | | Gastro-Intestinal Hemorrhage | Heart Failure | Hip Fracture | Pneumonia |
|---|---|---|---|---|---|---|---|---|---|---|
| | | | Total | Hemorrhagic | Ischemic | Subarachnoid | | | | |
| | | Note: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating. | | | | | | | | |
| | STATEWIDE | 5.9%<br>(2,660;44,767) | 8.5%<br>(5,089;59,573) | 19.9%<br>(2,341;11,790) | 4.8%<br>(2,170;45,165) | 22.1%<br>(578;2,618) | 2.8%<br>(1,415;51,109) | 2.9%<br>(2,795;97,448) | 1.5%<br>(385;25,657) | 6.3%<br>(6,055;95,457) |
| Los Angeles (continued) | Norwalk Community Hospital | 3.4%<br>(1;24) | 0.0%<br>(0;14) | | 0.0%<br>(0;14) | | 0.0%<br>(0;51) | 0.0%<br>(0;72) | 0.0%<br>(0;11) | **2.4%<br>(8;252)<br>Better** |
| | Olympia Medical Center | 11.2%<br>(6;48) | 0.0%<br>(0;19) | | 0.0%<br>(0;17) | | 3.3%<br>(2;72) | 2.0%<br>(3;154) | 0.0%<br>(0;15) | 6.4%<br>(30;330) |
| | PIH Health Hospital – Whittier | 5.3%<br>(18;397) | **11.9%<br>(95;703)<br>Worse** | **27.6%<br>(45;160)<br>Worse** | **6.8%<br>(38;500)<br>Worse** | 30.3%<br>(12;43) | 3.3%<br>(12;434) | 2.4%<br>(15;806) | 0.9%<br>(1;142) | 6.8%<br>(42;612) |
| | PIH Health Hospital – Downey | 9.2%<br>(14;156) | 4.0%<br>(2;130) | | 0.0%<br>(0;127) | | 1.7%<br>(4;289) | 3.2%<br>(10;413) | 4.9%<br>(4;86) | 4.7%<br>(17;372) |
| | Pacifica Hospital of the Valley | 0.0%<br>(0;5) | 0.0%<br>(0;7) | | 0.0%<br>(0;6) | | 6.7%<br>(1;26) | 0.0%<br>(0;27) | 11.7%<br>(1;7) | 14.2%<br>(7;119) |
| | Palmdale Regional Medical Center | 6.8%<br>(10;181) | 5.7%<br>(6;189) | 6.4%<br>(1;24) | 3.7%<br>(4;163) | | 5.6%<br>(9;200) | 3.2%<br>(8;379) | 1.9%<br>(1;64) | **10.4%<br>(31;371)<br>Worse** |
| | Pomona Valley Hospital Medical Center | 5.1%<br>(19;322) | 8.3%<br>(89;706) | 19.0%<br>(39;149) | 4.2%<br>(28;501) | 26.0%<br>(22;56) | 3.5%<br>(14;267) | 2.6%<br>(20;662) | 1.4%<br>(3;195) | 6.2%<br>(33;400) |
| | Providence Cedars-Sinai Tarzana Medical Center** | 8.3%<br>(10;147) | 6.3%<br>(5;118) | 21.5%<br>(2;11) | 3.0%<br>(3;106) | | 1.5%<br>(3;176) | 2.5%<br>(11;358) | 2.3%<br>(3;120) | 6.0%<br>(14;224) |
| | Providence Holy Cross Medical Center | 3.8%<br>(9;224) | 10.0%<br>(40;385) | 24.4%<br>(23;96) | 5.2%<br>(15;279) | 30.1%<br>(2;10) | 1.8%<br>(6;304) | **1.1%<br>(10;668)<br>Better** | 0.6%<br>(1;123) | **3.2%<br>(15;383)<br>Better** |
| | Providence Little Company of Mary Medical Center – San Pedro | 9.0%<br>(2;11) | 16.0%<br>(6;59) | 69.3%<br>(2;3) | 7.3%<br>(4;56) | | 2.3%<br>(4;106) | 3.0%<br>(5;152) | 2.8%<br>(2;50) | 7.8%<br>(12;175) |
| | Providence Little Company of Mary Medical Center – Torrance | 6.5%<br>(26;387) | 7.1%<br>(38;558) | 17.7%<br>(19;116) | 3.5%<br>(13;419) | 20.1%<br>(6;23) | 4.1%<br>(17;370) | 3.2%<br>(25;686) | 1.2%<br>(2;153) | **9.5%<br>(44;487)<br>Worse** |
| | Providence Saint John's Health Center | **10.2%<br>(16;164)<br>Worse** | 9.3%<br>(23;255) | 26.3%<br>(13;52) | 4.4%<br>(8;189) | 16.5%<br>(2;14) | 3.2%<br>(8;164) | 2.6%<br>(9;245) | 1.7%<br>(2;119) | 6.3%<br>(24;312) |
| | Providence Saint Joseph Medical Center | 8.0%<br>(19;222) | 7.7%<br>(39;481) | 15.4%<br>(16;89) | 5.3%<br>(20;366) | 13.9%<br>(3;26) | 2.1%<br>(7;313) | 2.6%<br>(17;508) | 2.3%<br>(5;193) | 6.0%<br>(30;482) |

**Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 2021, and HCAI hospital patient discharge data, 2020. See Technical Note for details about methodology.**

****Hospital name changed from 2019 to 2020, see Technical Note for details.**

**Note: Blank cells indicate that fewer than three conditions were treated**

**Definition of Terms: Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. Total Deaths: number of patients that died in this hospital. Total Cases: number of patients with this medical condition in this hospital.**

**Quality Ratings are risk-adjusted using age, gender, APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.**

# *California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators (Conditions), 2020*

HCAI Department of Health Care Access and Information

| County | Hospital Name | Acute Myocardial Infarction | Acute Stroke | | | | Gastro-Intestinal Hemorrhage | Heart Failure | Hip Fracture | Pneumonia |
|---|---|---|---|---|---|---|---|---|---|---|
| | | | Total | Hemorrhagic | Ischemic | Subarachnoid | | | | |
| | | Note: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating. | | | | | | | | |
| | STATEWIDE | 5.9%<br>(2,660;44,767) | 8.5%<br>(5,089;59,573) | 19.9%<br>(2,341;11,790) | 4.8%<br>(2,170;45,165) | 22.1%<br>(578;2,618) | 2.8%<br>(1,415;51,109) | 2.9%<br>(2,795;97,448) | 1.5%<br>(385;25,657) | 6.3%<br>(6,055;95,457) |
| Los Angeles (continued) | Ronald Reagan UCLA Medical Center | 6.0%<br>(20;199) | **4.0%**<br>**(26;357)**<br>**Better** | **7.5%**<br>**(11;114)**<br>**Better** | **2.7%**<br>**(10;211)**<br>**Better** | 13.1%<br>(5;32) | 2.8%<br>(11;212) | 2.0%<br>(12;426) | 2.6%<br>(1;35) | **3.2%**<br>**(19;376)**<br>**Better** |
| | Saint Francis Medical Center | 6.5%<br>(17;261) | 7.1%<br>(19;258) | 16.8%<br>(9;52) | 3.6%<br>(7;200) | 21.2%<br>(3;6) | 2.5%<br>(5;230) | 2.6%<br>(8;477) | 3.2%<br>(2;46) | 7.6%<br>(42;407) |
| | Saint Mary Medical Center – Long Beach | 4.7%<br>(17;249) | 9.2%<br>(24;252) | 22.7%<br>(11;63) | 4.4%<br>(6;162) | 25.0%<br>(7;27) | 1.9%<br>(4;184) | 1.7%<br>(6;457) | 0.0%<br>(0;43) | 7.1%<br>(21;290) |
| | Saint Vincent Medical Center | 0.0%<br>(0;8) | | | | | 0.0%<br>(0;9) | 0.0%<br>(0;11) | 0.0%<br>(0;4) | 0.0%<br>(0;11) |
| | San Dimas Community Hospital | 35.2%<br>(2;8) | 0.0%<br>(0;17) | | 0.0%<br>(0;17) | | 3.9%<br>(3;54) | 1.3%<br>(1;75) | 0.0%<br>(0;23) | 7.7%<br>(32;287) |
| | San Gabriel Valley Medical Center | 4.9%<br>(2;30) | 5.1%<br>(3;102) | 43.0%<br>(2;12) | 1.2%<br>(1;88) | | 2.8%<br>(5;159) | 3.0%<br>(7;169) | 1.2%<br>(1;81) | **4.3%**<br>**(26;428)**<br>**Better** |
| | Santa Monica – UCLA Medical Center and Orthopedic Hospital | 5.2%<br>(5;105) | **1.3%**<br>**(1;94)**<br>**Better** | 0.0%<br>(0;6) | **0.9%**<br>**(1;85)**<br>**Better** | 0.0%<br>(0;3) | 2.4%<br>(5;174) | 1.7%<br>(8;319) | 1.9%<br>(4;168) | **3.3%**<br>**(25;493)**<br>**Better** |
| | Sherman Oaks Hospital | 4.9%<br>(4;66) | 2.7%<br>(1;39) | 18.4%<br>(1;5) | 0.0%<br>(0;33) | | **7.2%**<br>**(8;72)**<br>**Worse** | **0.5%**<br>**(2;203)**<br>**Better** | 1.4%<br>(1;63) | **4.0%**<br>**(27;332)**<br>**Better** |
| | Southern California Hospital at Culver City | 2.4%<br>(4;242) | 2.1%<br>(1;75) | 34.7%<br>(1;8) | **0.0%**<br>**(0;67)**<br>**Better** | | 0.9%<br>(1;147) | 2.0%<br>(6;379) | 0.0%<br>(0;27) | **3.8%**<br>**(36;614)**<br>**Better** |
| | Southern California Hospital at Hollywood | 0.0%<br>(0;4) | 0.0%<br>(0;12) | | 0.0%<br>(0;12) | | 2.1%<br>(1;61) | 1.4%<br>(1;61) | 0.0%<br>(0;11) | **3.4%**<br>**(24;497)**<br>**Better** |
| | Torrance Memorial Medical Center | **3.7%**<br>**(16;400)**<br>**Better** | 10.1%<br>(47;506) | 25.5%<br>(27;110) | 5.6%<br>(18;383) | 13.2%<br>(2;13) | 3.3%<br>(14;470) | 3.0%<br>(36;934) | 1.2%<br>(3;227) | 7.6%<br>(63;726) |
| | University of Southern California Verdugo Hills Hospital | 0.0%<br>(0;13) | 3.8%<br>(2;96) | 16.8%<br>(2;10) | 0.0%<br>(0;86) | | 0.0%<br>(0;95) | 1.9%<br>(4;151) | 1.8%<br>(1;68) | 5.2%<br>(19;315) |
| | Valley Presbyterian Hospital | 6.4%<br>(13;169) | 14.6%<br>(6;96) | 28.2%<br>(2;12) | 9.5%<br>(4;83) | | 3.3%<br>(4;164) | 4.8%<br>(12;348) | 3.3%<br>(2;56) | **10.4%**<br>**(38;367)**<br>**Worse** |

**Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 2021, and HCAI hospital patient discharge data, 2020. See Technical Note for details about methodology.**

****Hospital name changed from 2019 to 2020, see Technical Note for details.**

**Note: Blank cells indicate that fewer than three conditions were treated**

**Definition of Terms: Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. Total Deaths: number of patients that died in this hospital. Total Cases: number of patients with this medical condition in this hospital.**

**Quality Ratings are risk-adjusted using age, gender, APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.**

# *California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators (Conditions), 2020*

HCAI Department of Health Care Access and Information

| County | Hospital Name | Acute Myocardial Infarction | Acute Stroke | | | | Gastro-Intestinal Hemorrhage | Heart Failure | Hip Fracture | Pneumonia |
|---|---|---|---|---|---|---|---|---|---|---|
| | | | Total | Hemorrhagic | Ischemic | Subarachnoid | | | | |
| | | Note: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating. | | | | | | | | |
| | STATEWIDE | 5.9%<br>(2,660;44,767) | 8.5%<br>(5,089;59,573) | 19.9%<br>(2,341;11,790) | 4.8%<br>(2,170;45,165) | 22.1%<br>(578;2,618) | 2.8%<br>(1,415;51,109) | 2.9%<br>(2,795;97,448) | 1.5%<br>(385;25,657) | 6.3%<br>(6,055;95,457) |
| Los Angeles (continued) | West Covina Medical Center | | | | | | | 0.0%<br>(0;3) | | 0.0%<br>(0;17) |
| | West Hills Hospital and Medical Center | 4.0%<br>(7;216) | **3.7%<br>(9;229)<br>Better** | **10.2%<br>(6;49)<br>Better** | 1.8%<br>(3;168) | 0.0%<br>(0;12) | 0.7%<br>(1;148) | 1.8%<br>(4;225) | 2.0%<br>(2;122) | **3.7%<br>(15;373)<br>Better** |
| | Whittier Hospital Medical Center | 4.7%<br>(1;19) | 10.6%<br>(2;44) | | 7.9%<br>(2;41) | | 2.4%<br>(2;140) | 1.1%<br>(2;211) | 2.5%<br>(1;40) | **11.3%<br>(30;242)<br>Worse** |
| Madera | Madera Community Hospital | 2.1%<br>(1;30) | 6.5%<br>(1;54) | | 3.6%<br>(1;54) | | 4.9%<br>(3;74) | 5.2%<br>(5;141) | 2.0%<br>(1;39) | 5.5%<br>(5;107) |
| Marin | Kaiser Foundation Hospital – San Rafael | 4.0%<br>(3;82) | 14.4%<br>(9;128) | 35.8%<br>(3;22) | 7.2%<br>(5;102) | 69.8%<br>(1;4) | 1.4%<br>(2;115) | 2.8%<br>(5;156) | 1.3%<br>(2;131) | 4.1%<br>(8;188) |
| | MarinHealth Medical Center | 4.2%<br>(7;189) | 11.0%<br>(21;185) | 26.5%<br>(13;35) | 5.7%<br>(6;147) | 31.1%<br>(2;3) | 2.0%<br>(5;135) | 1.7%<br>(7;206) | 1.1%<br>(1;107) | 5.2%<br>(17;257) |
| | Novato Community Hospital | 9.7%<br>(1;6) | 7.4%<br>(1;39) | | 0.0%<br>(0;38) | | 0.0%<br>(0;41) | 2.6%<br>(2;79) | 1.9%<br>(1;43) | 4.3%<br>(2;70) |
| Mariposa | John C. Fremont Healthcare District | | 0.0%<br>(0;6) | | 0.0%<br>(0;5) | | | 0.0%<br>(0;37) | | 8.6%<br>(1;23) |
| Mendocino | Adventist Health Howard Memorial | | 4.7%<br>(1;37) | | 2.7%<br>(1;36) | | 0.0%<br>(0;19) | 2.2%<br>(1;60) | 3.4%<br>(2;49) | 3.2%<br>(2;75) |
| | Adventist Health Mendocino Coast** | | 27.6%<br>(2;9) | | 27.1%<br>(1;8) | | 10.4%<br>(1;11) | 2.7%<br>(1;35) | 0.0%<br>(0;12) | 24.3%<br>(3;31) |
| | Adventist Health Ukiah Valley | 0.0%<br>(0;8) | 3.7%<br>(1;70) | | 0.0%<br>(0;68) | | 3.2%<br>(3;71) | 0.8%<br>(1;133) | 0.0%<br>(0;28) | 6.6%<br>(8;174) |
| Merced | Memorial Hospital Los Banos | | 4.1%<br>(1;35) | | 3.0%<br>(1;33) | | 0.0%<br>(0;3) | 5.6%<br>(2;43) | 0.0%<br>(0;28) | 12.8%<br>(4;57) |
| | Mercy Medical Center – Merced | 6.3%<br>(4;63) | 3.2%<br>(3;220) | 0.0%<br>(0;4) | 2.0%<br>(3;214) | | 1.2%<br>(1;182) | 3.1%<br>(15;594) | 2.2%<br>(2;104) | 7.5%<br>(31;540) |

**Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 2021, and HCAI hospital patient discharge data, 2020. See Technical Note for details about methodology.**

****Hospital name changed from 2019 to 2020, see Technical Note for details.**

**Note: Blank cells indicate that fewer than three conditions were treated**

**Definition of Terms: Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. Total Deaths: number of patients that died in this hospital. Total Cases: number of patients with this medical condition in this hospital.**

**Quality Ratings are risk-adjusted using age, gender, APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.**

# *California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators (Conditions), 2020*

HCAI Department of Health Care Access and Information

| County | Hospital Name | Acute Myocardial Infarction | Acute Stroke | | | | Gastro-Intestinal Hemorrhage | Heart Failure | Hip Fracture | Pneumonia |
|---|---|---|---|---|---|---|---|---|---|---|
| | | | Total | Hemorrhagic | Ischemic | Subarachnoid | | | | |
| | | Note: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating. | | | | | | | | |
| | STATEWIDE | 5.9%<br>(2,660;44,767) | 8.5%<br>(5,089;59,573) | 19.9%<br>(2,341;11,790) | 4.8%<br>(2,170;45,165) | 22.1%<br>(578;2,618) | 2.8%<br>(1,415;51,109) | 2.9%<br>(2,795;97,448) | 1.5%<br>(385;25,657) | 6.3%<br>(6,055;95,457) |
| Modoc | Modoc Medical Center | | | | | | | 0.0%<br>(0;15) | | 7.9%<br>(1;29) |
| Mono | Mammoth Hospital | | | | | | 9.3%<br>(1;15) | 0.0%<br>(0;7) | 0.0%<br>(0;6) | 0.0%<br>(0;11) |
| Monterey | Community Hospital of The Monterey Peninsula | 5.5%<br>(8;183) | **12.6%<br>(25;272)<br>Worse** | 21.9%<br>(8;35) | **8.5%<br>(16;235)<br>Worse** | | 3.1%<br>(7;290) | 4.2%<br>(18;436) | 3.7%<br>(4;105) | **9.3%<br>(48;538)<br>Worse** |
| | George L. Mee Memorial Hospital | | | | | | | 17.1%<br>(2;21) | | 0.0%<br>(0;19) |
| | Natividad Medical Center | 0.0%<br>(0;7) | 15.2%<br>(2;48) | 70.9%<br>(2;9) | 0.0%<br>(0;38) | | 3.6%<br>(3;118) | 1.9%<br>(1;99) | 2.0%<br>(1;61) | 7.7%<br>(4;95) |
| | Salinas Valley Memorial Hospital | 4.0%<br>(7;235) | **12.3%<br>(30;297)<br>Worse** | 21.0%<br>(13;66) | **9.8%<br>(15;225)<br>Worse** | 38.1%<br>(2;6) | 2.0%<br>(3;260) | 4.6%<br>(16;452) | 1.7%<br>(1;53) | 7.0%<br>(22;359) |
| Napa | Adventist Health Saint Helena | 4.9%<br>(5;129) | 5.7%<br>(1;16) | | 3.2%<br>(1;16) | | 0.0%<br>(0;49) | 1.0%<br>(1;92) | 3.4%<br>(1;21) | 7.3%<br>(5;59) |
| | Queen of the Valley Medical Center | 6.9%<br>(5;97) | 11.4%<br>(13;140) | **40.7%<br>(7;18)<br>Worse** | 4.5%<br>(6;122) | | 3.4%<br>(6;149) | 2.1%<br>(3;141) | 0.0%<br>(0;97) | 9.3%<br>(14;190) |
| Nevada | Sierra Nevada Memorial Hospital | 7.6%<br>(2;17) | 7.3%<br>(5;90) | 18.2%<br>(1;4) | 3.9%<br>(3;84) | | 0.9%<br>(1;109) | 2.6%<br>(6;207) | 0.8%<br>(1;112) | 4.2%<br>(7;190) |
| | Tahoe Forest Hospital | | 0.0%<br>(0;3) | | 0.0%<br>(0;3) | | 9.2%<br>(2;23) | 7.7%<br>(2;31) | 0.0%<br>(0;23) | 4.9%<br>(1;59) |
| Orange | AHMC Anaheim Regional Medical Center | 6.6%<br>(12;187) | 5.9%<br>(2;52) | 21.0%<br>(1;3) | 2.4%<br>(1;49) | | 4.9%<br>(6;155) | 3.3%<br>(7;284) | 0.0%<br>(0;44) | 7.8%<br>(24;340) |
| | Anaheim Global Medical Center | 4.7%<br>(2;41) | 8.3%<br>(1;9) | | 0.0%<br>(0;8) | | 0.0%<br>(0;37) | 0.0%<br>(0;43) | 0.0%<br>(0;5) | 7.6%<br>(9;113) |
| | Chapman Global Medical Center | | 14.2%<br>(3;11) | | 10.0%<br>(1;9) | | **22.9%<br>(3;19)<br>Worse** | **11.6%<br>(4;42)<br>Worse** | 0.0%<br>(0;11) | **12.8%<br>(14;81)<br>Worse** |

**Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 2021, and HCAI hospital patient discharge data, 2020. See Technical Note for details about methodology.**

****Hospital name changed from 2019 to 2020, see Technical Note for details.**

**Note: Blank cells indicate that fewer than three conditions were treated**

**Definition of Terms: Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. Total Deaths: number of patients that died in this hospital. Total Cases: number of patients with this medical condition in this hospital.**

**Quality Ratings are risk-adjusted using age, gender, APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.**

# *California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators (Conditions), 2020*

HCAI Department of Health Care Access and Information

| County | Hospital Name | Acute Myocardial Infarction | Acute Stroke | | | | Gastro-Intestinal Hemorrhage | Heart Failure | Hip Fracture | Pneumonia |
|---|---|---|---|---|---|---|---|---|---|---|
| | | | Total | Hemorrhagic | Ischemic | Subarachnoid | | | | |
| | | Note: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating. | | | | | | | | |
| | STATEWIDE | 5.9%<br>(2,660;44,767) | 8.5%<br>(5,089;59,573) | 19.9%<br>(2,341;11,790) | 4.8%<br>(2,170;45,165) | 22.1%<br>(578;2,618) | 2.8%<br>(1,415;51,109) | 2.9%<br>(2,795;97,448) | 1.5%<br>(385;25,657) | 6.3%<br>(6,055;95,457) |
| Orange (continued) | Foothill Regional Medical Center | 0.0%<br>(0;5) | 0.0%<br>(0;22) | 0.0%<br>(0;4) | 0.0%<br>(0;18) | | 0.0%<br>(0;54) | 0.0%<br>(0;61) | 0.0%<br>(0;28) | **1.8%<br>(7;278)<br>Better** |
| | Fountain Valley Regional Hospital and Medical Center – Euclid | 5.9%<br>(19;211) | 7.5%<br>(41;392) | 18.7%<br>(23;80) | 3.5%<br>(13;297) | 25.1%<br>(5;15) | 3.2%<br>(9;286) | 1.9%<br>(9;522) | 2.5%<br>(2;86) | 6.6%<br>(47;653) |
| | Garden Grove Hospital and Medical Center | 10.1%<br>(4;18) | 0.0%<br>(0;43) | | 0.0%<br>(0;42) | | 2.8%<br>(5;124) | 2.8%<br>(2;87) | 1.6%<br>(1;36) | 5.3%<br>(29;322) |
| | Hoag Memorial Hospital Presbyterian | 6.9%<br>(23;412) | **5.1%<br>(27;680)<br>Better** | 16.6%<br>(18;109) | **1.3%<br>(5;545)<br>Better** | 18.6%<br>(4;26) | 2.6%<br>(15;633) | 2.6%<br>(23;745) | 1.4%<br>(4;331) | **4.7%<br>(43;986)<br>Better** |
| | Hoag Orthopedic Institute | | | | | | | | 0.0%<br>(0;79) | |
| | Huntington Beach Hospital | 9.1%<br>(1;12) | 0.0%<br>(0;15) | | 0.0%<br>(0;15) | | 3.8%<br>(2;46) | 3.5%<br>(2;68) | 0.0%<br>(0;11) | 4.0%<br>(10;189) |
| | Kaiser Foundation Hospital – Orange County – Anaheim | 7.3%<br>(10;201) | **4.0%<br>(17;446)<br>Better** | **6.7%<br>(5;116)<br>Better** | **2.6%<br>(10;301)<br>Better** | 12.9%<br>(2;29) | 2.9%<br>(5;279) | 2.0%<br>(13;539) | **0.0%<br>(0;267)<br>Better** | 6.0%<br>(30;551) |
| | La Palma Intercommunity Hospital | 5.0%<br>(1;29) | 0.0%<br>(0;22) | | 0.0%<br>(0;22) | | 1.6%<br>(1;71) | 1.1%<br>(1;95) | 0.0%<br>(0;30) | 4.3%<br>(12;208) |
| | Los Alamitos Medical Center | 5.5%<br>(11;169) | 8.5%<br>(49;426) | 20.7%<br>(26;85) | 4.3%<br>(16;318) | 24.5%<br>(7;23) | 3.3%<br>(5;177) | 2.9%<br>(13;339) | 0.7%<br>(1;165) | 8.0%<br>(27;319) |
| | MemorialCare Orange Coast Medical Center | 6.6%<br>(17;195) | 8.1%<br>(8;146) | 18.1%<br>(4;18) | 5.7%<br>(4;124) | 0.0%<br>(0;4) | 3.1%<br>(8;233) | 3.2%<br>(13;357) | 2.1%<br>(3;106) | **3.8%<br>(13;343)<br>Better** |
| | MemorialCare Saddleback Medical Center | 5.0%<br>(12;200) | 9.5%<br>(37;328) | 20.8%<br>(17;64) | 5.6%<br>(16;251) | 27.0%<br>(4;13) | 2.5%<br>(9;264) | 3.0%<br>(14;371) | 1.2%<br>(2;162) | 5.4%<br>(20;309) |
| | Orange County Global Medical Center | 4.5%<br>(4;104) | 8.3%<br>(24;194) | 14.4%<br>(8;39) | 6.5%<br>(15;146) | 8.8%<br>(1;9) | 3.4%<br>(3;85) | 4.4%<br>(5;171) | 0.0%<br>(0;35) | 6.2%<br>(20;237) |
| | Placentia Linda Hospital | 5.2%<br>(3;41) | 19.3%<br>(3;31) | | 11.9%<br>(3;29) | | 5.1%<br>(2;84) | 3.1%<br>(4;129) | 0.0%<br>(0;67) | 5.7%<br>(11;217) |
| | Providence Mission Hospital** | 7.7%<br>(24;282) | 10.6%<br>(54;463) | 23.7%<br>(23;88) | 6.7%<br>(26;347) | 20.1%<br>(5;28) | 2.4%<br>(9;316) | 3.5%<br>(15;314) | 2.1%<br>(6;252) | **8.5%<br>(50;482)<br>Worse** |

**Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 2021, and HCAI hospital patient discharge data, 2020. See Technical Note for details about methodology.**

****Hospital name changed from 2019 to 2020, see Technical Note for details.**

**Note: Blank cells indicate that fewer than three conditions were treated**

**Definition of Terms: Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. Total Deaths: number of patients that died in this hospital. Total Cases: number of patients with this medical condition in this hospital.**

**Quality Ratings are risk-adjusted using age, gender, APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.**

# *California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators (Conditions), 2020*

HCAI Department of Health Care Access and Information

| County | Hospital Name | Acute Myocardial Infarction | Acute Stroke | | | | Gastro-Intestinal Hemorrhage | Heart Failure | Hip Fracture | Pneumonia |
|---|---|---|---|---|---|---|---|---|---|---|
| | | | Total | Hemorrhagic | Ischemic | Subarachnoid | | | | |
| | | Note: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating. | | | | | | | | |
| | STATEWIDE | 5.9%<br>(2,660;44,767) | 8.5%<br>(5,089;59,573) | 19.9%<br>(2,341;11,790) | 4.8%<br>(2,170;45,165) | 22.1%<br>(578;2,618) | 2.8%<br>(1,415;51,109) | 2.9%<br>(2,795;97,448) | 1.5%<br>(385;25,657) | 6.3%<br>(6,055;95,457) |
| Orange (continued) | Providence Mission Hospital – Laguna Beach** | | 0.0%<br>(0;3) | | | | 0.0%<br>(0;31) | 0.0%<br>(0;40) | 0.0%<br>(0;55) | 2.8%<br>(2;80) |
| | Providence St. Joseph Hospital** | 6.0%<br>(17;295) | 9.0%<br>(23;366) | 23.4%<br>(10;64) | 5.2%<br>(13;292) | 0.0%<br>(0;10) | 1.1%<br>(3;264) | 3.8%<br>(22;479) | 1.1%<br>(2;193) | 6.6%<br>(38;554) |
| | Providence St. Jude Medical Center** | 6.2%<br>(27;322) | 8.0%<br>(43;549) | 14.0%<br>(16;126) | 6.1%<br>(23;385) | 17.4%<br>(4;38) | **4.7%<br>(20;299)<br>Worse** | 3.4%<br>(17;443) | 0.0%<br>(0;201) | **8.7%<br>(85;673)<br>Worse** |
| | South Coast Global Medical Center | 16.1%<br>(1;5) | 0.0%<br>(0;22) | | 0.0%<br>(0;21) | | 0.0%<br>(0;27) | 6.2%<br>(4;51) | 0.0%<br>(0;3) | 9.3%<br>(12;133) |
| | UC Irvine Medical Center | 6.3%<br>(16;199) | **2.0%<br>(14;469)<br>Better** | **3.5%<br>(5;133)<br>Better** | **1.5%<br>(6;300)<br>Better** | **5.3%<br>(3;36)<br>Better** | **0.7%<br>(3;299)<br>Better** | **0.5%<br>(2;445)<br>Better** | 0.0%<br>(0;56) | **2.0%<br>(6;301)<br>Better** |
| | West Anaheim Medical Center | 7.2%<br>(13;141) | 11.3%<br>(2;41) | 25.5%<br>(1;3) | 6.6%<br>(1;38) | | 3.7%<br>(3;94) | 2.5%<br>(5;177) | 0.0%<br>(0;50) | **4.5%<br>(29;459)<br>Better** |
| Placer | Kaiser Foundation Hospital – Roseville | 5.2%<br>(15;465) | **13.4%<br>(25;311)<br>Worse** | **40.5%<br>(11;37)<br>Worse** | 6.5%<br>(13;270) | 24.7%<br>(1;4) | 3.3%<br>(11;402) | **5.5%<br>(37;697)<br>Worse** | 1.5%<br>(5;351) | 7.0%<br>(39;709) |
| | Sutter Auburn Faith Hospital | 0.0%<br>(0;9) | 0.0%<br>(0;90) | 0.0%<br>(0;3) | 0.0%<br>(0;84) | 0.0%<br>(0;3) | 2.8%<br>(1;44) | 1.9%<br>(3;134) | 1.4%<br>(1;67) | 3.1%<br>(3;144) |
| | Sutter Roseville Medical Center | **3.9%<br>(17;481)<br>Better** | 9.3%<br>(42;523) | 19.1%<br>(18;95) | 5.7%<br>(22;422) | 31.6%<br>(2;6) | 1.6%<br>(6;517) | 3.3%<br>(21;631) | 1.3%<br>(4;295) | 6.8%<br>(33;566) |
| Plumas | Eastern Plumas Hospital – Portola Campus | | | | | | | 0.0%<br>(0;3) | | 0.0%<br>(0;25) |
| | Plumas District Hospital | 0.0%<br>(0;3) | 0.0%<br>(0;4) | | 0.0%<br>(0;4) | | 0.0%<br>(0;7) | 0.0%<br>(0;8) | | 0.0%<br>(0;12) |
| | Seneca District Hospital | | | | | | | | | 0.0%<br>(0;9) |

**Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 2021, and HCAI hospital patient discharge data, 2020. See Technical Note for details about methodology.**

****Hospital name changed from 2019 to 2020, see Technical Note for details.**

**Note: Blank cells indicate that fewer than three conditions were treated**

**Definition of Terms: Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. Total Deaths: number of patients that died in this hospital. Total Cases: number of patients with this medical condition in this hospital.**

**Quality Ratings are risk-adjusted using age, gender, APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.**

# *California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators (Conditions), 2020*

HCAI Department of Health Care Access and Information

| County | Hospital Name | Acute Myocardial Infarction | Acute Stroke | | | | Gastro-Intestinal Hemorrhage | Heart Failure | Hip Fracture | Pneumonia |
|---|---|---|---|---|---|---|---|---|---|---|
| | | | Total | Hemorrhagic | Ischemic | Subarachnoid | | | | |
| | | Note: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating. | | | | | | | | |
| | STATEWIDE | 5.9%<br>(2,660;44,767) | 8.5%<br>(5,089;59,573) | 19.9%<br>(2,341;11,790) | 4.8%<br>(2,170;45,165) | 22.1%<br>(578;2,618) | 2.8%<br>(1,415;51,109) | 2.9%<br>(2,795;97,448) | 1.5%<br>(385;25,657) | 6.3%<br>(6,055;95,457) |
| Riverside | Corona Regional Medical Center – Main | 6.0%<br>(7;117) | 7.3%<br>(5;119) | 0.0%<br>(0;5) | 4.5%<br>(5;114) | | **0.5%<br>(1;203)<br>Better** | 3.0%<br>(10;353) | 1.0%<br>(1;108) | 7.4%<br>(24;353) |
| | Desert Regional Medical Center | 8.0%<br>(30;451) | **10.7%<br>(61;432)<br>Worse** | 25.5%<br>(33;112) | 6.1%<br>(23;298) | 22.7%<br>(5;22) | 4.2%<br>(12;357) | 3.7%<br>(14;371) | 2.1%<br>(5;246) | 6.9%<br>(34;478) |
| | Doctors Hospital of Riverside** | 12.3%<br>(3;16) | 8.1%<br>(1;40) | | 0.0%<br>(0;39) | | 5.5%<br>(6;114) | 5.0%<br>(7;229) | 0.0%<br>(0;57) | 9.6%<br>(23;324) |
| | Eisenhower Medical Center | 4.2%<br>(14;392) | 7.2%<br>(29;579) | 21.3%<br>(17;95) | 2.9%<br>(9;468) | 20.5%<br>(3;16) | 1.9%<br>(8;484) | 2.6%<br>(18;780) | 0.4%<br>(1;297) | 5.5%<br>(43;880) |
| | Hemet Global Medical Center | 4.4%<br>(3;51) | 0.0%<br>(0;52) | | 0.0%<br>(0;51) | | 3.2%<br>(5;215) | 1.7%<br>(7;480) | 2.3%<br>(3;125) | 4.8%<br>(24;607) |
| | John F. Kennedy Memorial Hospital | 5.1%<br>(8;121) | **17.7%<br>(9;96)<br>Worse** | | 10.1%<br>(8;95) | | 3.1%<br>(7;113) | 2.0%<br>(4;206) | 0.0%<br>(0;93) | 9.2%<br>(14;198) |
| | Kaiser Foundation Hospital – Moreno Valley | 5.6%<br>(2;48) | 5.8%<br>(3;86) | 0.0%<br>(0;12) | 3.6%<br>(2;73) | | 0.0%<br>(0;64) | 1.7%<br>(4;147) | 0.0%<br>(0;34) | 7.1%<br>(8;132) |
| | Kaiser Foundation Hospital – Riverside | 6.4%<br>(5;67) | 3.3%<br>(3;121) | 9.9%<br>(1;21) | 1.7%<br>(2;98) | | 4.3%<br>(5;108) | 2.2%<br>(5;190) | 2.7%<br>(2;90) | 7.5%<br>(15;200) |
| | Loma Linda University Medical Center – Murrieta | 7.0%<br>(26;404) | 8.6%<br>(8;145) | 27.8%<br>(7;23) | 1.6%<br>(1;122) | | **5.4%<br>(11;212)<br>Worse** | 2.9%<br>(9;424) | 1.5%<br>(1;88) | **9.8%<br>(22;357)<br>Worse** |
| | Menifee Global Medical Center | 10.0%<br>(1;10) | 10.2%<br>(2;9) | 20.1%<br>(1;4) | 7.0%<br>(1;5) | | 0.0%<br>(0;68) | 0.0%<br>(0;118) | 2.0%<br>(1;44) | 8.7%<br>(12;159) |
| | Palo Verde Hospital | 21.3%<br>(1;4) | | | | | 15.8%<br>(1;16) | 0.0%<br>(0;9) | | 0.0%<br>(0;11) |
| | Riverside Community Hospital | 5.0%<br>(40;719) | **5.8%<br>(73;802)<br>Better** | **14.6%<br>(31;169)<br>Better** | **2.6%<br>(29;582)<br>Better** | 23.6%<br>(13;51) | 3.1%<br>(14;299) | **0.5%<br>(4;633)<br>Better** | 0.8%<br>(2;189) | **3.4%<br>(24;566)<br>Better** |
| | Riverside University Health System – Medical Center | 10.1%<br>(7;49) | 9.3%<br>(32;299) | 23.9%<br>(24;84) | 4.1%<br>(8;214) | | 1.4%<br>(5;247) | 2.7%<br>(13;522) | 1.6%<br>(2;106) | 9.8%<br>(18;235) |

**Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 2021, and HCAI hospital patient discharge data, 2020. See Technical Note for details about methodology.**

****Hospital name changed from 2019 to 2020, see Technical Note for details.**

**Note: Blank cells indicate that fewer than three conditions were treated**

**Definition of Terms: Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. Total Deaths: number of patients that died in this hospital. Total Cases: number of patients with this medical condition in this hospital.**

**Quality Ratings are risk-adjusted using age, gender, APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.**

# *California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators (Conditions), 2020*

HCAI Department of Health Care Access and Information

| County | Hospital Name | Acute Myocardial Infarction | Acute Stroke | | | | Gastro-Intestinal Hemorrhage | Heart Failure | Hip Fracture | Pneumonia |
|---|---|---|---|---|---|---|---|---|---|---|
| | | | Total | Hemorrhagic | Ischemic | Subarachnoid | | | | |
| | | Note: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating. | | | | | | | | |
| | STATEWIDE | 5.9%<br>(2,660;44,767) | 8.5%<br>(5,089;59,573) | 19.9%<br>(2,341;11,790) | 4.8%<br>(2,170;45,165) | 22.1%<br>(578;2,618) | 2.8%<br>(1,415;51,109) | 2.9%<br>(2,795;97,448) | 1.5%<br>(385;25,657) | 6.3%<br>(6,055;95,457) |
| Riverside (continued) | San Gorgonio Memorial Hospital | 0.0%<br>(0;7) | 0.0%<br>(0;15) | | 0.0%<br>(0;15) | | **11.7%<br>(5;78)<br>Worse** | 2.4%<br>(2;139) | 0.0%<br>(0;58) | 11.4%<br>(9;132) |
| | Southwest Healthcare System – Murrieta | 9.9%<br>(3;24) | 7.6%<br>(28;388) | 21.1%<br>(18;63) | 3.1%<br>(8;316) | 22.1%<br>(2;9) | 1.7%<br>(4;273) | 3.2%<br>(14;414) | 1.0%<br>(2;208) | 5.8%<br>(20;499) |
| | Temecula Valley Hospital | 5.2%<br>(18;436) | 8.0%<br>(33;332) | 18.6%<br>(13;65) | 4.0%<br>(11;227) | 24.6%<br>(9;40) | 3.3%<br>(8;208) | 2.8%<br>(16;499) | 1.0%<br>(1;98) | 7.2%<br>(20;311) |
| | Vibra Rehabilitation Hospital of Rancho Mirage | | 0.0%<br>(0;10) | 0.0%<br>(0;6) | 0.0%<br>(0;3) | | | 0.0%<br>(0;3) | | |
| Sacramento | Kaiser Foundation Hospital – Sacramento | 8.2%<br>(11;102) | 8.9%<br>(70;681) | 20.3%<br>(35;195) | 6.1%<br>(29;412) | 13.2%<br>(6;74) | 2.7%<br>(8;288) | 3.9%<br>(16;461) | 2.3%<br>(4;148) | 4.6%<br>(19;404) |
| | Kaiser Foundation Hospital – South Sacramento | 5.8%<br>(17;362) | 9.7%<br>(21;325) | 26.2%<br>(12;55) | 4.7%<br>(9;268) | | 5.0%<br>(12;217) | 3.1%<br>(19;670) | 0.5%<br>(1;188) | 5.7%<br>(30;545) |
| | Mercy General Hospital | 5.3%<br>(43;990) | 8.6%<br>(20;244) | 19.5%<br>(14;89) | 5.4%<br>(6;154) | | 3.2%<br>(7;231) | 2.4%<br>(22;802) | 0.0%<br>(0;89) | 6.3%<br>(19;368) |
| | Mercy Hospital of Folsom | 4.0%<br>(2;21) | 10.9%<br>(8;158) | 37.9%<br>(2;6) | 5.5%<br>(6;152) | | 4.2%<br>(5;182) | 1.9%<br>(4;295) | 0.0%<br>(0;122) | 5.0%<br>(9;258) |
| | Mercy San Juan Medical Center** | 6.5%<br>(30;352) | 9.0%<br>(130;889) | 19.8%<br>(69;282) | 6.0%<br>(39;530) | 21.6%<br>(22;77) | 2.3%<br>(8;329) | 3.9%<br>(28;768) | 2.9%<br>(5;176) | 6.5%<br>(35;547) |
| | Methodist Hospital of Sacramento | 3.8%<br>(3;50) | 9.8%<br>(10;191) | 20.2%<br>(3;17) | 5.4%<br>(6;173) | | 1.0%<br>(2;185) | 4.0%<br>(16;409) | 5.7%<br>(3;55) | 7.6%<br>(19;288) |
| | Sutter Medical Center, Sacramento | 3.9%<br>(17;534) | 8.9%<br>(54;536) | 22.6%<br>(26;129) | 4.9%<br>(20;372) | 19.3%<br>(8;35) | 2.4%<br>(8;379) | 3.3%<br>(31;865) | 2.6%<br>(5;167) | 6.2%<br>(18;398) |
| | UC Davis Medical Center | 6.0%<br>(10;198) | **4.7%<br>(32;438)<br>Better** | **9.6%<br>(15;147)<br>Better** | **2.3%<br>(9;252)<br>Better** | 22.5%<br>(8;39) | 2.8%<br>(10;305) | 2.1%<br>(19;904) | 1.3%<br>(2;132) | 4.3%<br>(14;351) |
| San Benito | Hazel Hawkins Memorial Hospital | 0.0%<br>(0;4) | 33.7%<br>(3;26) | 49.3%<br>(1;4) | 26.9%<br>(2;22) | | 0.0%<br>(0;18) | 5.4%<br>(1;39) | 0.0%<br>(0;16) | 12.8%<br>(3;58) |
| San Bernardino | Arrowhead Regional Medical Center<br>Right click to view Comment Letter | 8.9%<br>(4;55) | 10.9%<br>(35;244) | 26.6%<br>(25;80) | 6.8%<br>(10;157) | 0.0%<br>(0;7) | 2.9%<br>(5;207) | 2.4%<br>(7;458) | 0.8%<br>(1;85) | 8.0%<br>(30;347) |

**Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 2021, and HCAI hospital patient discharge data, 2020. See Technical Note for details about methodology.**

****Hospital name changed from 2019 to 2020, see Technical Note for details.**

**Note: Blank cells indicate that fewer than three conditions were treated**

**Definition of Terms: Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. Total Deaths: number of patients that died in this hospital. Total Cases: number of patients with this medical condition in this hospital.**

**Quality Ratings are risk-adjusted using age, gender, APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.**

# *California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators (Conditions), 2020*

HCAI Department of Health Care Access and Information

| County | Hospital Name | Acute Myocardial Infarction | Acute Stroke | | | | Gastro-Intestinal Hemorrhage | Heart Failure | Hip Fracture | Pneumonia |
|---|---|---|---|---|---|---|---|---|---|---|
| | | | Total | Hemorrhagic | Ischemic | Subarachnoid | | | | |
| | | Note: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating. | | | | | | | | |
| | STATEWIDE | 5.9%<br>(2,660;44,767) | 8.5%<br>(5,089;59,573) | 19.9%<br>(2,341;11,790) | 4.8%<br>(2,170;45,165) | 22.1%<br>(578;2,618) | 2.8%<br>(1,415;51,109) | 2.9%<br>(2,795;97,448) | 1.5%<br>(385;25,657) | 6.3%<br>(6,055;95,457) |
| San Bernardino (continued) | Barstow Community Hospital | 0.0%<br>(0;8) | 0.0%<br>(0;7) | | 0.0%<br>(0;7) | | 7.2%<br>(1;34) | 4.1%<br>(2;79) | 0.0%<br>(0;16) | 8.3%<br>(6;113) |
| | Bear Valley Community Hospital | | | | | | | 0.0%<br>(0;5) | | 0.0%<br>(0;12) |
| | Chino Valley Medical Center | 4.4%<br>(2;59) | 0.0%<br>(0;31) | | 0.0%<br>(0;30) | | 2.0%<br>(1;79) | 0.8%<br>(1;135) | 0.0%<br>(0;13) | 3.9%<br>(10;272) |
| | Colorado River Medical Center | | | | | | 0.0%<br>(0;7) | 0.0%<br>(0;18) | | 17.8%<br>(1;25) |
| | Community Hospital of San Bernardino | 3.9%<br>(3;26) | 0.0%<br>(0;32) | | 0.0%<br>(0;31) | | **8.8%<br>(5;62)<br>Worse** | 4.9%<br>(7;196) | 5.4%<br>(1;25) | 4.3%<br>(7;211) |
| | Desert Valley Hospital | 5.1%<br>(17;307) | **0.0%<br>(0;86)<br>Better** | 0.0%<br>(0;3) | 0.0%<br>(0;83) | | 1.8%<br>(4;193) | **0.6%<br>(3;556)<br>Better** | 1.0%<br>(1;60) | **2.8%<br>(21;562)<br>Better** |
| | Hi-Desert Medical Center | 0.0%<br>(0;6) | 0.0%<br>(0;15) | | 0.0%<br>(0;15) | | | 6.6%<br>(1;36) | 0.0%<br>(0;29) | **27.0%<br>(8;91)<br>Worse** |
| | Kaiser Foundation Hospital – Fontana | 6.1%<br>(19;443) | 10.0%<br>(57;618) | 17.1%<br>(19;157) | **7.0%<br>(31;415)<br>Worse** | 29.1%<br>(7;46) | 3.7%<br>(14;443) | **4.7%<br>(30;623)<br>Worse** | 2.7%<br>(6;222) | **11.6%<br>(58;623)<br>Worse** |
| | Loma Linda University Medical Center | 6.3%<br>(43;614) | 7.2%<br>(63;583) | 17.8%<br>(37;186) | 4.3%<br>(18;338) | 13.4%<br>(8;59) | 2.9%<br>(8;269) | **4.3%<br>(36;921)<br>Worse** | 0.0%<br>(0;119) | 4.4%<br>(16;383) |
| | Montclair Hospital Medical Center | 8.8%<br>(1;9) | | | | | 1.0%<br>(1;48) | 2.0%<br>(2;91) | 0.0%<br>(0;7) | 4.5%<br>(13;257) |
| | Mountains Community Hospital | | | | | | | 0.0%<br>(0;4) | | 0.0%<br>(0;28) |
| | Redlands Community Hospital | 10.7%<br>(5;32) | **14.6%<br>(36;246)<br>Worse** | **35.9%<br>(16;44)<br>Worse** | **8.6%<br>(16;186)<br>Worse** | 26.1%<br>(4;16) | 2.9%<br>(6;234) | 3.5%<br>(10;276) | 2.6%<br>(3;138) | **9.1%<br>(49;472)<br>Worse** |
| | Saint Bernardine Medical Center | 4.7%<br>(29;544) | **5.8%<br>(25;374)<br>Better** | 16.0%<br>(16;87) | **2.0%<br>(5;275)<br>Better** | 18.1%<br>(4;12) | 2.4%<br>(6;276) | 1.9%<br>(12;716) | 3.7%<br>(2;65) | 5.8%<br>(42;741) |

**Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 2021, and HCAI hospital patient discharge data, 2020. See Technical Note for details about methodology.**

****Hospital name changed from 2019 to 2020, see Technical Note for details.**

**Note: Blank cells indicate that fewer than three conditions were treated**

**Definition of Terms: Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. Total Deaths: number of patients that died in this hospital. Total Cases: number of patients with this medical condition in this hospital.**

**Quality Ratings are risk-adjusted using age, gender, APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.**

# *California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators (Conditions), 2020*

HCAI Department of Health Care Access and Information

| County | Hospital Name | Acute Myocardial Infarction | Acute Stroke | | | | Gastro-Intestinal Hemorrhage | Heart Failure | Hip Fracture | Pneumonia |
|---|---|---|---|---|---|---|---|---|---|---|
| | | | Total | Hemorrhagic | Ischemic | Subarachnoid | | | | |
| | | Note: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating. | | | | | | | | |
| | STATEWIDE | 5.9%<br>(2,660;44,767) | 8.5%<br>(5,089;59,573) | 19.9%<br>(2,341;11,790) | 4.8%<br>(2,170;45,165) | 22.1%<br>(578;2,618) | 2.8%<br>(1,415;51,109) | 2.9%<br>(2,795;97,448) | 1.5%<br>(385;25,657) | 6.3%<br>(6,055;95,457) |
| San Bernardino (continued) | Saint Mary Medical Center – Apple Valley | 6.5%<br>(21;347) | 7.3%<br>(7;238) | 59.0%<br>(2;5) | 3.2%<br>(5;233) | | 0.6%<br>(1;273) | 2.0%<br>(11;616) | 0.0%<br>(0;89) | 8.0%<br>(40;536) |
| | San Antonio Regional Hospital | 6.4%<br>(33;578) | 10.8%<br>(35;432) | 21.8%<br>(17;84) | 6.8%<br>(16;340) | 45.8%<br>(2;8) | 3.2%<br>(11;342) | 2.8%<br>(16;599) | 0.8%<br>(1;160) | 5.4%<br>(37;888) |
| | Victor Valley Global Medical Center | 9.2%<br>(6;40) | 9.8%<br>(2;37) | | 5.5%<br>(2;37) | | 1.6%<br>(1;82) | 1.0%<br>(1;155) | 6.7%<br>(2;28) | 9.2%<br>(21;265) |
| San Diego | Alvarado Hospital | 5.0%<br>(5;124) | 7.7%<br>(3;58) | 20.2%<br>(1;7) | 5.4%<br>(2;50) | | 0.7%<br>(1;100) | 2.5%<br>(6;189) | 3.1%<br>(1;23) | **3.5%<br>(11;283)<br>Better** |
| | Grossmont Hospital | 6.9%<br>(19;324) | **11.5%<br>(85;687)<br>Worse** | 24.9%<br>(35;110) | **6.8%<br>(36;539)<br>Worse** | 32.0%<br>(14;38) | 2.1%<br>(10;385) | 2.8%<br>(13;692) | 1.9%<br>(3;164) | 7.3%<br>(47;820) |
| | Kaiser Foundation Hospital – San Diego – Zion | 8.6%<br>(9;104) | 11.2%<br>(43;455) | 17.6%<br>(14;105) | **7.9%<br>(24;326)<br>Worse** | 46.5%<br>(5;24) | 1.0%<br>(2;268) | 3.4%<br>(24;778) | 2.8%<br>(11;448) | 5.1%<br>(28;609) |
| | Palomar Health Downtown Campus | 6.7%<br>(28;408) | **14.8%<br>(81;543)<br>Worse** | **27.4%<br>(32;124)<br>Worse** | **10.4%<br>(43;404)<br>Worse** | 34.4%<br>(6;15) | 3.1%<br>(12;442) | **4.6%<br>(28;613)<br>Worse** | 2.1%<br>(5;244) | **8.5%<br>(43;656)<br>Worse** |
| | Palomar Medical Center Poway** | 7.6%<br>(2;8) | **36.8%<br>(9;56)<br>Worse** | 74.1%<br>(1;4) | **20.1%<br>(7;51)<br>Worse** | | **8.0%<br>(8;97)<br>Worse** | 2.7%<br>(3;127) | 4.7%<br>(3;65) | 5.8%<br>(11;216) |
| | Paradise Valley Hospital | 3.0%<br>(1;41) | 9.7%<br>(4;59) | 25.6%<br>(3;14) | 4.0%<br>(1;45) | | 0.0%<br>(0;79) | 1.2%<br>(2;201) | 0.0%<br>(0;17) | 4.1%<br>(12;266) |
| | Scripps Green Hospital | 12.0%<br>(2;6) | 7.1%<br>(4;86) | 10.7%<br>(1;16) | 3.5%<br>(2;68) | | 1.3%<br>(2;155) | 0.0%<br>(0;77) | 0.0%<br>(0;74) | 6.4%<br>(9;150) |
| | Scripps Memorial Hospital – Encinitas | 2.5%<br>(4;190) | **3.5%<br>(6;243)<br>Better** | 8.8%<br>(2;34) | **1.0%<br>(2;206)<br>Better** | 42.1%<br>(2;3) | 1.0%<br>(2;279) | **1.2%<br>(6;431)<br>Better** | 0.5%<br>(1;195) | **3.8%<br>(14;381)<br>Better** |
| | Scripps Memorial Hospital – La Jolla | 4.5%<br>(33;749) | **10.6%<br>(74;526)<br>Worse** | 21.0%<br>(25;112) | 6.1%<br>(30;357) | **34.1%<br>(19;57)<br>Worse** | 1.3%<br>(4;306) | 2.8%<br>(27;709) | 2.5%<br>(5;215) | 7.6%<br>(25;360) |

**Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 2021, and HCAI hospital patient discharge data, 2020. See Technical Note for details about methodology.**

****Hospital name changed from 2019 to 2020, see Technical Note for details.**

**Note: Blank cells indicate that fewer than three conditions were treated**

**Definition of Terms: Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. Total Deaths: number of patients that died in this hospital. Total Cases: number of patients with this medical condition in this hospital.**

**Quality Ratings are risk-adjusted using age, gender, APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.**

# *California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators (Conditions), 2020*

HCAI Department of Health Care Access and Information

| County | Hospital Name | Acute Myocardial Infarction | Acute Stroke | | | | Gastro-Intestinal Hemorrhage | Heart Failure | Hip Fracture | Pneumonia |
|---|---|---|---|---|---|---|---|---|---|---|
| | | | Total | Hemorrhagic | Ischemic | Subarachnoid | | | | |
| | | Note: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating. | | | | | | | | |
| | STATEWIDE | 5.9%<br>(2,660;44,767) | 8.5%<br>(5,089;59,573) | 19.9%<br>(2,341;11,790) | 4.8%<br>(2,170;45,165) | 22.1%<br>(578;2,618) | 2.8%<br>(1,415;51,109) | 2.9%<br>(2,795;97,448) | 1.5%<br>(385;25,657) | 6.3%<br>(6,055;95,457) |
| San Diego (continued) | Scripps Mercy Hospital | 4.3%<br>(17;336) | 7.7%<br>(47;557) | 21.9%<br>(32;135) | **2.5%<br>(10;413)<br>Better** | 27.6%<br>(5;9) | 2.2%<br>(15;516) | 2.1%<br>(23;958) | 0.8%<br>(2;250) | 6.1%<br>(51;843) |
| | Sharp Chula Vista Medical Center | 8.1%<br>(22;265) | 10.3%<br>(25;304) | 25.3%<br>(14;52) | 5.4%<br>(8;244) | 25.7%<br>(3;8) | 2.7%<br>(8;294) | 1.6%<br>(7;601) | 1.1%<br>(1;87) | 7.0%<br>(29;540) |
| | Sharp Coronado Hospital and Healthcare Center | 0.0%<br>(0;3) | 15.0%<br>(2;24) | | 5.1%<br>(1;23) | | 7.5%<br>(3;54) | 0.0%<br>(0;38) | 5.7%<br>(1;24) | 7.4%<br>(7;91) |
| | Sharp Memorial Hospital | 6.1%<br>(17;357) | **11.3%<br>(46;525)<br>Worse** | 23.2%<br>(17;84) | 6.8%<br>(26;435) | 36.6%<br>(3;6) | 2.4%<br>(9;400) | 2.2%<br>(14;661) | 1.6%<br>(6;280) | 5.4%<br>(33;676) |
| | Tri-City Medical Center – Oceanside | 7.9%<br>(17;208) | **13.4%<br>(41;283)<br>Worse** | **33.7%<br>(22;55)<br>Worse** | 5.9%<br>(13;222) | **51.1%<br>(6;6)<br>Worse** | 3.6%<br>(8;185) | 2.2%<br>(8;287) | 0.6%<br>(1;123) | 6.0%<br>(29;455) |
| | UC San Diego Health System – Hillcrest Medical Center | 3.9%<br>(13;320) | **4.8%<br>(31;518)<br>Better** | 14.3%<br>(16;113) | **2.1%<br>(10;355)<br>Better** | **11.5%<br>(5;50)<br>Better** | **1.5%<br>(10;503)<br>Better** | 1.9%<br>(18;991) | 1.2%<br>(2;174) | **3.5%<br>(30;858)<br>Better** |
| San Francisco | California Pacific Medical Center – Davies Campus Hospital | | 9.2%<br>(40;406) | 22.3%<br>(19;103) | 6.4%<br>(20;264) | **4.4%<br>(1;39)<br>Better** | 0.0%<br>(0;44) | 6.6%<br>(3;57) | 0.0%<br>(0;6) | 12.4%<br>(7;64) |
| | California Pacific Medical Center – Mission Bernal Campus | 0.0%<br>(0;8) | 11.8%<br>(2;26) | 0.0%<br>(0;5) | 10.6%<br>(2;21) | | 0.0%<br>(0;93) | 1.0%<br>(2;213) | 0.9%<br>(1;111) | 3.4%<br>(5;170) |
| | California Pacific Medical Center – Van Ness Campus | 4.3%<br>(12;180) | 6.4%<br>(11;174) | 19.7%<br>(8;40) | 1.6%<br>(2;130) | 55.0%<br>(1;4) | 2.5%<br>(12;260) | 3.4%<br>(25;448) | 0.0%<br>(0;9) | **9.2%<br>(25;205)<br>Worse** |
| | Chinese Hospital | 12.1%<br>(1;5) | 0.0%<br>(0;44) | | 0.0%<br>(0;43) | | 2.2%<br>(1;49) | 2.1%<br>(3;101) | 3.9%<br>(1;25) | 9.6%<br>(10;94) |
| | Kaiser Foundation Hospital – San Francisco | 4.4%<br>(14;480) | 12.7%<br>(18;201) | 30.0%<br>(8;37) | 6.7%<br>(9;161) | 58.6%<br>(1;3) | 1.7%<br>(5;216) | 3.4%<br>(16;422) | 1.0%<br>(1;79) | 8.8%<br>(19;210) |
| | Laguna Honda Hospital and Rehabilitation Center | | | | | | | | | 6.1%<br>(2;33) |
| | Priscilla Chan & Mark Zuckerberg San Francisco General Hospital & Trauma Center | 6.3%<br>(8;110) | **12.0%<br>(59;366)<br>Worse** | **27.4%<br>(27;94)<br>Worse** | 7.1%<br>(22;251) | 28.5%<br>(10;21) | **6.5%<br>(9;184)<br>Worse** | 3.3%<br>(11;563) | 2.2%<br>(3;125) | 7.2%<br>(19;365) |

**Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 2021, and HCAI hospital patient discharge data, 2020. See Technical Note for details about methodology.**

****Hospital name changed from 2019 to 2020, see Technical Note for details.**

**Note: Blank cells indicate that fewer than three conditions were treated**

**Definition of Terms: Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. Total Deaths: number of patients that died in this hospital. Total Cases: number of patients with this medical condition in this hospital.**

**Quality Ratings are risk-adjusted using age, gender, APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.**

# *California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators (Conditions), 2020*

HCAI Department of Health Care Access and Information

| County | Hospital Name | Acute Myocardial Infarction | Acute Stroke | | | | Gastro-Intestinal Hemorrhage | Heart Failure | Hip Fracture | Pneumonia |
|---|---|---|---|---|---|---|---|---|---|---|
| | | | Total | Hemorrhagic | Ischemic | Subarachnoid | | | | |
| | | Note: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating. | | | | | | | | |
| | STATEWIDE | 5.9%<br>(2,660;44,767) | 8.5%<br>(5,089;59,573) | 19.9%<br>(2,341;11,790) | 4.8%<br>(2,170;45,165) | 22.1%<br>(578;2,618) | 2.8%<br>(1,415;51,109) | 2.9%<br>(2,795;97,448) | 1.5%<br>(385;25,657) | 6.3%<br>(6,055;95,457) |
| San Francisco (continued) | Saint Francis Memorial Hospital | 0.0%<br>(0;7) | 7.7%<br>(4;48) | 20.6%<br>(3;17) | 3.2%<br>(1;30) | | 0.0%<br>(0;53) | 2.8%<br>(3;118) | 3.3%<br>(1;18) | **1.9%<br>(3;213)<br>Better** |
| | St. Mary's Medical Center** | 6.6%<br>(9;68) | 9.1%<br>(4;66) | 22.1%<br>(2;7) | 4.9%<br>(2;59) | | 2.6%<br>(3;89) | **0.9%<br>(3;198)<br>Better** | 1.7%<br>(1;66) | **2.5%<br>(5;149)<br>Better** |
| | UC San Francisco Medical Center | 6.1%<br>(14;165) | **6.6%<br>(52;453)<br>Better** | **14.2%<br>(23;133)<br>Better** | 4.1%<br>(18;264) | 17.7%<br>(11;56) | 1.5%<br>(7;299) | 2.3%<br>(17;542) | 0.0%<br>(0;76) | 5.2%<br>(38;528) |
| San Joaquin | Adventist Health Lodi Memorial | 9.3%<br>(5;34) | 7.5%<br>(6;141) | 23.0%<br>(2;7) | 3.8%<br>(4;134) | | 2.8%<br>(4;175) | 2.8%<br>(8;280) | 1.2%<br>(1;72) | 5.0%<br>(13;293) |
| | Dameron Hospital | **17.1%<br>(13;104)<br>Worse** | 20.1%<br>(4;52) | | 11.3%<br>(4;52) | | 4.1%<br>(3;86) | 2.0%<br>(3;237) | 4.7%<br>(1;27) | 6.6%<br>(12;263) |
| | Doctors Hospital of Manteca | 5.4%<br>(1;23) | 7.7%<br>(2;63) | 0.0%<br>(0;4) | 7.0%<br>(2;59) | | 4.5%<br>(2;103) | 1.0%<br>(1;164) | 0.0%<br>(0;29) | **13.5%<br>(14;165)<br>Worse** |
| | Kaiser Foundation Hospital – Manteca | 5.3%<br>(4;96) | 7.2%<br>(6;161) | 15.9%<br>(2;20) | 4.1%<br>(4;141) | | 3.8%<br>(6;175) | 3.6%<br>(13;426) | 2.3%<br>(3;153) | 6.4%<br>(24;462) |
| | Saint Joseph's Medical Center of Stockton | 5.2%<br>(37;693) | 6.3%<br>(18;326) | 19.3%<br>(10;23) | **2.5%<br>(7;302)<br>Better** | | 1.9%<br>(11;421) | **1.4%<br>(14;911)<br>Better** | 0.0%<br>(0;67) | **2.5%<br>(18;626)<br>Better** |
| | San Joaquin General Hospital | 5.6%<br>(2;31) | 10.8%<br>(19;157) | 28.7%<br>(15;43) | 3.3%<br>(3;111) | 97.1%<br>(1;3) | 3.0%<br>(5;120) | 2.2%<br>(4;300) | 1.1%<br>(1;113) | **11.6%<br>(19;282)<br>Worse** |
| | Sutter Tracy Community Hospital | 0.0%<br>(0;15) | 0.0%<br>(0;66) | | 0.0%<br>(0;64) | | 4.3%<br>(3;88) | 2.4%<br>(5;136) | 4.2%<br>(1;24) | 5.6%<br>(11;211) |
| San Luis Obispo | French Hospital Medical Center | 7.1%<br>(15;254) | 5.0%<br>(2;50) | 35.9%<br>(2;3) | 0.0%<br>(0;47) | | 2.5%<br>(2;100) | 3.3%<br>(11;235) | 5.3%<br>(3;46) | 6.2%<br>(12;161) |
| | Marian Regional Medical Center – Arroyo Grande | 0.0%<br>(0;5) | 5.8%<br>(2;47) | 27.2%<br>(2;6) | 0.0%<br>(0;41) | | 0.0%<br>(0;59) | 4.3%<br>(3;77) | 0.0%<br>(0;59) | 5.9%<br>(5;102) |

**Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 2021, and HCAI hospital patient discharge data, 2020. See Technical Note for details about methodology.**

****Hospital name changed from 2019 to 2020, see Technical Note for details.**

**Note: Blank cells indicate that fewer than three conditions were treated**

**Definition of Terms: Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. Total Deaths: number of patients that died in this hospital. Total Cases: number of patients with this medical condition in this hospital.**

**Quality Ratings are risk-adjusted using age, gender, APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.**

# *California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators (Conditions), 2020*

HCAI Department of Health Care Access and Information

| County | Hospital Name | Acute Myocardial Infarction | Acute Stroke | | | | Gastro-Intestinal Hemorrhage | Heart Failure | Hip Fracture | Pneumonia |
|---|---|---|---|---|---|---|---|---|---|---|
| | | | Total | Hemorrhagic | Ischemic | Subarachnoid | | | | |
| | | Note: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating. | | | | | | | | |
| | STATEWIDE | 5.9%<br>(2,660;44,767) | 8.5%<br>(5,089;59,573) | 19.9%<br>(2,341;11,790) | 4.8%<br>(2,170;45,165) | 22.1%<br>(578;2,618) | 2.8%<br>(1,415;51,109) | 2.9%<br>(2,795;97,448) | 1.5%<br>(385;25,657) | 6.3%<br>(6,055;95,457) |
| San Luis Obispo (continued) | Sierra Vista Regional Medical Center | 19.8%<br>(1;3) | 7.6%<br>(16;156) | **9.6%<br>(5;54)<br>Better** | 7.2%<br>(11;100) | | 2.2%<br>(2;96) | 5.3%<br>(5;88) | 0.0%<br>(0;58) | 5.2%<br>(7;149) |
| | Tenet Health Central Coast Twin Cities Community Hospital** | 0.0%<br>(0;12) | 8.0%<br>(4;88) | | 4.0%<br>(3;85) | | 2.9%<br>(3;104) | 2.6%<br>(5;166) | 0.0%<br>(0;80) | 8.9%<br>(17;204) |
| San Mateo | Kaiser Foundation Hospital – Redwood City | 2.6%<br>(1;44) | **13.5%<br>(75;571)<br>Worse** | 22.9%<br>(30;202) | **10.7%<br>(30;282)<br>Worse** | **41.5%<br>(15;87)<br>Worse** | 4.1%<br>(3;89) | 4.5%<br>(8;197) | 1.5%<br>(1;70) | 8.3%<br>(10;131) |
| | Kaiser Foundation Hospital – South San Francisco | 5.7%<br>(2;33) | 7.8%<br>(11;198) | 20.7%<br>(5;50) | 2.9%<br>(4;141) | 57.6%<br>(2;7) | 3.0%<br>(3;154) | 3.2%<br>(14;387) | 1.3%<br>(1;64) | 5.4%<br>(21;330) |
| | Peninsula Medical Center | 4.3%<br>(11;227) | 7.3%<br>(32;354) | 16.9%<br>(15;59) | 3.7%<br>(12;285) | 26.1%<br>(5;10) | **0.7%<br>(2;269)<br>Better** | 3.1%<br>(17;400) | 2.0%<br>(3;145) | 7.3%<br>(11;158) |
| | San Mateo Medical Center | 16.2%<br>(1;3) | 14.7%<br>(1;28) | 0.0%<br>(0;3) | 10.1%<br>(1;24) | | 0.0%<br>(0;41) | 1.9%<br>(1;83) | 0.0%<br>(0;7) | **18.1%<br>(11;102)<br>Worse** |
| | Sequoia Hospital | 1.5%<br>(1;55) | 8.8%<br>(5;84) | 31.3%<br>(4;17) | 2.2%<br>(1;66) | | 2.1%<br>(1;72) | 3.2%<br>(6;164) | 2.3%<br>(1;55) | 3.7%<br>(4;130) |
| | Seton Medical Center | 8.3%<br>(14;154) | 6.2%<br>(7;103) | 13.9%<br>(4;27) | 3.6%<br>(3;76) | | 2.2%<br>(4;115) | 2.7%<br>(7;201) | 2.7%<br>(1;37) | 5.2%<br>(9;151) |
| Santa Barbara | Goleta Valley Cottage Hospital | | | | | | | 9.3%<br>(1;13) | 0.0%<br>(0;54) | 10.0%<br>(2;41) |
| | Lompoc Valley Medical Center | | 0.0%<br>(0;40) | 0.0%<br>(0;4) | 0.0%<br>(0;36) | | 2.6%<br>(1;83) | 7.2%<br>(5;90) | 0.0%<br>(0;24) | **15.7%<br>(8;72)<br>Worse** |
| | Marian Regional Medical Center | 7.3%<br>(11;162) | 9.2%<br>(15;166) | 24.1%<br>(11;56) | 4.0%<br>(4;108) | | 2.9%<br>(7;180) | 4.1%<br>(14;350) | 4.1%<br>(4;121) | **3.1%<br>(9;274)<br>Better** |
| | Santa Barbara Cottage Hospital | 5.8%<br>(13;252) | 8.0%<br>(66;533) | 17.6%<br>(27;110) | 5.4%<br>(29;363) | 14.9%<br>(10;60) | 2.2%<br>(8;336) | 3.1%<br>(16;427) | 0.0%<br>(0;173) | 8.1%<br>(32;403) |
| | Santa Ynez Valley Cottage Hospital | | 0.0%<br>(0;3) | | 0.0%<br>(0;3) | | | 0.0%<br>(0;7) | | 21.6%<br>(2;14) |

**Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 2021, and HCAI hospital patient discharge data, 2020. See Technical Note for details about methodology.**

****Hospital name changed from 2019 to 2020, see Technical Note for details.**

**Note: Blank cells indicate that fewer than three conditions were treated**

**Definition of Terms: Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. Total Deaths: number of patients that died in this hospital. Total Cases: number of patients with this medical condition in this hospital.**

**Quality Ratings are risk-adjusted using age, gender, APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.**

# *California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators (Conditions), 2020*

HCAI Department of Health Care Access and Information

| County | Hospital Name | Acute Myocardial Infarction | Acute Stroke | | | | Gastro-Intestinal Hemorrhage | Heart Failure | Hip Fracture | Pneumonia |
|---|---|---|---|---|---|---|---|---|---|---|
| | | | Total | Hemorrhagic | Ischemic | Subarachnoid | | | | |
| | | Note: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating. | | | | | | | | |
| | STATEWIDE | 5.9%<br>(2,660;44,767) | 8.5%<br>(5,089;59,573) | 19.9%<br>(2,341;11,790) | 4.8%<br>(2,170;45,165) | 22.1%<br>(578;2,618) | 2.8%<br>(1,415;51,109) | 2.9%<br>(2,795;97,448) | 1.5%<br>(385;25,657) | 6.3%<br>(6,055;95,457) |
| Santa Clara | El Camino Health | 7.7%<br>(15;207) | 8.7%<br>(21;343) | 21.0%<br>(10;46) | 4.7%<br>(9;292) | 23.9%<br>(2;5) | 3.6%<br>(11;349) | 2.7%<br>(15;532) | 0.5%<br>(1;177) | 6.7%<br>(28;446) |
| | Good Samaritan Hospital – San Jose | 5.9%<br>(19;279) | **5.1%<br>(28;489)<br>Better** | 13.7%<br>(14;101) | **2.0%<br>(8;367)<br>Better** | 19.2%<br>(6;21) | 2.7%<br>(7;187) | **1.0%<br>(5;429)<br>Better** | 1.4%<br>(2;164) | 5.4%<br>(28;428) |
| | Kaiser Foundation Hospital – San Jose | 5.0%<br>(10;190) | 7.9%<br>(11;254) | 23.5%<br>(5;42) | 3.3%<br>(4;204) | 24.8%<br>(2;8) | 2.1%<br>(3;224) | 3.9%<br>(21;432) | 0.6%<br>(1;125) | 6.6%<br>(22;373) |
| | Kaiser Foundation Hospital – Santa Clara | 6.1%<br>(34;649) | 11.0%<br>(20;247) | **36.4%<br>(11;50)<br>Worse** | 4.8%<br>(9;192) | 0.0%<br>(0;5) | 3.6%<br>(10;306) | 3.0%<br>(26;707) | 2.3%<br>(4;156) | 5.5%<br>(24;390) |
| | O'Connor Hospital | 5.0%<br>(10;174) | 6.4%<br>(5;135) | 15.5%<br>(3;33) | 1.8%<br>(1;99) | 85.0%<br>(1;3) | 2.5%<br>(5;216) | 3.5%<br>(16;486) | 0.0%<br>(0;74) | 8.4%<br>(30;351) |
| | Regional Medical Center of San Jose | **4.0%<br>(16;258)<br>Better** | **4.9%<br>(46;611)<br>Better** | **11.3%<br>(18;130)<br>Better** | **2.6%<br>(18;442)<br>Better** | **14.3%<br>(10;39)<br>Better** | 2.8%<br>(10;251) | **1.2%<br>(7;588)<br>Better** | 3.8%<br>(3;94) | **2.7%<br>(20;583)<br>Better** |
| | Saint Louise Regional Hospital | 2.3%<br>(1;12) | 0.0%<br>(0;66) | 0.0%<br>(0;4) | 0.0%<br>(0;62) | | 1.4%<br>(1;91) | 2.1%<br>(2;133) | 0.0%<br>(0;32) | 2.9%<br>(4;195) |
| | Santa Clara Valley Medical Center | 4.0%<br>(5;142) | 9.8%<br>(17;263) | 26.1%<br>(11;66) | 3.4%<br>(3;186) | 31.0%<br>(3;11) | 2.8%<br>(8;297) | 2.8%<br>(14;602) | 0.0%<br>(0;59) | 8.0%<br>(20;317) |
| | Stanford Health Care | 5.6%<br>(19;261) | 8.2%<br>(46;535) | 13.0%<br>(10;123) | 5.0%<br>(28;356) | 30.4%<br>(8;56) | 2.2%<br>(13;395) | 3.1%<br>(33;663) | 0.9%<br>(2;191) | 5.7%<br>(37;501) |
| Santa Cruz | Dominican Hospital | 3.9%<br>(7;194) | 10.2%<br>(20;239) | 14.5%<br>(6;58) | 7.3%<br>(11;178) | **51.5%<br>(3;3)<br>Worse** | **0.0%<br>(0;207)<br>Better** | 3.9%<br>(12;334) | 1.8%<br>(2;136) | 4.3%<br>(9;275) |
| | Watsonville Community Hospital | 35.8%<br>(1;3) | 6.9%<br>(3;53) | 27.1%<br>(2;4) | 0.0%<br>(0;48) | | 5.3%<br>(3;72) | 1.1%<br>(1;111) | 0.0%<br>(0;39) | 9.1%<br>(10;161) |
| Shasta | Mayers Memorial Hospital | | | | | | | 0.0%<br>(0;5) | | 26.2%<br>(1;17) |
| | Mercy Medical Center – Redding | 4.7%<br>(15;304) | **14.8%<br>(69;490)<br>Worse** | 26.9%<br>(21;78) | **10.5%<br>(43;394)<br>Worse** | 23.3%<br>(5;18) | 3.9%<br>(9;264) | **6.0%<br>(26;354)<br>Worse** | 3.6%<br>(7;179) | **10.1%<br>(47;465)<br>Worse** |

**Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 2021, and HCAI hospital patient discharge data, 2020. See Technical Note for details about methodology.**

****Hospital name changed from 2019 to 2020, see Technical Note for details.**

**Note: Blank cells indicate that fewer than three conditions were treated**

**Definition of Terms: Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. Total Deaths: number of patients that died in this hospital. Total Cases: number of patients with this medical condition in this hospital.**

**Quality Ratings are risk-adjusted using age, gender, APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.**

# *California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators (Conditions), 2020*

HCAI Department of Health Care Access and Information

| County | Hospital Name | Acute Myocardial Infarction | Acute Stroke | | | | Gastro-Intestinal Hemorrhage | Heart Failure | Hip Fracture | Pneumonia |
|---|---|---|---|---|---|---|---|---|---|---|
| | | | Total | Hemorrhagic | Ischemic | Subarachnoid | | | | |
| | | Note: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating. | | | | | | | | |
| | STATEWIDE | 5.9%<br>(2,660;44,767) | 8.5%<br>(5,089;59,573) | 19.9%<br>(2,341;11,790) | 4.8%<br>(2,170;45,165) | 22.1%<br>(578;2,618) | 2.8%<br>(1,415;51,109) | 2.9%<br>(2,795;97,448) | 1.5%<br>(385;25,657) | 6.3%<br>(6,055;95,457) |
| Shasta (continued) | Shasta Regional Medical Center | 8.7%<br>(16;218) | 9.1%<br>(4;127) | | 4.1%<br>(3;126) | | 2.5%<br>(4;223) | 2.7%<br>(11;451) | 0.0%<br>(0;75) | 7.9%<br>(31;400) |
| Siskiyou | Fairchild Medical Center | 14.3%<br>(1;7) | 0.0%<br>(0;33) | | 0.0%<br>(0;31) | | 6.1%<br>(1;23) | 6.8%<br>(2;46) | 0.0%<br>(0;29) | 5.2%<br>(2;70) |
| | Mercy Medical Center Mt. Shasta | | 10.6%<br>(1;14) | | 7.6%<br>(1;13) | | 0.0%<br>(0;13) | 11.4%<br>(2;23) | 0.0%<br>(0;17) | 10.7%<br>(3;55) |
| Solano | Kaiser Foundation Hospital & Rehabilitation Center – Vallejo | 3.7%<br>(8;257) | **19.7%<br>(22;166)<br>Worse** | **56.7%<br>(10;36)<br>Worse** | **9.3%<br>(11;126)<br>Worse** | 56.9%<br>(1;4) | 3.2%<br>(6;165) | 4.6%<br>(19;351) | 1.8%<br>(2;103) | 5.0%<br>(10;235) |
| | Kaiser Foundation Hospital – Vacaville | 5.2%<br>(3;40) | 11.2%<br>(15;143) | 24.5%<br>(9;51) | 7.0%<br>(6;92) | | 2.4%<br>(4;116) | **7.0%<br>(19;255)<br>Worse** | 2.2%<br>(2;86) | 8.4%<br>(13;209) |
| | NorthBay Medical Center | 7.5%<br>(19;252) | 10.4%<br>(24;248) | 28.8%<br>(14;42) | 5.1%<br>(10;203) | 0.0%<br>(0;3) | **5.7%<br>(12;214)<br>Worse** | 3.9%<br>(14;398) | 1.7%<br>(2;93) | 7.2%<br>(36;556) |
| | Sutter Solano Medical Center | 15.8%<br>(3;15) | 0.0%<br>(0;83) | 0.0%<br>(0;6) | 0.0%<br>(0;77) | | 3.6%<br>(3;109) | 5.0%<br>(6;193) | 5.6%<br>(1;27) | 2.7%<br>(3;152) |
| Sonoma | Healdsburg District Hospital | | 53.1%<br>(2;15) | | **39.6%<br>(2;14)<br>Worse** | | 0.0%<br>(0;3) | 0.0%<br>(0;20) | 0.0%<br>(0;10) | 11.8%<br>(4;51) |
| | Kaiser Foundation Hospital – Santa Rosa | 7.6%<br>(4;56) | 12.9%<br>(15;172) | 37.9%<br>(7;33) | 6.3%<br>(7;134) | 23.0%<br>(1;5) | 1.9%<br>(3;235) | 2.1%<br>(7;329) | 0.7%<br>(1;152) | 3.6%<br>(10;359) |
| | Petaluma Valley Hospital | 15.7%<br>(2;5) | 23.2%<br>(4;29) | | 11.1%<br>(3;27) | | 11.6%<br>(1;7) | 2.5%<br>(2;81) | 2.7%<br>(1;36) | 9.4%<br>(7;93) |
| | Santa Rosa Memorial Hospital – Montgomery | 7.2%<br>(19;317) | **11.9%<br>(39;331)<br>Worse** | 28.4%<br>(19;85) | 6.8%<br>(19;237) | 18.4%<br>(1;9) | 3.9%<br>(10;285) | **5.5%<br>(23;406)<br>Worse** | 1.3%<br>(2;157) | **11.5%<br>(36;356)<br>Worse** |
| | Sonoma Specialty Hospital | | | | | | | | | 12.8%<br>(2;4) |
| | Sonoma Valley Hospital | | 0.0%<br>(0;17) | | 0.0%<br>(0;17) | | 0.0%<br>(0;5) | 1.5%<br>(1;30) | 0.0%<br>(0;30) | 5.8%<br>(2;30) |

**Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 2021, and HCAI hospital patient discharge data, 2020. See Technical Note for details about methodology.**

****Hospital name changed from 2019 to 2020, see Technical Note for details.**

**Note: Blank cells indicate that fewer than three conditions were treated**

**Definition of Terms: Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. Total Deaths: number of patients that died in this hospital. Total Cases: number of patients with this medical condition in this hospital.**

**Quality Ratings are risk-adjusted using age, gender, APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.**

# *California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators (Conditions), 2020*

HCAI Department of Health Care Access and Information

| County | Hospital Name | Acute Myocardial Infarction | Acute Stroke | | | | Gastro-Intestinal Hemorrhage | Heart Failure | Hip Fracture | Pneumonia |
|---|---|---|---|---|---|---|---|---|---|---|
| | | | Total | Hemorrhagic | Ischemic | Subarachnoid | | | | |
| | | Note: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating. | | | | | | | | |
| | STATEWIDE | 5.9%<br>(2,660;44,767) | 8.5%<br>(5,089;59,573) | 19.9%<br>(2,341;11,790) | 4.8%<br>(2,170;45,165) | 22.1%<br>(578;2,618) | 2.8%<br>(1,415;51,109) | 2.9%<br>(2,795;97,448) | 1.5%<br>(385;25,657) | 6.3%<br>(6,055;95,457) |
| Sonoma (continued) | Sutter Santa Rosa Regional Hospital | 3.8%<br>(7;193) | 11.2%<br>(7;97) | 19.6%<br>(1;5) | 7.0%<br>(5;90) | | 3.0%<br>(5;179) | 2.2%<br>(4;189) | 2.4%<br>(1;59) | 7.6%<br>(8;143) |
| Stanislaus | Doctors Medical Center | 7.4%<br>(43;460) | **7.0%<br>(87;806)<br>Better** | 17.0%<br>(37;181) | 4.1%<br>(40;567) | **13.8%<br>(10;58)<br>Better** | 2.4%<br>(9;392) | **4.9%<br>(35;701)<br>Worse** | 1.4%<br>(2;160) | **8.3%<br>(57;721)<br>Worse** |
| | Emanuel Medical Center | 4.2%<br>(15;356) | 7.2%<br>(4;65) | 40.0%<br>(3;6) | 1.5%<br>(1;59) | | 2.0%<br>(2;159) | 2.7%<br>(12;450) | 0.0%<br>(0;53) | 6.7%<br>(33;532) |
| | Memorial Medical Center – Modesto | 7.6%<br>(30;314) | 6.4%<br>(20;381) | 16.8%<br>(13;90) | 2.8%<br>(6;289) | | 2.7%<br>(12;520) | 2.8%<br>(21;754) | 1.1%<br>(2;200) | **9.1%<br>(34;536)<br>Worse** |
| | Oak Valley District Hospital | | 0.0%<br>(0;12) | | 0.0%<br>(0;12) | | 13.0%<br>(1;4) | 0.0%<br>(0;34) | | 15.2%<br>(5;71) |
| Tehama | Saint Elizabeth Community Hospital | 14.9%<br>(2;8) | 3.1%<br>(1;60) | 0.0%<br>(0;3) | 2.1%<br>(1;57) | | 7.8%<br>(2;46) | 5.3%<br>(7;135) | 0.0%<br>(0;51) | 5.7%<br>(5;134) |
| Trinity | Trinity Hospital | | 51.4%<br>(1;4) | | 28.9%<br>(1;4) | | | | | 0.0%<br>(0;14) |
| Tulare | Adventist Health Tulare | 0.0%<br>(0;7) | 0.0%<br>(0;27) | | 0.0%<br>(0;25) | | | 3.3%<br>(4;105) | | 5.7%<br>(3;82) |
| | Kaweah Delta Medical Center | **8.2%<br>(39;482)<br>Worse** | 9.0%<br>(30;462) | 23.1%<br>(19;92) | 4.5%<br>(11;368) | | 3.2%<br>(17;477) | 3.9%<br>(37;1,043) | 0.4%<br>(1;217) | **9.0%<br>(52;762)<br>Worse** |
| | Sierra View Medical Center | 5.8%<br>(6;80) | 8.8%<br>(5;105) | | 5.0%<br>(5;105) | | 1.8%<br>(1;126) | 5.3%<br>(5;137) | 0.0%<br>(0;58) | **10.2%<br>(31;387)<br>Worse** |
| Tuolumne | Adventist Health Sonora – Greenley | 8.7%<br>(5;41) | 5.7%<br>(2;71) | 78.1%<br>(1;4) | 1.7%<br>(1;67) | | 3.1%<br>(3;97) | **5.8%<br>(12;173)<br>Worse** | 1.7%<br>(2;122) | **10.2%<br>(21;223)<br>Worse** |

**Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 2021, and HCAI hospital patient discharge data, 2020. See Technical Note for details about methodology.**

****Hospital name changed from 2019 to 2020, see Technical Note for details.**

**Note: Blank cells indicate that fewer than three conditions were treated**

**Definition of Terms: Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. Total Deaths: number of patients that died in this hospital. Total Cases: number of patients with this medical condition in this hospital.**

**Quality Ratings are risk-adjusted using age, gender, APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.**

# *California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators (Conditions), 2020*

HCAI Department of Health Care Access and Information

| County | Hospital Name | Acute Myocardial Infarction | Acute Stroke | | | | Gastro-Intestinal Hemorrhage | Heart Failure | Hip Fracture | Pneumonia |
|---|---|---|---|---|---|---|---|---|---|---|
| | | | Total | Hemorrhagic | Ischemic | Subarachnoid | | | | |
| | | Note: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating. | | | | | | | | |
| | STATEWIDE | 5.9%<br>(2,660;44,767) | 8.5%<br>(5,089;59,573) | 19.9%<br>(2,341;11,790) | 4.8%<br>(2,170;45,165) | 22.1%<br>(578;2,618) | 2.8%<br>(1,415;51,109) | 2.9%<br>(2,795;97,448) | 1.5%<br>(385;25,657) | 6.3%<br>(6,055;95,457) |
| Ventura | Adventist Health Simi Valley | 1.3%<br>(1;103) | 6.9%<br>(6;125) | 23.5%<br>(4;18) | 2.5%<br>(2;104) | 0.0%<br>(0;3) | 0.0%<br>(0;135) | 1.9%<br>(6;276) | 2.7%<br>(2;77) | **3.4%<br>(10;343)<br>Better** |
| | Community Memorial Hospital – San Buenaventura | 7.5%<br>(10;230) | 9.3%<br>(13;266) | 13.8%<br>(3;38) | 5.8%<br>(9;226) | | 3.8%<br>(9;262) | **1.1%<br>(4;374)<br>Better** | 3.1%<br>(5;187) | 8.1%<br>(17;315) |
| | Los Robles Hospital and Medical Center | 5.3%<br>(16;267) | 8.6%<br>(33;376) | 23.5%<br>(17;56) | 4.9%<br>(15;286) | 6.2%<br>(1;34) | 3.7%<br>(8;174) | 2.1%<br>(10;391) | 2.6%<br>(4;181) | **4.5%<br>(28;529)<br>Better** |
| | Ojai Valley Community Hospital | | | | | | | 0.0%<br>(0;5) | | 0.0%<br>(0;18) |
| | Saint John's Pleasant Valley Hospital | 8.8%<br>(2;22) | **0.0%<br>(0;66)<br>Better** | 0.0%<br>(0;6) | 0.0%<br>(0;59) | | 1.2%<br>(1;86) | 1.1%<br>(2;140) | 0.0%<br>(0;86) | 4.5%<br>(9;246) |
| | Saint John's Regional Medical Center | 3.6%<br>(7;236) | 8.0%<br>(41;361) | 14.6%<br>(15;74) | 5.3%<br>(22;275) | 25.1%<br>(4;12) | 3.7%<br>(11;320) | 2.4%<br>(10;414) | 1.7%<br>(2;86) | **3.9%<br>(15;335)<br>Better** |
| | Ventura County Medical Center | 3.3%<br>(1;42) | 7.0%<br>(3;88) | 21.0%<br>(2;13) | 2.7%<br>(1;75) | | 1.1%<br>(1;95) | 2.1%<br>(2;165) | 0.0%<br>(0;42) | 1.5%<br>(1;140) |
| Yolo | Sutter Davis Hospital | 22.8%<br>(2;8) | 5.2%<br>(1;45) | | 0.0%<br>(0;42) | | 4.5%<br>(3;82) | 0.8%<br>(1;123) | 0.0%<br>(0;46) | 9.5%<br>(5;68) |
| | Woodland Memorial Hospital | 0.0%<br>(0;10) | 21.3%<br>(4;47) | 36.5%<br>(1;3) | 13.6%<br>(3;44) | | 2.8%<br>(2;57) | 6.1%<br>(6;102) | **7.5%<br>(3;55)<br>Worse** | 4.2%<br>(3;96) |
| Yuba | Adventist Health and Rideout | 4.6%<br>(9;222) | 7.6%<br>(10;232) | 18.4%<br>(2;8) | 4.2%<br>(7;222) | | 2.9%<br>(6;202) | 3.5%<br>(18;657) | 3.7%<br>(4;113) | 7.6%<br>(32;585) |

**Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 2021, and HCAI hospital patient discharge data, 2020. See Technical Note for details about methodology.**

****Hospital name changed from 2019 to 2020, see Technical Note for details.**

**Note: Blank cells indicate that fewer than three conditions were treated**

**Definition of Terms: Risk Adjustment: The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. Total Deaths: number of patients that died in this hospital. Total Cases: number of patients with this medical condition in this hospital.**

**Quality Ratings are risk-adjusted using age, gender, APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.**