

4142 HW 5

Duncan Wilkie

28 October 2022

Problem 1. Consider the energy matrix

$$\begin{pmatrix} a & 0 & 0 & -b \\ 0 & c & id & 0 \\ 0 & -id & -c & 0 \\ -b & 0 & 0 & -a \end{pmatrix}$$

Find the eigenvalues and eigenvectors of this Hamiltonian. It will help to not approach this as a direct diagonalization but regard the elements above as coupling between four states.

Solution. Looking at what will happen when the basis vectors, it's clear this is actually is two non-interacting problems: vectors in combinations of the first and last basis vectors get mapped to each other, and likewise for the second and third. The individual problems are

$$\begin{pmatrix} a & -b \\ -b & -a \end{pmatrix} \Rightarrow \det \begin{vmatrix} a - \lambda & -b \\ -b & -a - \lambda \end{vmatrix} = 0 \Leftrightarrow \lambda^2 - a^2 - b^2 = 0 \Rightarrow \lambda = \pm \sqrt{a^2 + b^2}$$

and

$$\begin{pmatrix} c & id \\ -id & -c \end{pmatrix} \Rightarrow \det \begin{vmatrix} c - \lambda & id \\ -id & -c - \lambda \end{vmatrix} = 0 \Leftrightarrow \lambda^2 - c^2 - d^2 = 0 \Rightarrow \lambda = \pm \sqrt{c^2 + d^2}$$

□

Problem 2. At time $t = 0$, a spin-1/2 particle is in the state $|S_z = +\rangle$.

1. If S_x is measured at $t = 0$, what is the probability of getting a value $\hbar/2$?
2. Instead, with no measurement at $t = 0$, suppose the system evolves in a magnetic field $\vec{B} = B_0 \hat{e}_y$. Use the S_z basis to calculate the state of the system at time t .
3. Suppose you now measure S_x at t . What is the probability of getting the value $\hbar/2$?

Solution. In general,

$$\chi = \begin{pmatrix} a \\ b \end{pmatrix} = \frac{a+b}{\sqrt{2}} \chi_+^{(x)} + \frac{a-b}{\sqrt{2}} \chi_-^{(x)},$$

so in the state $S_z = + \Rightarrow a = 1, b = 0$, so the probability that a measurement of S_x yields $\hbar/2$ is

$$P(S_x = +) = \frac{1}{\sqrt{2}}.$$

The Hamiltonian in the presence of the magnetic field is

$$H = \gamma B_0 S_y = -\frac{\gamma B_0 \hbar}{2} \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}$$

The eigenstates of this are the usual eigenstates of S_y : in the S_z basis,

$$\chi_+^{(y)} = \begin{pmatrix} -i/\sqrt{2} \\ 1/\sqrt{2} \end{pmatrix}, \quad \chi_-^{(y)} = \begin{pmatrix} i/\sqrt{2} \\ 1/\sqrt{2} \end{pmatrix}$$

and accordingly

$$\chi = \frac{b + ia}{\sqrt{2}} \chi_+^{(y)} + \frac{b - ia}{\sqrt{2}} \chi_-^{(y)}$$

Since this is time-independent, the time-dependent solution expressed in the S_y basis is

$$\chi(t) = \begin{pmatrix} ae^{-iE_+t/\hbar} \\ be^{-iE_-t/\hbar} \end{pmatrix}$$

where a, b are specified by the initial condition (in this case, $\chi(0) = |S_z = +\rangle \Rightarrow a = \frac{i}{\sqrt{2}}, b = \frac{-i}{\sqrt{2}}$) and E_{\pm} are the energies of the $\chi_{\pm}^{(y)}$ eigenstates, which are $\pm \frac{\gamma B_0 \hbar}{2}$ by the eigenvalues of the states:

$$\chi(t) = \begin{pmatrix} \frac{i}{\sqrt{2}} e^{i\gamma B_0 t/2} \\ \frac{-i}{\sqrt{2}} e^{-i\gamma B_0 t/2} \end{pmatrix}.$$

Changing back to the S_z basis,

$$\begin{aligned} \chi(t) &= \left(\frac{i}{\sqrt{2}} e^{i\gamma B_0 t/2} \right) \left(\frac{-i}{\sqrt{2}} \chi_+ + \frac{1}{\sqrt{2}} \chi_- \right) + \left(\frac{-i}{\sqrt{2}} e^{-i\gamma B_0 t/2} \right) \left(\frac{i}{\sqrt{2}} \chi_+ + \frac{1}{\sqrt{2}} \chi_- \right) \\ &= \frac{e^{i\gamma B_0 t/2}}{2} \chi_+ + \frac{ie^{i\gamma B_0 t/2}}{2} \chi_- + \frac{e^{-i\gamma B_0 t/2}}{2} \chi_+ - \frac{ie^{-i\gamma B_0 t/2}}{2} \chi_- \\ &= \begin{pmatrix} \cos(\gamma B_0 t/2) \\ -\sin(\gamma B_0 t/2) \end{pmatrix} \end{aligned}$$

This looks a lot like a clockwise parameterization of a circle.

The change of basis from S_z to S_x is

$$\chi = \frac{a + b}{\sqrt{2}} \chi_+^{(x)} + \frac{a - b}{\sqrt{2}} \chi_-^{(x)},$$

so the coefficient of $\chi_+^{(x)}$ using the above expression for the S_z basis is

$$\frac{\cos(\gamma B_0 t/2) - \sin(\gamma B_0 t/2)}{\sqrt{2}}$$

The corresponding probability of a measurement resulting in $\chi_+^{(x)}$ (which has eigenvalue $\hbar/2$) is

$$\left(\frac{\cos(\gamma B_0 t/2) - \sin(\gamma B_0 t/2)}{\sqrt{2}} \right)^2 = \frac{1}{2} (1 - 2 \sin(\gamma B_0 t))$$

□

Problem 3. Three quarks, each of spin $1/2$, form a baryon. What are the allowed values of baryon spin?

Solution. Distinguish one pair of quarks. This combination has a spin of either 1 or 0, as those are the only integers between $\frac{1}{2} + \frac{1}{2}$ and $\frac{1}{2} - \frac{1}{2}$. The spin of the whole system is a combination of the spin of the pair with the spin of the remaining quark; it's either $1 + \frac{1}{2} \Rightarrow \frac{3}{2}, \frac{1}{2}$ or $0 + \frac{1}{2} \Rightarrow \frac{1}{2}, -\frac{1}{2}$. Accordingly, the only possibly baryon spins are $\frac{1}{2}$ and $\frac{3}{2}$. \square

Problem 4. Suppose an electron is in potentials $\vec{A} = B_0(x\hat{j} - y\hat{i})$, $\Phi = Kz^2$.

1. Find the electric and magnetic fields.
2. Find the allowed energies of the electron.

Solution. Since Φ is time-independent, $\vec{B} = \nabla \times \vec{A}$ and $\vec{E} = -\nabla\Phi$:

$$\nabla \times \vec{A} = \det \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ -B_0y & B_0x & 0 \end{vmatrix} = 2B_0\hat{k},$$

$$\nabla\Phi = 2Kz\hat{k}.$$

The minimal coupling Hamiltonian is

$$\begin{aligned} H &= \frac{1}{2m}(-i\hbar\nabla - e\vec{A})^2 + e\Phi = \frac{1}{2m}(-i\hbar\nabla + eB_0(x\hat{j} - y\hat{i}))^2 + eKz^2 \\ &= \frac{1}{2m}(i\hbar\nabla + eB_0(x\hat{j} - y\hat{i})) \cdot (i\hbar\nabla + eB_0(x\hat{j} - y\hat{i})) + eKz^2 \\ &= \frac{1}{2m} \left[-\hbar^2\nabla^2 + i\hbar eB_0(\nabla \cdot (x\hat{j} - y\hat{i}) + (x\hat{j} - y\hat{i}) \cdot \nabla) + e^2B_0^2(x^2 + y^2) \right] + eKz^2 \\ &= \frac{1}{2m} \left[-\hbar^2\nabla^2 + i\hbar eB_0 \left(x \frac{\partial}{\partial y} - y \frac{\partial}{\partial x} \right) + e^2B_0^2(x^2 + y^2) \right] + eKz^2 \\ &= \frac{1}{2m} [-\hbar^2\nabla^2 - eB_0L_z + e^2B_0^2(x^2 + y^2)] + eKz^2 \\ &= \frac{1}{2m} [-\hbar^2\nabla^2 - eB_0L_z] + \left[\frac{e^2B_0^2}{2m}\hat{i} + \frac{e^2B_0^2}{2m}\hat{j} + eK\hat{z} \right] \cdot [x^2\hat{i} + y^2\hat{j} + z^2\hat{k}] \end{aligned}$$

Applying this to a wave function ψ ,

$$H\psi = \frac{1}{2m} [-\hbar^2\nabla^2\psi - eB_0L_z\psi] + \left[\frac{e^2B_0^2}{2m}\hat{i} + \frac{e^2B_0^2}{2m}\hat{j} + eK\hat{z} \right] \cdot [x^2\hat{i} + y^2\hat{j} + z^2\hat{k}]\psi = E\psi$$

It's clear that this problem has an angular component, from the presence of L_z , and that the z -axis is distinguished, from the dot-product term. Accordingly, we move to cylindrical coordinates, in which $L_z = -i\hbar\frac{\partial}{\partial\theta}$.

$$\frac{1}{2m} \left[-\hbar^2 \left(\frac{1}{r} \frac{\partial}{\partial r} \left[r \frac{\partial\psi}{\partial r} \right] + \frac{1}{r^2} \frac{\partial^2\psi}{\partial\theta^2} + \frac{\partial^2\psi}{\partial z^2} \right) + ie\hbar B_0 \frac{\partial\psi}{\partial\theta} \right] + \frac{e^2B_0^2}{2m} r^2\psi + eKz^2\psi = E\psi.$$

$$\frac{-\hbar^2}{2m} \left[\left(\frac{1}{r} \frac{\partial}{\partial r} \left[r \frac{\partial \psi}{\partial r} \right] - \frac{e^2 B_0^2}{\hbar^2} r^2 \psi \right) + \left(\frac{1}{r^2} \frac{\partial^2 \psi}{\partial \theta^2} - \frac{ieB_0}{\hbar} \frac{\partial \psi}{\partial \theta} \right) + \left(\frac{\partial^2 \psi}{\partial z^2} - \frac{2meKz^2}{\hbar^2} \psi \right) \right] = E\psi.$$

Assuming the problem separates, i.e. $\psi = R(r)\Theta(\theta)Z(z)$ where each factor depends only on the indicated variable, this becomes

$$\begin{aligned} & \frac{-\hbar^2}{2m} \left[\Theta(\theta)Z(z) \left(\frac{1}{r} \frac{\partial}{\partial r} \left[r \frac{\partial R}{\partial r} \right] - \frac{e^2 B_0^2}{\hbar^2} r^2 R(r) \right) + R(r)Z(z) \left(\frac{1}{r^2} \frac{\partial^2 \Theta}{\partial \theta^2} - \frac{ieB_0}{\hbar} \frac{\partial \Theta}{\partial \theta} \right) \right. \\ & \quad \left. + R(r)\Theta(\theta) \left(\frac{\partial^2 Z}{\partial z^2} - \frac{2meKz^2}{\hbar^2} Z(z) \right) \right] = ER(r)\Theta(\theta)Z(z) \\ \Leftrightarrow & \frac{-\hbar^2}{2m} \left[\frac{1}{R(r)} \left(\frac{1}{r} \frac{\partial}{\partial r} \left[r \frac{\partial R}{\partial r} \right] - \frac{e^2 B_0^2}{\hbar^2} r^2 R(r) \right) + \frac{1}{\Theta(\theta)} \left(\frac{1}{r^2} \frac{\partial^2 \Theta}{\partial \theta^2} - \frac{ieB_0}{\hbar} \frac{\partial \Theta}{\partial \theta} \right) \right. \\ & \quad \left. + \frac{1}{Z(z)} \left(\frac{\partial^2 Z}{\partial z^2} - \frac{2meKz^2}{\hbar^2} Z(z) \right) \right] = E. \end{aligned}$$

Each term must be equal to a constant, since the sum is: if any varied with respect to its variable, the others don't depend on that variable, and so can't vary to compensate and make the entire expression equal E . The problem then splits into three ODEs:

$$\begin{aligned} \frac{1}{r} \frac{d}{dr} \left(r \frac{dR}{dr} \right) &= \left(c_1 + \frac{e^2 B_0^2}{\hbar^2} r^2 \right) R(r), \\ \frac{1}{r^2} \frac{d^2 \Theta}{d\theta^2} - \frac{ieB_0}{\hbar} \frac{d\Theta}{d\theta} - c_2 \Theta(\theta) &= 0, \\ \frac{d^2 Z}{dz^2} &= \left(c_3 + \frac{2meK}{\hbar^2} z^2 \right) Z(z) \end{aligned}$$

□

Problem 5. Consider the observables $A = x^2$ and $B = L_z$.

1. Find the uncertainty principle governing A and B , that is, $\Delta A \Delta B$.
2. Evaluate ΔB for the hydrogenic $|n\ell m\rangle$ state.
3. Therefore, what can you conclude about $\langle xy \rangle$ in this state?