

4123 HW 4

Duncan Wilkie

9 November 2021

1a

The Lagrangian for this system is

$$L = \frac{1}{2}m(\dot{x}^2 + \dot{y}^2) + mgy$$

The modified Euler-Lagrange equations are

$$\frac{\partial L}{\partial x} + \lambda \frac{\partial C}{\partial x} = \frac{d}{dt} \frac{\partial L}{\partial \dot{x}} \Leftrightarrow 2\lambda x = m\ddot{x}$$

and

$$\frac{\partial L}{\partial y} + \lambda \frac{\partial C}{\partial y} = \frac{d}{dt} \frac{\partial L}{\partial \dot{y}} \Leftrightarrow mg + 2\lambda y = m\ddot{y}$$

Adding these together and solving for λ , we obtain

$$mg + 2\lambda(x + y) = m(\ddot{x} + \ddot{y}) \Leftrightarrow \lambda = \frac{m}{2} \frac{\ddot{x} + \ddot{y} - g}{x + y}$$

1b

Differentiating the constraint twice,

$$C = 0 \Rightarrow x\dot{x} + y\dot{y} = 0 \Rightarrow x\ddot{x} + \dot{x}^2 + y\ddot{y} + \dot{y}^2 = 0$$

Solving for \ddot{x} and \ddot{y} ,

$$\begin{aligned}\ddot{x} &= -\frac{y\ddot{y} + \dot{x}^2 + \dot{y}^2}{x} \\ \ddot{y} &= -\frac{x\ddot{x} + \dot{x}^2 + \dot{y}^2}{y}\end{aligned}$$

Adding,

$$\begin{aligned}\ddot{x} + \ddot{y} &= -\left(\frac{y^2\ddot{y} + y\dot{x}^2 + y\dot{y}^2 + x^2\ddot{x} + x\dot{x}^2 + x\dot{y}^2}{xy}\right) \\ &= \frac{-1}{xy} (x^2\ddot{x} + y^2\ddot{y} + (x + y)(\dot{x}^2 + \dot{y}^2))\end{aligned}$$

Plugging this in to the equation for λ ,

$$\lambda = -\frac{m}{2} \left(\frac{x^2\ddot{x} + y^2\ddot{y}}{x^2y + y^2x} + \frac{\dot{x}^2 + \dot{y}^2}{xy} + \frac{g}{x+y} \right)$$

Multiplying by $1 = \frac{l^2}{l^2}$ and noting that $l^2 = x^2 + y^2 = (x+y)^2 - 2xy$,

=

2a

Since there is no “real” potential, there is no force on m_2 and it must move with constant velocity.

2b

The resulting Lagrangian in the CM frame (which is approximately the m_2 frame for the same reason) is

$$L = \frac{1}{2}m_2\dot{r}^2 - \frac{l^2}{2m_2r^2}$$

The resulting Euler-Lagrange equation of motion is

$$\frac{\partial L}{\partial r} = \frac{d}{dt} \frac{\partial L}{\partial \dot{r}} \Leftrightarrow \frac{l^2}{m_2r^3} = m_2\ddot{r} \Leftrightarrow \frac{l^2}{m_2^2} = r^3\ddot{r}$$

There is a centrifugal force term present. This is a second-order autonomous equation with no first-derivative dependence, and so can be solved as

$$\begin{aligned} t(r) + c_1 &= \pm \int \frac{dr}{\sqrt{2 \int \frac{l^2}{m_2^2 r^3} dr + c_2}} = \pm \frac{m_2}{l} \int \frac{dr}{\sqrt{c_2 - \frac{1}{r^2}}} = \pm \frac{m_2}{l} \frac{r \sqrt{c_2 - \frac{1}{r^2}}}{c_2} \\ &\Rightarrow \sqrt{r^2 c_2 - 1} = \pm \left(c_2 t \frac{l}{m_2} + c_1 c_2 \right) \\ &\Leftrightarrow r^2 = 1 + \left(t \frac{l}{m_2} + c \right)^2 \end{aligned}$$

This is a hyperbola, translated in the t -axis by c . This is not consistent with a constant-velocity trajectory, but since hyperbolas have oblique asymptotes it becomes approximately a constant-velocity trajectory. Therefore, the deviance we see is a breakdown of the approximation of the CM frame by the m_1 frame.

3

The ordinary central-force radial Lagrangian is

$$L = \frac{1}{2}\mu\dot{r}^2 - \frac{l^2}{2\mu r^2} - V(r)$$

The corresponding Euler-Lagrange equation is

$$\frac{\partial L}{\partial r} = \frac{d}{dt} \frac{\partial L}{\partial \dot{r}} \Leftrightarrow -\frac{\partial V}{\partial r} + \frac{l^2}{\mu r^3} = \mu\ddot{r} \Leftrightarrow F(r) + \frac{l^2}{\mu r^3} = \mu\ddot{r}$$

Taking r to be a function of θ alone, $\frac{d^2 s}{d\theta^2} = \frac{\partial s}{\partial r} r'' + \frac{\partial^2 s}{\partial \theta \partial r} r' = \frac{-r''}{r^2}$ (since s is not directly dependent on θ). The target equation may be written in terms of r as

$$\begin{aligned} \frac{-r''}{r^2} + \frac{1}{r} &= \frac{-\mu r^2}{l^2} F(r) \Leftrightarrow \frac{r'' - r}{r^2} \frac{l^2}{\mu r^2} = F(r) \\ &\Leftrightarrow \frac{l^2 r''}{\mu r^4} - \frac{l^2}{\mu r^3} = F(r) \end{aligned}$$

Noting that by the chain rule $\ddot{r} = r'\ddot{\theta} + r''\dot{\theta}^2 = r''\dot{\theta}^2 = r''\left(\frac{l}{\mu r^2}\right)^2$ (constant angular momentum \Rightarrow no angular acceleration), $\frac{l^2 r''}{\mu r^4} = \mu\ddot{r}$. Therefore, the target equation is

$$\mu\ddot{r} - \frac{l^2}{\mu r^3} = F(r)$$

which is clearly equivalent to the the Euler-Lagrange equation found above.

4

We have $s = \frac{1}{k\theta^2}$, so $\frac{d^2 s}{d\theta^2} = \frac{6}{k\theta^4}$. The above equation is then

$$\begin{aligned} \frac{6}{k\theta^4} + \frac{1}{k\theta^2} &= \frac{-\mu k^2 \theta^4}{l^2} F(r) \\ \Leftrightarrow F(r) &= \frac{-l^2}{\mu k^2 \theta^4} \left(\frac{6}{k\theta^4} + \frac{1}{k\theta^2} \right) \end{aligned}$$