1. VERIFY

Confirm the order in CPRS

Service	Order	Start / Stop	Provider	Nurse	Clerk	0
Inpt. Meds	INFLUENZA INJ 0.5 ml IM ONCE For age 65 or older administer Fluzone HIGH DOSE Trivalent vaccine. For age 64 or under administer Fluarix Trivalent vaccine. View appropriateness of vaccine PRIOR to completing IMMUNIZATION TEMPLATE. Document in both BCMA and Influenza Immunization template as applicable. Indication: INFLUENZA VACCINATION		Geurkink,Eric A			

and BCMA


INFLUENZA INJ INFLUENZA VACCINE,QUAD (PF) SYR 0.5ML	1 SYRINGE,	INTRAM	DUE
For age 65 or older administer Fluzone HIGH DOSE Quadrivalent 0.7 mL	ONCE		
syringe. For age 64 or under administer Fluzone Quadrivalent 0.5 mL syringe. Use immunization template to document			
administration.			

Vaccine orders are activated from the Admission orders after being verified by pharmacy (**The BCMA example listed in this education does not list the current year's influenza vaccine, however, the process is still the same)

Locate the Influenza Immunization information:

> Immunization History

Reports tab > Health Summary > Immunizations Minneapolis



2. SCREEN

Per available stock and ACIP 2024-25 recommendations, the VA will provide two options:

Sanofi Fluzone High-dose: adults aged ≥65 years

(A high dose includes 3-4 times as much flu virus antigen for an increased immune response)



 Fluarix – standard influenza vaccine recommended for adults ≤64 years





Vaccine FYI:

- The standard trivalent dose is available to persons ≥65 years by request only
- Screen each patient per department process prior to giving vaccine
- Eggs and latex allergies are NOT considered contraindications to receiving the vaccine, the only contraindication to the flu vaccination is severe allergic reaction (ie. Anaphylactic reactions)
- Review link for community options to receive flu vaccines:
 Flu Shots Near You National Center for Health Promotion and Disease Prevention (va.gov)

3. SUPPLIES



Supplies needed:

- Prefilled syringe in fridge
- Safety locking needle (22g-25g, 1-1½ inch)
- Alcohol prep
- Gloves
- Face Shield (optional)
- Vaccine Information Sheet (VIS)
- Bandaid/cotton balls






- The High-Dose is a 0.5 ml dose and comes in packages of two
- The Standard dose is a 0.5 ml dose and comes in a package with a row of ten syringes

Both vaccines should be:

- Stored in a refrigerator between 2º and 8ºC (36º and 46ºF)
- Do not freeze, discard if the vaccine has been frozen

4 ADMINISTRATION

- Administer flu vaccine in deltoid muscle
- This muscle can be used for more than one IM injection including Covid-19 vaccine
- Separate injection sites by 1 inch or more, if possible



- Administer vaccine to the patient's deltoid (locate the acromion process, which is the prominent part of the shoulder bone; then aim for administration at approximately 2-3 finger widths below)
- Administer at a 90- degree angle
- Do not aspirate

5. MONITOR

• Monitor for any suspected allergic reactions:


Anaphylactic	Symptoms of mild <u>local</u> reactions:	Actions for mild local reactions:
Reaction to	Redness	 Ice pack to localized area
Vaccines and	Itching	
medications	Tenderness	
	Localized edema at site	
	Symptoms of systemic reactions including:	Actions for any systemic reaction:
	Hives	Call RRT: 612-919-5795
	Dizziness	Code Blue/SMART: 31-1911
	Shortness of breath Tongue/throat/angioedema	Maintain airway
	Nausea	Document adverse reaction in CPRS
	Weakness	Complete Joint Patient Safety
	Hypotension	Reporting (JPSR)
	Loss of Consciousness	

 Health care providers are required to report to VAERS any adverse event listed by the vaccine manufacturer as a contraindication to further doses of vaccine and adverse events listed: https://vaers.hhs.gov/

6. DOCUMENT


CPRS: Influenza Vaccine Template

- The VIS hyperlink takes you to the most up-to-date version from the CDC
- If the patient is
 - o ≥65 years old both doses will be listed
- After selecting a dose, fill out the VIMM




7. The VIMM

- Verify the correct vaccination, lot number, and expriation date
 - If there are any discrepancies contact pharmacy
- <u>All</u> asterisked fields are required



8. <u>BCMA</u>

- Scan vaccine in BCMA.
 - Make sure to scan in BCMA before signing your note in CPRS
- Only mark "held" if the patient has already received the vaccine.
- If the patient declines/refuses, then mark in BCMA as refused
 - Reminder: refusals will still need to be documented in CPRS



(**The BCMA example listed in this education does not list the current year's influenza vaccine, however, the process is still the same)

If there are any further questions, please reach out to your manager