



Tarlatamab

- Must stay/live within 1 hour of a healthcare setting for a total of 48 hours (2 days) from the start of the infusion following the first and second infusions.
- 24 hour caregiver requirement: A dedicated caregiver should be present for at least 48 hours (2 days) following the first and second infusions, to observe for vital sign and neurologic changes.

Home Monitoring

- You will receive home monitoring equipment prior to your start date & this will be done through our home telehealth program *Note this will just be to document the information. You and your caregiver will be responsible for self-assessment and seeking **immediate** care with changes
- This will include blood pressure machine, thermometer, pulse oximeter.

Vital signs: blood pressure, temperature, pulse, pulse oximetry to be completed and uploaded to home telehealth 4 times DAILY. Neurologic function to be assessed by patient or caregiver TWICE DAILY

Seek **IMMEDIATE** care if:

CRS (Cytokine release syndrome) symptoms arise including any of the following:

- | | | |
|--|--|------------------------------------|
| -Temperature of 100.4 (38 C) or higher | -Low blood pressure: SPB <90 | -Fast heart rate: greater than 130 |
| -Shortness of breath/trouble breathing | -Pulse oximetry reading of 90% or less | |
| -Issues with balance/movement | -Dizziness | -Headache |
| -Confusion/restlessness/anxious | -Nausea/Vomiting | -Prolonged bleeding |

Any neurological changes, including:

- Confusion, word-finding difficulty, tremors, somnolence, and/or disorientation.

Handwriting changes when writing the sentence "Our national bird is the bald eagle"

- Write sentence on worksheet twice daily.

For IMMEDIATE care please call 911 or go to your nearest emergency department. Notify providers you are receiving Tarlatamab.