

Root systems

Antoine de Saint Germain and Ambrose Tang

May 14, 2026

0.1 Root pairings

0.1.1 Definitions

Recall that a *perfect pairing* is a quadruple $(R, M, N, \mathcal{L})$ where R is a ring, M and N are R -modules, and $\mathcal{L} : M \times N \rightarrow R$ is a bilinear map such that the induced maps $M \rightarrow \text{Hom}_R(N, R)$ and $N \rightarrow \text{Hom}_R(M, R)$ are isomorphisms of R -modules.

If I is a set, denote by $\text{Perm}(I)$ the group of permutations of I .

Definition 1. A *root pairing* is a tuple $(R, M, N, \mathcal{L}, I, \alpha, \alpha^\vee, s)$ where $(R, M, N, \mathcal{L})$ is a perfect pairing, I is a set, $\alpha : I \rightarrow M, i \mapsto \alpha_i$, $\alpha^\vee : I \rightarrow N, i \mapsto \alpha_i^\vee$, and $s : I \rightarrow \text{Perm}(I), i \mapsto s_i$, with the conditions that

1. $\alpha, \alpha^\vee$ are injective,
2. For all $i \in I$, $\mathcal{L}(\alpha_i, \alpha_i^\vee) = 2$,
3. For all $i, j \in I$, we have

$$\begin{aligned}\alpha_{s_i(j)} &= \alpha_j - \mathcal{L}(\alpha_j, \alpha_i^\vee) \alpha_i, \\ \alpha_{s_i(j)}^\vee &= \alpha_j^\vee - \mathcal{L}(\alpha_i, \alpha_j^\vee) \alpha_i^\vee.\end{aligned}$$

0.1.2 Root pairing of type A

Consider the perfect pairing $(R, M, N, \mathcal{L}) = (\mathbb{Z}, \mathbb{Z}^n, (\mathbb{Z}^n)^*, (\cdot, \cdot))$ where $(\cdot, \cdot)$ is the canonical pairing between $\mathbb{Z}^n$ and its algebraic dual $(\mathbb{Z}^n)^* := \text{Hom}_{\mathbb{Z}}(\mathbb{Z}^n, \mathbb{Z})$, i.e. the dot product.

We denote by $e_1, \dots, e_n$ the standard basis of $\mathbb{Z}^n$ and by $e_1^*, \dots, e_n^*$ the dual basis of $(\mathbb{Z}^n)^*$.

Write $[1, n] = \{1, \dots, n\}$. An *interval* in $[1, n]$ is a subset of the form $[i, j] \stackrel{\text{denotes}}{=} \{i, i+1, \dots, j\}$ for some $1 \leq i \leq j \leq n$. We denote by I_n the set of intervals in $[1, n]$. Write $[i] := [i, i]$ for $1 \leq i \leq n$.

A *signed interval* in $[1, n]$ is an element of $I_n := I_n \times \{1, -1\}$. We denote by I_n^+ the set of signed intervals of the form $(J, 1)$ and by I_n^- the set of signed intervals of the form $(J, -1)$. It is clear that $I_n = I_n^+ \sqcup I_n^-$. Throughout the rest of this file, we write J for the signed interval $(J, 1)$ and $-J$ for $(J, -1)$.

Let $A = (a_{i,j})$ be the Cartan matrix of type A_n defined by $a_{i,i} = 2$ for all $i \in [1, n]$, $a_{i,j} = -1$ if $|i - j| = 1$, and $a_{i,j} = 0$ otherwise.

Define

$$\begin{aligned}\alpha : I_n &\rightarrow \mathbb{Z}^n, & \forall J \in I_n^+, \quad \alpha_{\pm J} &= \pm \left(\sum_{j \in J} A e_j \right), \\ \alpha^\vee : I_n &\rightarrow (\mathbb{Z}^n)^*, & \forall J \in I_n^+, \quad \alpha_{\pm J}^\vee &= \pm \left(\sum_{j \in J} e_j^* \right).\end{aligned}$$

Finally, define $s : I_n^+ \rightarrow \text{Perm}(I_n)$ as follows. Fix $J, K \in I_n^+$. It is clear that either $J = K$, $J \subset K$, $K \subset J$, or and set $L = (J \cup K) \setminus (J \cap K)$. It is clear that J, K, L satisfy exactly one of the following four conditions.

R1. $L = \emptyset$.

R2. $L \in I_n$ and $K \subset J$, and

R3. $L \in I_n$ and $K \not\subset J$.

R4. $L \notin I_n \cup \{\emptyset\}$.

Then,

$$s_J(K) = \begin{cases} -K & \text{in Case \textbf{R1},} \\ -L & \text{in Case \textbf{R2},} \\ L & \text{in Case \textbf{R3}.} \\ K & \text{in Case \textbf{R4},} \end{cases} \quad (1)$$

If $J = [i, j]$ and $K = [k, l]$, then (1) can be rewritten as

$$s_{[i,j]}[k, l] = \begin{cases} -[k, l] & \text{if } (i, j) = (k, l) \\ -[l + 1, j] & \text{if } i = k, j > l \\ [j + 1, l] & \text{if } i = k, j < l \\ -[i, k - 1] & \text{if } i < k, j = l \\ [k, i - 1] & \text{if } i > k, j = l \\ [k, j] & \text{if } i = l + 1 \\ [i, l] & \text{if } k = j + 1 \\ [k, l] & \text{otherwise.} \end{cases}$$

We extend s_J to I_n by setting $s_J(-K) = -s_J(K)$ for all $K \in I_n^+$. We extend s to I_n by setting $s_{-J} = s_J$ for all $J \in I_n^+$.

Lemma 2. *For all J , we have $s_J \in \text{Perm}(I_n)$.*

Proof. Fix $J \in I_n$. We prove that $s_J^2 = \text{id}$ by a straightforward case-by-case check using the definition of s_J and the four cases above. Then $s_J^{-1} = s_J$, and our claim is proved. $\square$

Theorem 3. *The 8-tuple $(\mathbb{Z}, \mathbb{Z}^l, (\mathbb{Z}^l)^*, (\cdot, \cdot), I_n, \alpha, \alpha^\vee, s)$ is a root pairing.*

Proof. $\square$