

The short-lived ^{53}Mn - ^{53}Cr chronology and its application in the origin of angrite parent body.

Ke Zhu (朱柯)^{1*}, Frédéric Moynier¹, Jean-Alix Barrat², Martin Bizzarro³

¹Institut de Physique du Globe de Paris, Université Sorbonne Paris Cité, CNRS, 1 rue Jussieu, Paris 75005, France. ²Laboratoire Géosciences Océan (UMR CNRS 6538), Université de Bretagne Occidentale et Institut Universitaire Européen de la Mer, Place Nicolas Copernic, 29280 Plouzané, France. ³Centre for Star and Planet Formation and Natural History Museum of Denmark, University of Copenhagen, Øster Voldgade 5–7, Copenhagen DK-1350, Denmark. (*corresponding author: zhu@ipgp.fr)

Abstract: Chromium (Cr) isotopes is probably one of the best suited system to study the timing and genetic relationships. The short-lived nuclide ^{53}Mn , with a half-life of 3.7 ± 0.2 Myrs, decays to ^{53}Cr (radiogenic origin) and therefore the ^{53}Mn - ^{53}Cr decay system is a useful chronometer to date early Solar System events. On the other hand, the variation of $^{54}\text{Cr}/^{52}\text{Cr}$ ratios (nucleosynthetic origin) can be a good tracer to track the kinship of Solar System materials.

Angrite meteorites are some of the oldest materials in the solar system. They provide important information on the earliest evolution of the solar system and accretion timescales of protoplanets. Here, we show that the $^{54}\text{Cr}/^{52}\text{Cr}$ ratio is homogeneously distributed among angrite meteorites within 13 parts per million, indicating that precursor materials must have experienced a global-scale melting such as a magma ocean. The $^{53}\text{Cr}/^{52}\text{Cr}$ and Mn/Cr ratios are correlated which is evidence for an initial $^{53}\text{Mn}/^{55}\text{Mn}$ ratio of $(3.17 \pm 0.21) \times 10^{-6}$. When anchored to the U-corrected Pb-Pb age for the D'Orbigny angrite, this initial $^{53}\text{Mn}/^{55}\text{Mn}$ corresponds to an absolute age of 4563.3 ± 0.4 Ma, i.e. 4.0 ± 0.4 Ma after CAI-formation. This age is distinct from the one of the volatile depletion events dated by the $^{87}\text{Sr}/^{86}\text{Sr}$ initial ratio and therefore must correspond to the age of crystallization of the magma ocean and crust formation of angrite parent body (APB), which can also constrain a bigger size of APB than that of Vesta. Furthermore, this age is similar to those obtained from internal isochrons of the oldest volcanic angrites that cooled rapidly at the surface of the parent body (with ages of $4564 \sim 4563$ Ma), while older than those obtained from plutonic angrites ($4561 \sim 4556$ Ma) that cooled down slowly, located deeper within the parent body. This implies that cooling of the angrite parent body took at least ~ 8 Myrs after its formation.

Key words: Geo and cosmochemistry, Sun and Solar system.