

Uncovering the sulfur isotopic composition of fluids from HP-metamorphic slab

Genot I., Cartigny P., Angiboust S.

In order to better understand the fate and behavior of sulfur in subduction zones and the sulfur isotopic signature in arcs, we analyzed the $^{33}\text{S}/^{32}\text{S}$, $^{34}\text{S}/^{32}\text{S}$ and $^{36}\text{S}/^{32}\text{S}$ ratios (expressed as $\delta^{34}\text{S} = (^{34}\text{R}_{\text{sample}}/^{34}\text{R}_{\text{std}} - 1) * 1000$; $\Delta^{33}\text{S} = \delta^{33}\text{S} - ((1 + \delta^{34}\text{S}/1000)^{0.515} - 1) * 1000$ and $\Delta^{36}\text{S} = \delta^{36}\text{S} - ((1 + \delta^{34}\text{S}/1000)^{1.89} - 1) * 1000$) in eclogite breccias, serpentinites, metasediments from the Lower Shear Zone (LSZ) of Monviso (Alps, Italy) recording evidence of a channelized fluid flow at the metamorphic peak $\sim 550^\circ\text{C}$ and 2.6 GPa, as well as metabasalt and metagabbros unaffected by these fluids.

The $\delta^{34}\text{S}$ ranges from -12.4 to $20.8 \pm 0.2\text{‰}$, $\Delta^{33}\text{S}$ from 0 to 0.07 ± 0.01 and $\Delta^{36}\text{S}$ from -0.3 to $0.4 \pm 0.2\text{‰}$. Metabasalt and metagabbros non-affected by fluids have a $\delta^{34}\text{S}$ between 0 and 2‰ that suggests a magmatic origin and/or hydrothermal alteration on the seafloor whereas other samples in the LSZ are ^{33}S - ^{34}S -enriched compared to the former, except a metasediment being ^{34}S -depleted. Furthermore, the absence of sulfate and the magnetite precipitation in the LSZ suggest that the fluid was reduced, with sulfur as HS^- or H_2S . We can interpret our data using at least two sources of S that were mixed in the fluid. A first one is ^{33}S - ^{34}S -enriched and could derive from sulfates from either evaporite of the Dora Maira unit or from antigorite breakdown ($\delta^{34}\text{S} \sim 21\text{‰}$). The second source is ^{34}S -depleted and ^{33}S -enriched and could be typified by sulfate reduced by microorganisms on the seafloor in sediments and altered oceanic crust.

This study clearly illustrates that arc lavas could have variable $\delta^{34}\text{S}$ from negative values (biogenic sulfides in altered oceanic crust and sediments) to positive values (evaporite and/or antigorite breakdown).

Keywords: Geochemistry, stable isotope, subduction zone