

# Why Symplectic Singularities?

Antoine Bourget  
IPhT, CEA, Saclay

Fields and String Seminar, Munich

November 26, 2023

Based on long time collaboration with P. Argyres, S. Cabrera, A. Collinucci, S. Giacomelli, J. Grimminger, A. Hanany, R. Kalveks, M. Martone, S. Schäfer-Nameki, M. Sperling, G. Zafrir, Z. Zhong.  
[AB, Grimminger, 2209.15016]

# Why Symplectic Singularities ?

- Symplectic singularities are a basic kind of geometric spaces.
- They feature in moduli space of vacua of supersymmetric theories.
- They provide a geometric perspective to understand
  - the phases of a given theory
  - the landscape of all theories
  - and the relations between them.

# Why Symplectic Singularities ?

- 1 Introduction and Examples
- 2 Definition of Conical Symplectic Singularities
- 3 Moduli Space of Vacua
  - SCFTs and Moduli Space of Vacua
  - Classification of SCFTs
  - Phases of SCFTs
- 4 CSS stratification and fibrations
  - General Strategy
  - Phase diagrams from Magnetic Quivers
  - The quest for the elementary building blocks
- 5 Conclusion

## 1 Introduction and Examples

## 2 Definition of Conical Symplectic Singularities

## 3 Moduli Space of Vacua

- SCFTs and Moduli Space of Vacua
- Classification of SCFTs
- Phases of SCFTs

## 4 CSS stratification and fibrations

- General Strategy
- Phase diagrams from Magnetic Quivers
- The quest for the elementary building blocks

## 5 Conclusion

# Example 1 : Gravitational Instantons

Start from Asymptotically Locally Euclidean HyperKähler 4-manifolds.  
For instance Eguchi-Hanson space:  $T^*\mathbb{P}^1$  with metric

$$ds^2 = \frac{dr^2}{1 - \frac{a}{r^4}} + \frac{r^2}{4} \left(1 - \frac{a}{r^4}\right) \sigma_3^2 + \frac{r^2}{4} (\sigma_1^2 + \sigma_2^2)$$

with

$$\sigma_1 = \sin \psi d\theta - \cos \psi \sin \theta d\phi$$

$$\sigma_2 = \cos \psi d\theta + \sin \psi \sin \theta d\phi$$

$$\sigma_3 = d\psi + \cos \theta d\phi$$

# Example 1 : Gravitational Instantons

Start from Asymptotically Locally Euclidean HyperKähler 4-manifolds.  
For instance Eguchi-Hanson space:  $T^*\mathbb{P}^1$  with metric

$$ds^2 = \frac{dr^2}{1 - \frac{a}{r^4}} + \frac{r^2}{4} \left(1 - \frac{a}{r^4}\right) \sigma_3^2 + \frac{r^2}{4} (\sigma_1^2 + \sigma_2^2)$$

with

$$\sigma_1 = \sin \psi d\theta - \cos \psi \sin \theta d\phi$$

$$\sigma_2 = \cos \psi d\theta + \sin \psi \sin \theta d\phi$$

$$\sigma_3 = d\psi + \cos \theta d\phi$$

When  $a \rightarrow 0$ :

$$T^*\mathbb{P}^1 \rightarrow \mathbb{R}^4 / \mathbb{Z}_2.$$

This is the  $A_1$  singularity.

## Example 2 : Moduli Space of Gauge Instantons

Let  $\mathcal{M}_{k,N}$  be the moduli space of  $k$   $SU(N)$  framed instantons on  $\mathbb{R}^4$ . It is a singular HyperKähler manifold,

$$\dim_{\mathbb{R}} \mathcal{M}_{k,N} = 4kN .$$

## Example 2 : Moduli Space of Gauge Instantons


Let  $\mathcal{M}_{k,N}$  be the moduli space of  $k$   $SU(N)$  framed instantons on  $\mathbb{R}^4$ . It is a singular HyperKähler manifold,

$$\dim_{\mathbb{R}} \mathcal{M}_{k,N} = 4kN.$$

For  $k = 1, N = 2$  :

$$\mathcal{M}_{1,2} = \mathbb{R}^4 \times \mathbb{R}^4 / \mathbb{Z}_2 = \mathbb{H} \times A_1.$$

Singularity : small instanton transition.





## Example 3 : Higgs branches

Supersymmetric QFT in spacetime dimension  $3 \leq d \leq 6$  with 8 supercharges have *Higgs branches*.

## Example 3 : Higgs branches

Supersymmetric QFT in spacetime dimension  $3 \leq d \leq 6$  with 8 supercharges have *Higgs branches*.

Exemple: SQED with gauge group  $U(1)$  and  $N_f = 2$ :

$$\mathbb{C}[\mathcal{H}] = \left( \frac{\mathbb{C}[Q_{2 \times 1}, \tilde{Q}_{1 \times 2}]}{(\tilde{Q}Q = 0, |Q|^2 - |\tilde{Q}|^2 = 0)} \right)^{U(1)} \quad \mathcal{H} = \mathbb{C}^4 // U(1) = \mathbb{C}^2 / \mathbb{Z}_2.$$


## Example 3 : Higgs branches

Supersymmetric QFT in spacetime dimension  $3 \leq d \leq 6$  with 8 supercharges have *Higgs branches*.

Exemple: SQED with gauge group  $U(1)$  and  $N_f = 2$ :

$$\mathbb{C}[\mathcal{H}] = \left( \frac{\mathbb{C}[Q_{2 \times 1}, \tilde{Q}_{1 \times 2}]}{(\tilde{Q}Q = 0, |Q|^2 - |\tilde{Q}|^2 = 0)} \right)^{U(1)} \quad \mathcal{H} = \mathbb{C}^4 // U(1) = \mathbb{C}^2 / \mathbb{Z}_2.$$

$$\mathcal{H} = \{M_{2 \times 2} | M^2 = 0, \text{tr}(M) = 0\}, \text{ with } M = Q\tilde{Q}.$$





Resolution / Deformation by Fayet-Iliopoulos terms.

# Example 3 : Higgs branches


Brane system for 3d  $\mathcal{N} = 4$  system (Hanany-Witten):

	$x^0$	$x^1$	$x^2$	$x^3$	$x^4$	$x^5$	$x^6$	$x^7$	$x^8$	$x^9$
NS5	x	x	x	x	x	x				
D3	x	x	x				x			
D5	x	x	x					x	x	x



# Example 3 : Higgs branches

Full moduli space of vacua for 3d  $\mathcal{N} = 4$  SQED with  $N_f = 2$ :



## Example 4 : Nilpotent Orbits

Nilpotent cone:

$$\mathcal{N}(\mathfrak{sl}_N) = \{M \in \text{Mat}(N, \mathbb{C}) \mid M^N = 0\}$$

## Example 4 : Nilpotent Orbits

Nilpotent cone:

$$\mathcal{N}(\mathfrak{sl}_N) = \{M \in \text{Mat}(N, \mathbb{C}) \mid M^N = 0\}$$

Exemple :  $N = 2$

$$\mathcal{N}(\mathfrak{sl}_2) = \left\{ M = \begin{bmatrix} a & b \\ c & -a \end{bmatrix} \in \text{Mat}(N, \mathbb{C}) \mid a^2 + bc = 0 \right\}$$

# Example 4 : Nilpotent Orbits


Nilpotent cone:

$$\mathcal{N}(\mathfrak{sl}_N) = \{M \in \text{Mat}(N, \mathbb{C}) \mid M^N = 0\}$$

Exemple :  $N = 2$

$$\mathcal{N}(\mathfrak{sl}_2) = \left\{ M = \begin{bmatrix} a & b \\ c & -a \end{bmatrix} \in \text{Mat}(N, \mathbb{C}) \mid a^2 + bc = 0 \right\}$$


Springer resolution:  $T^*\text{Flag}_N \rightarrow \mathcal{N}(\mathfrak{sl}_N)$






# Example 4 : Nilpotent Orbits

Nilpotent orbits and stratification of the nilpotent cone:



# Example 4 : Nilpotent Orbits

Nilpotent orbits and stratification of the nilpotent cone:



Minimal nilpotent orbit closure  $\overline{\mathcal{O}_{\min}(\mathfrak{g})}$ :

$$\dim_{\mathbb{R}} \overline{\mathcal{O}_{\min}(\mathfrak{g})} = 4(h^{\vee}(\mathfrak{g}) - 1)$$

$$\mathcal{M}_{1,G} = \mathbb{R}^4 \times \overline{\mathcal{O}_{\min}(\mathfrak{g})}$$

- 1 Introduction and Examples
- 2 Definition of Conical Symplectic Singularities
- 3 Moduli Space of Vacua
  - SCFTs and Moduli Space of Vacua
  - Classification of SCFTs
  - Phases of SCFTs
- 4 CSS stratification and fibrations
  - General Strategy
  - Phase diagrams from Magnetic Quivers
  - The quest for the elementary building blocks
- 5 Conclusion

# Common features / Definition

- Presence of singularities ;
- Complex symplectic structure on smooth locus ;
- Singular limit of a HyperKähler manifolds ;
- (Complexified) scaling:  $\mathbb{C}^*$ -action compatible with symplectic structure.

# Common features / Definition

- Presence of singularities ;
- Complex symplectic structure on smooth locus ;
- Singular limit of a HyperKähler manifolds ;
- (Complexified) scaling:  $\mathbb{C}^*$ -action compatible with symplectic structure.

A Poisson variety  $(X, \{, \})$  is a *Conical Symplectic Singularity* if:

- 1 The Poisson structure is non-degenerate on  $X^{\text{reg}}$  : there is a symplectic form  $\omega$  ;
- 2 For some / any resolution of singularities  $\pi : Y \rightarrow X$ ,  $\pi^*\omega$  extends to a regular 2-form on  $Y$ ;
- 3 There is a  $\mathbb{C}^*$  action that contracts  $X$  to a point under which  $\omega$  has positive integer weight.

# Common features / Definition

- Presence of singularities ;
- Complex symplectic structure on smooth locus ;
- Singular limit of a HyperKähler manifolds ;
- (Complexified) scaling:  $\mathbb{C}^*$ -action compatible with symplectic structure.

A Poisson variety  $(X, \{, \})$  is a *Conical Symplectic Singularity* if:


- 1 The Poisson structure is non-degenerate on  $X^{\text{reg}}$  : there is a symplectic form  $\omega$  ;
- 2 For some / any resolution of singularities  $\pi : Y \rightarrow X$ ,  $\pi^*\omega$  extends to a regular 2-form on  $Y$ ;
- 3 There is a  $\mathbb{C}^*$  action that contracts  $X$  to a point under which  $\omega$  has positive integer weight.

A Conical Symplectic Singularity is *isolated* (ICSS) if the singular locus is a point.

# Stratification and Transverse Slices

ICSS are the building blocks of all CSS:

- A CSS has a stratification into symplectic leaves  $\rightarrow$  Hasse diagram
- The *transverse slices* are ICSS.




<sup>1</sup>Normal and with trivial local fundamental group.

# Stratification and Transverse Slices

ICSS are the building blocks of all CSS:

- A CSS has a stratification into symplectic leaves  $\rightarrow$  Hasse diagram
- The *transverse slices* are ICSS.



Open question: is there a classification of ICSS?<sup>1</sup>


<sup>1</sup>Normal and with trivial local fundamental group.



# Stratification and Transverse Slices

ICSS are the building blocks of all CSS:

- A CSS has a stratification into symplectic leaves  $\rightarrow$  Hasse diagram
- The *transverse slices* are ICSS.



Open question: is there a classification of ICSS?<sup>1</sup>

Only known example is  $\overline{\mathcal{O}_{\min}(\mathfrak{g})}$  [Beauville, 2000]

But see later...

<sup>1</sup>Normal and with trivial local fundamental group.

- 1 Introduction and Examples
- 2 Definition of Conical Symplectic Singularities
- 3 **Moduli Space of Vacua**
  - SCFTs and Moduli Space of Vacua
  - Classification of SCFTs
  - Phases of SCFTs
- 4 CSS stratification and fibrations
  - General Strategy
  - Phase diagrams from Magnetic Quivers
  - The quest for the elementary building blocks
- 5 Conclusion

# SCFTs and Moduli Space of Vacua

Dimension	Susy	Bosonic subalgebra	SCA
$d = 6$	$\mathcal{N} = (1, 0)$	$\mathfrak{so}(6, 2) \oplus \mathfrak{su}(2)_H$	$\subset \mathfrak{osp}(6, 2 1)$
$d = 5$	$\mathcal{N} = 1$	$\mathfrak{so}(5, 2) \oplus \mathfrak{su}(2)_H$	$\subset \mathfrak{f}(4)$
$d = 4$	$\mathcal{N} = 2$	$\mathfrak{so}(4, 2) \oplus \mathfrak{su}(2)_H \oplus \mathfrak{u}(1)_C$	$\subset \mathfrak{su}(2, 2 2)$
$d = 3$	$\mathcal{N} = 4$	$\mathfrak{so}(3, 2) \oplus \mathfrak{su}(2)_H \oplus \mathfrak{su}(2)_C$	$\subset \mathfrak{osp}(4 4)$

# SCFTs and Moduli Space of Vacua

Dimension	Susy	Bosonic subalgebra	SCA
$d = 6$	$\mathcal{N} = (1, 0)$	$\mathfrak{so}(6, 2) \oplus \mathfrak{su}(2)_H$	$\subset \mathfrak{osp}(6, 2 1)$
$d = 5$	$\mathcal{N} = 1$	$\mathfrak{so}(5, 2) \oplus \mathfrak{su}(2)_H$	$\subset \mathfrak{f}(4)$
$d = 4$	$\mathcal{N} = 2$	$\mathfrak{so}(4, 2) \oplus \mathfrak{su}(2)_H \oplus \mathfrak{u}(1)_C$	$\subset \mathfrak{su}(2, 2 2)$
$d = 3$	$\mathcal{N} = 4$	$\mathfrak{so}(3, 2) \oplus \mathfrak{su}(2)_H \oplus \mathfrak{su}(2)_C$	$\subset \mathfrak{osp}(4 4)$

SCFTs are

- "Rare" in 6d / 5d – isolated, rely on exceptional isomorphisms, non Lagrangian.
- More common in 4d (some Lagrangian ; existence of conformal manifolds). Classification?
- Very large number in 3d.

# SCFTs and Moduli Space of Vacua

Dimension	Susy	Bosonic subalgebra	SCA
$d = 6$	$\mathcal{N} = (1, 0)$	$\mathfrak{so}(6, 2) \oplus \mathfrak{su}(2)_H$	$\subset \mathfrak{osp}(6, 2 1)$
$d = 5$	$\mathcal{N} = 1$	$\mathfrak{so}(5, 2) \oplus \mathfrak{su}(2)_H$	$\subset \mathfrak{f}(4)$
$d = 4$	$\mathcal{N} = 2$	$\mathfrak{so}(4, 2) \oplus \mathfrak{su}(2)_H \oplus \mathfrak{u}(1)_C$	$\subset \mathfrak{su}(2, 2 2)$
$d = 3$	$\mathcal{N} = 4$	$\mathfrak{so}(3, 2) \oplus \mathfrak{su}(2)_H \oplus \mathfrak{su}(2)_C$	$\subset \mathfrak{osp}(4 4)$


SCFTs are

- "Rare" in 6d / 5d – isolated, rely on exceptional isomorphisms, non Lagrangian.
- More common in 4d (some Lagrangian ; existence of conformal manifolds). Classification?
- Very large number in 3d.

Existence of **Moduli space of vacua**, always contains **Higgs branch**.


# SCFTs and Moduli Space of Vacua

Example:  $SU(2)$  gauge theory with  $N_f = 4$ :



# SCFTs and Moduli Space of Vacua

Example:  $SU(2)$  gauge theory with  $N_f = 4$ :



## Lesson 1

The Moduli Space of Vacua can be used to "solve" certain sectors of the theory, and go beyond perturbation theory.

# Classification of SCFTs

Idea: classify possible MSV (bottom-up approach).

Example: "simplest" 4d  $\mathcal{N} = 2$  SCFTs moduli spaces

- Coulomb branch : dimension 1.

$$II^*, \quad III^*, \quad IV^*, \quad I_0^*, \quad IV, \quad III, \quad II.$$



# Classification of SCFTs

Idea: classify possible MSV (bottom-up approach).

Example: "simplest" 4d  $\mathcal{N} = 2$  SCFTs moduli spaces

- Coulomb branch : dimension 1.

$$II^*, \quad III^*, \quad IV^*, \quad I_0^*, \quad IV, \quad III, \quad II.$$

- Higgs branch : Isolated Conical Symplectic Singularities that appear as associated varieties to VOAs:

$$\mathfrak{e}_8, \quad \mathfrak{e}_7, \quad \mathfrak{e}_6, \quad \mathfrak{f}_4, \quad \mathfrak{d}_4, \quad \mathfrak{g}_2, \quad \mathfrak{a}_2, \quad \mathfrak{a}_1.$$

# Classification of SCFTs

Idea: classify possible MSV (bottom-up approach).

Example: "simplest" 4d  $\mathcal{N} = 2$  SCFTs moduli spaces

- Coulomb branch : dimension 1.

$$II^*, \quad III^*, \quad IV^*, \quad I_0^*, \quad IV, \quad III, \quad II.$$

One needs to take into account deformations! Example:  $SU(2)$   $N_f = 4$  vs  $\mathcal{N} = 4$ .

- Higgs branch : Isolated Conical Symplectic Singularities that appear as associated varieties to VOAs:

$$\mathfrak{e}_8, \quad \mathfrak{e}_7, \quad \mathfrak{e}_6, \quad \mathfrak{f}_4, \quad \mathfrak{d}_4, \quad \mathfrak{g}_2, \quad \mathfrak{a}_2, \quad \mathfrak{a}_1.$$

To match with CB classification, needs to also consider non-isolated singularities!

# Classification of SCFTs

Classification of 4d rank 1  $\mathcal{N} = 2$  SCFT Coulomb branch geometries:

Flavor	CB geometry and deformation	$\Delta(u)$
$E_8$	$II^* \rightarrow \{I_1^{10}\}$	6
$E_7$	$III^* \rightarrow \{I_1^9\}$	4
$E_6$	$IV^* \rightarrow \{I_1^8\}$	3
$D_4$	$I_0^* \rightarrow \{I_1^6\}$	2
$A_2$	$IV \rightarrow \{I_1^4\}$	$3/2$
$A_1$	$III \rightarrow \{I_1^3\}$	$4/3$
$\emptyset$	$II \rightarrow \{I_1^3\}$	$6/5$
$C_5$	$II^* \rightarrow \{I_1^6, I_4\}$	6
$C_3 A_1$	$III^* \rightarrow \{I_1^5, I_4\}$	4
$C_2 U_1$	$IV^* \rightarrow \{I_1^4, I_4\}$	3
$C_1$	$I_0^* \rightarrow \{I_1^2, I_4\}$	2
$A_3 \rtimes \mathbb{Z}_2$	$II^* \rightarrow \{I_1^3, I_1^*\}$	6
$A_1 U_1 \rtimes \mathbb{Z}_2$	$III^* \rightarrow \{I_1^2, I_1^*\}$	4
$U_1$	$IV^* \rightarrow \{I_1^1, I_1^*\}$	3
$A_2 \rtimes \mathbb{Z}_2$	$II^* \rightarrow \{I_1^2, IV_{Q=1}^*\}$	6
$U_1 \rtimes \mathbb{Z}_2$	$III^* \rightarrow \{I_1, IV_{Q=1}^*\}$	4
$\emptyset$	$IV_{Q=1}^*$	3
$C_1$	$I_0^* \rightarrow \{I_2^3\}$	2

[Argyres, Lotito, Lü, Martone 18]

# Classification of SCFTs

## Lesson 2

**"Conjecture": SCFTs with 8 supercharges in dimensions  $3 \leq d \leq 6$  are entirely characterized by their full moduli space of vacua.**


Example:  $C_5$  theory.

# Classification of SCFTs

## Lesson 2

**"Conjecture": SCFTs with 8 supercharges in dimensions  $3 \leq d \leq 6$  are entirely characterized by their full moduli space of vacua.**

Example:  $C_5$  theory.



# Phases of SCFTs

## Lesson 3




**Phases of the theory**  $\equiv$  **Smooth loci of the MSV**  
**Phase transitions**  $\equiv$  **Singularities**

# Phases of SCFTs

## Lesson 3

Phases of the theory  $\equiv$  Smooth loci of the MSV  
 Phase transitions  $\equiv$  Singularities

Example:  $d = 6$ ,  $\mathcal{N} = (1, 0)$  SCFT from  $SU(2)$  with  $N_f = 4$ . At the SCFT point, Spin(7) symmetry! [AB, Grimminger, 2022]




- 1 Introduction and Examples
- 2 Definition of Conical Symplectic Singularities
- 3 Moduli Space of Vacua
  - SCFTs and Moduli Space of Vacua
  - Classification of SCFTs
  - Phases of SCFTs
- 4 CSS stratification and fibrations
  - General Strategy
  - Phase diagrams from Magnetic Quivers
  - The quest for the elementary building blocks
- 5 Conclusion



# General Strategy

- 1 Fibration of the full moduli space:



- 2 For each fiber, compute singular stratification.
- 3 The phase diagram is obtained by deleting non-singular transitions.

[AB, Grimminger, 22]

- 1 Fibration of the full moduli space:


$$\begin{array}{ccc} \text{CSS} & \longrightarrow & \text{MSV} \\ & & \downarrow \\ & & \text{Base} \end{array}$$

- 2 For each fiber, **compute singular stratification**. How to do this?
- 3 The phase diagram is obtained by deleting non-singular transitions.

[AB, Grimminger, 22]

# Magnetic Quivers


The Coulomb branch of a 3d  $\mathcal{N} = 4$  SCFT is a CSS due to  $\mathfrak{su}(2)_C$ .



[Cremonesi, Hanany, Zaffaroni 14] , [Bullimore, Dimofte, Gaiotto 15] , [Braverman, Finkelberg, Nakajima 15]

# Magnetic Quivers

The Coulomb branch of a 3d  $\mathcal{N} = 4$  SCFT is a CSS due to  $\mathfrak{su}(2)_C$ .




[Cremonesi, Hanany, Zaffaroni 14] , [Bullimore, Dimofte, Gaiotto 15] , [Braverman, Finkelberg, Nakajima 15]

Let  $X$  be a CSS. We say that the (generalized) quiver  $Q$  is a **magnetic quiver** for  $X$  if

$$\mathcal{C}^{3d \mathcal{N}=4}(Q) = X.$$

# Magnetic Quivers

The Coulomb branch of a 3d  $\mathcal{N} = 4$  SCFT is a CSS due to  $\mathfrak{su}(2)_C$ .



[Cremonesi, Hanany, Zaffaroni 14] , [Bullimore, Dimofte, Gaiotto 15] , [Braverman, Finkelberg, Nakajima 15]

Let  $X$  be a CSS. We say that the (generalized) quiver  $Q$  is a **magnetic quiver** for  $X$  if

$$\mathcal{C}^{3d \mathcal{N}=4}(Q) = X.$$

## Lesson 4

**A magnetic quiver is a simple combinatorial object that encodes CSS and which gives access to its singular stratification.**

# Rank-1 4d $\mathcal{N} = 2$ magnetic quivers

Flavor	$\dim_{\mathbb{H}}(HB)$	Magnetic Quiver
$E_8$	29	Affine
$E_7$	17	
$E_6$	11	
$D_4$	5	
$A_2$	2	Dynkin
$A_1$	1	
$\emptyset$	0	

[AB, Grimminger, Hanany, Sperling, Zafir, Zhong 20]

Flavor	$\dim_{\mathbb{H}}(HB)$	Magnetic Quiver
$C_5$	16	
$C_3A_1$	8	
$C_2U_1$	4	
$C_1$	1	
$A_3 \rtimes \mathbb{Z}_2$	9	
$A_1U_1 \rtimes \mathbb{Z}_2$	3	
$U_1$	1	
$A_2 \rtimes \mathbb{Z}_2$	5	
$U_1 \rtimes \mathbb{Z}_2$	1	
$\emptyset$	0	
$C_1$	1	

# Quiver Subtraction

From a magnetic quiver, the stratification is obtained via a *quiver algorithm*.

**Physically corresponds to Higgsing graph between SCFTs.**

# Quiver Subtraction

From a magnetic quiver, the stratification is obtained via a *quiver algorithm*.

**Physically corresponds to Higgsing graph between SCFTs.**



Different algorithms:

- **Quiver subtraction** [AB, Cabrera, Grimminger, Hanany, Sperling, Zajac, Zhong, 2019]: needs a list of ICSS as an input / or alternatively a brane realization.
- **Quiver fission and decay** [AB, Sperling, Zhong, 2023]: no input needed. ICSS correspond to *stable* quivers (which can't fission or decay).

Example : 6d  $\mathcal{N} = (1, 0)$  theory from  $SU(6)$  gauge theory with  $N_f = 14$  and one antisymmetric hyper.



# Quiver Subtraction



## New elementary slices


Using brane systems / quiver fission and decay, one can construct **new Isolated Conical Symplectic Singularities!**

$$\mathcal{J}_{k_1, k_2} = \mathcal{C} \left( \begin{array}{ccc} & k_1 & k_2 \\ \circ & \leftarrow & \circ & \rightarrow & \circ \\ 1 & & 2 & & 1 \end{array} \right)$$

$k_2 \backslash k_1$	$\geq 4$	3	2	1
$\geq 4$				
3				
2				
1				bad

See also [Bellamy, Bonafé, Fu, Levy, Juteau, Sommers 2022]

# New elementary slices



Example :

$$\text{HS}(\mathcal{J}_{33}) = \frac{1 + 3t^2 + 18t^4 + 14t^6 + 18t^8 + 3t^{10} + t^{12}}{(1 - t^2)^3(1 - t^4)^3}.$$

# Outline

- 1 Introduction and Examples
- 2 Definition of Conical Symplectic Singularities
- 3 Moduli Space of Vacua
  - SCFTs and Moduli Space of Vacua
  - Classification of SCFTs
  - Phases of SCFTs
- 4 CSS stratification and fibrations
  - General Strategy
  - Phase diagrams from Magnetic Quivers
  - The quest for the elementary building blocks
- 5 Conclusion

# Summary

- **The Moduli Space of Vacua is a central object in the study of various aspects of SCFTs.**

# Summary

- **The Moduli Space of Vacua is a central object in the study of various aspects of SCFTs.**
- **It has the structure of a fibration by Conical Symplectic Singularities, which is the geometric incarnation of the phase diagram of the theory.**

# Summary

- The Moduli Space of Vacua is a central object in the study of various aspects of SCFTs.
- It has the structure of a fibration by Conical Symplectic Singularities, which is the geometric incarnation of the phase diagram of the theory.
- This structure can be uncovered using combinatorial objects called magnetic quivers.

# Summary

- The Moduli Space of Vacua is a central object in the study of various aspects of SCFTs.
- It has the structure of a fibration by Conical Symplectic Singularities, which is the geometric incarnation of the phase diagram of the theory.
- This structure can be uncovered using combinatorial objects called magnetic quivers.
- In turn, this allows to discover new phase transitions and new Conical Symplectic Singularities, of interest to physicists and mathematicians.



# Summary

- The Moduli Space of Vacua is a central object in the study of various aspects of SCFTs.
- It has the structure of a fibration by Conical Symplectic Singularities, which is the geometric incarnation of the phase diagram of the theory.
- This structure can be uncovered using combinatorial objects called magnetic quivers.
- In turn, this allows to discover new phase transitions and new Conical Symplectic Singularities, of interest to physicists and mathematicians.

Not covered today:

- Magnetic quivers can be seen as *intersection graphs* in brane systems.
- Links with (generalized) toric polygons [AB, Collinucci, Schäfer-Nameki, 22]
- RG flows between SCFTs (e.g. [AB, Giacomelli, Grimminger, 22])
- Monopole formula and Hilbert series [Cremonesi, Hanany, Zaffaroni, 14]

# Future Directions

- How to construct magnetic quivers in general (e.g. from pure geometry) ?
- List all ICSS from stable quivers.
- Monopole Formula for arbitrary transverse slices.
- Bootstrap the geometry to non-isolated CSS.
- Understand the corresponding dynamics of QFTs.
- Beyond classical geometry: scheme and stack aspects.

# Future Directions

- How to construct magnetic quivers in general (e.g. from pure geometry) ?
- List all ICSS from stable quivers.
- Monopole Formula for arbitrary transverse slices.
- Bootstrap the geometry to non-isolated CSS.
- Understand the corresponding dynamics of QFTs.
- Beyond classical geometry: scheme and stack aspects.

Thank you for your attention!