

The Power Spectral Density


Andreas Jakobsson

The power spectral density

The *power spectral density* (PSD) of a WSS stochastic process is defined as

$$\phi_y(\omega) = \sum_{k=-\infty}^{\infty} r_y(k) e^{-i\omega k}$$

over the frequencies $-\pi < \omega \leq \pi$.



The power spectral density

The *power spectral density* (PSD) of a WSS stochastic process is defined as

$$\phi_y(\omega) = \sum_{k=-\infty}^{\infty} r_y(k) e^{-i\omega k}$$

over the frequencies $-\pi < \omega \leq \pi$. The inverse transform recovers $r_y(k)$

$$r_y(k) = \frac{1}{2\pi} \int_{-\pi}^{\pi} \phi_y(\omega) e^{i\omega k} d\omega$$

It is worth noting that

$$r_y(0) = \frac{1}{2\pi} \int_{-\pi}^{\pi} \phi_y(\omega) d\omega$$

which, for a zero-mean process, measures the power of y_t , i.e.,

$$r_y(0) = E\{|y_t|^2\}$$

The PSD is *real-valued* and *non-negative*. For a real-valued process, the PSD is *symmetric*, whereas it is *non-symmetric* for a complex-valued process.

Example:

As a white noise is uncorrelated, $r_x(k) = \sigma_x^2 \delta_K(k)$, where $\delta_K(k)$ is the Kronecker delta. Thus, the PSD of a white noise is

$$\phi_x(\omega) = \sum_{k=-\infty}^{\infty} r_x(k) e^{-i\omega k} = \sigma_x^2$$

The power spectral density

Example:

Consider a sinusoidal process

$$y_t = A \cos(\omega_0 t + \phi) + w_t$$

where w_t is an AWGN. The ACF of y_t is then

$$r_y(k) = \frac{A^2}{2} \cos(\omega_0 k) + \sigma_w^2 \delta_K(k)$$

This implies that the PSD of y_t is

$$\phi_y(k) = \frac{A^2}{4} \delta_D(\omega - \omega_0) + \frac{A^2}{4} \delta_D(\omega + \omega_0) + \sigma_w^2$$

where $\sigma_D(\omega)$ is the Dirac delta, satisfying

$$f(a) = \int_{-\infty}^{\infty} f(x) \delta_D(x - a) dx$$

Estimating the power spectral density

Under the weak assumption that

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{k=-N}^N |k| |r_y(k)| = 0$$

the PSD can be expressed equivalently as

$$\phi_y(\omega) = \lim_{N \rightarrow \infty} E \left\{ \left| \frac{1}{N} \sum_{t=1}^N y_t e^{-i\omega t} \right|^2 \right\}$$

Estimating the power spectral density

Under the weak assumption that

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{k=-N}^N |k| |r_y(k)| = 0$$

the PSD can be expressed equivalently as

$$\phi_y(\omega) = \lim_{N \rightarrow \infty} E \left\{ \left| \frac{1}{N} \sum_{t=1}^N y_t e^{-i\omega t} \right|^2 \right\}$$


This suggests two natural estimators, namely as the *periodogram*

$$\hat{\phi}_p^y(\omega) = \frac{1}{N} \left| \sum_{t=1}^N y_t e^{-i\omega t} \right|^2$$

and the *correlogram*

$$\hat{\phi}_c^y(\omega) = \sum_{k=-N+1}^{N-1} \hat{r}_y(k) e^{-i\omega k}$$

Note that $\hat{r}_y(k)$ should be the *biased* estimator.



The annual number of sunspots

Estimating the power spectral density

Under the weak assumption that

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{k=-N}^N |k| |r_y(k)| = 0$$

the PSD can be expressed equivalently as

$$\phi_y(\omega) = \lim_{N \rightarrow \infty} E \left\{ \left| \frac{1}{N} \sum_{t=1}^N y_t e^{-i\omega t} \right|^2 \right\}$$

This suggests two natural estimators, namely as the *periodogram*


$$\hat{\phi}_p^y(\omega) = \frac{1}{N} \left| \sum_{t=1}^N y_t e^{-i\omega t} \right|^2$$

and the *correlogram*


$$\hat{\phi}_c^y(\omega) = \sum_{k=-N+1}^{N-1} \hat{r}_y(k) e^{-i\omega k}$$

Note that $\hat{r}_y(k)$ should be the *biased* estimator.

These estimators are equivalent, and are *asymptotically* unbiased, but not consistent. Notably, the variance of the estimate is about the same as the true spectrum squared.



Zero-padding



Generally, it is preferable to *zero-pad* the data before computing the periodogram estimate. Adding zeros does obviously not yield new information as such, but it interpolates the spectral estimate, yielding an estimate on a finer grid, often revealing features that are not otherwise visible in the estimate.

Windowing

It is often preferable to use some other time or lag window, for the *periodogram*

$$\hat{\phi}_y^p(\omega) = \frac{1}{N} \left| \sum_{t=1}^N y_t e^{-i\omega t} \right|^2 = \frac{1}{N} \left| \sum_{t=-\infty}^{\infty} v_t y_t e^{-i\omega t} \right|^2$$

and the *correlogram*

$$\hat{\phi}_y^c(\omega) = \sum_{k=-N+1}^{N-1} \hat{r}_y(k) e^{-i\omega k} = \sum_{k=-\infty}^{\infty} w_k \hat{r}_y(k) e^{-i\omega k}$$

where the *time* and *lag* windows, v_t and w_k , take the value 1 in the given range, and zero otherwise.

Such windows widens the main lobe, but lowers the sidelobes.

Windowing

It is often preferable to use some other time or lag window, for the *periodogram*



$$\hat{\phi}_y^p(\omega) = \frac{1}{N} \left| \sum_{t=1}^N y_t e^{-i\omega t} \right|^2 = \frac{1}{N} \left| \sum_{t=-\infty}^{\infty} v_t y_t e^{-i\omega t} \right|^2$$

and the *correlogram*

$$\hat{\phi}_y^c(\omega) = \sum_{k=-N+1}^{N-1} \hat{r}_y(k) e^{-i\omega k} = \sum_{k=-\infty}^{\infty} w_k \hat{r}_y(k) e^{-i\omega k}$$

where the *time* and *lag* windows, v_t and w_k , take the value 1 in the given range, and zero otherwise.

Such windows widens the main lobe, but lowers the sidelobes.



Windowing

It is often preferable to use some other time or lag window, for the *periodogram*

$$\hat{\phi}_y^p(\omega) = \frac{1}{N} \left| \sum_{t=1}^N y_t e^{-i\omega t} \right|^2 = \frac{1}{N} \left| \sum_{t=-\infty}^{\infty} v_t y_t e^{-i\omega t} \right|^2$$


and the *correlogram*

$$\hat{\phi}_y^c(\omega) = \sum_{k=-N+1}^{N-1} \hat{r}_y(k) e^{-i\omega k} = \sum_{k=-\infty}^{\infty} w_k \hat{r}_y(k) e^{-i\omega k}$$

where the *time* and *lag* windows, v_t and w_k , take the value 1 in the given range, and zero otherwise.



Such windows widens the main lobe, but lowers the sidelobes.

Window Name	Defining Equation	Approx. Main Lobe Width (radians)	Sidelobe Level (dB)
Rectangular	$w(k) = 1$	$2\pi/M$	-13
Bartlett	$w(k) = \frac{M-k}{M}$	$4\pi/M$	-25
Hanning	$w(k) = 0.5 + 0.5 \cos\left(\frac{2\pi k}{M}\right)$	$4\pi/M$	-31
Hamming	$w(k) = 0.54 + 0.46 \cos\left(\frac{2\pi k}{M-1}\right)$	$4\pi/M$	-41
Blackman	$w(k) = 0.42 + 0.5 \cos\left(\frac{2\pi k}{M-1}\right) + 0.08 \cos\left(\frac{4\pi k}{M-1}\right)$	$6\pi/M$	-57




You can read more in the excellent textbook by Stoica and Moses (2005).

Voiced speech



High-resolution estimators





Periodogram



Capon



Synthetic aperture radar



The spectrogram

