

WILL Part II

Relational Cosmology and Galactic Dynamics

Anton Rize
antonrize@willrg.com

January 2026

Abstract

In this work, we apply the principles of WILL Relational Geometry (RG) to the domain of cosmology and galactic dynamics. By enforcing geometric closure conditions with **zero free parameters**, we establish a sequential unbroken chain of derivations from first principles to observational evidence.

1. **Hubble parameter:** we derive ($H_0 \approx 68.15$ km/s/Mpc) solely from the CMB temperature and the fine-structure constant (α) providing the direct bridge between scales and landing within 1% of Planck 2018 mission measurement ($H_0 \approx 67.4$ km/s/Mpc).
2. **Distant Supernova Flux Levels:** using this H_0 value along with kinetic and potential geometric weights we compare the curve with **Pantheon+** dataset - deviations remains below 0.015 mag across the entire redshift range.
3. **CMB Acoustic Spectrum:** utilizing same derived horizon scale, we reconstruct the CMB acoustic spectrum as the resonant harmonics of an S^2 topology loaded by $\approx 4.2\%$ baryons, naturally resolving the "Low Quadrupole" anomaly via vacuum tension.
4. **Galactic Rotation Curves:** we translate the global horizon into a local acceleration scale ($a_\kappa = cH_0/3\pi$), which we apply to the SPARC database (175 galaxies). This rigid geometric prescription predicts rotation curves, the Radial Acceleration Relation (RAR), and the linear scaling of Phantom Inertia with equal or higher precision than MOND phenomenology, without any fitting.
5. **Dark Lensing:** we extend this framework to the gravitational lensing of "dark" potentials showing that the same Phantom Inertia responsible for galactic dynamics, also stands behind the "dark lensing" phenomena.
6. **Wide Binary Stars:** applied to this dynamic systems the theory correctly predicts the kinetic resonance scale ($a_\beta = cH_0/6\pi$) matching recent Gaia DR3 within observational uncertainties.

These are only few results achieved by RG in macro scale domain. They showing that the framework reproduces the considered empirical constraints across approximately 20 orders of magnitude. Taken together, these results *motivate* considering a **relational-geometric ontology** as a viable alternative to dark-sector phenomenology, and they *warrant further* empirical stress-testing across independent datasets.

Contents

1	Stage I: Galactic Dynamics	5
1.1	Main Results Established in WILL RG Part I (Recap)	5
1.2	The Relational Carriers: S^1 and S^2	6
1.3	The Energetic Closure Condition	7
1.4	Total Relational Shift and Self-Centering Reciprocity	7
2	The Necessity of Global Resonance	8
3	Deriving H_0 from CMB Temperature and α	9
3.1	Prerequisite: The Geometric Identity of α (Summary of Part III)	9
3.2	Input Parameters and Constants	9
3.3	Step-by-Step Derivation	9
3.4	Step 1: Radiation Density Calculation (ρ_γ)	9
3.5	Step 2: Maximal Structural Density	10
3.6	Step 3: The Hubble Parameter (H_0)	10
3.7	Results and Numerical Calculation	10
3.8	Unit Conversion	11
4	Galactic Dynamics: The Resonant Horizon	11
4.1	The Fundamental Tone (f_0)	11
4.2	Bifurcation of Resonance: Structural vs. Kinetic	11
4.3	Geometric Scale Coupling (The Resonant Bridge)	12
4.4	Conclusion	13
5	Derivation of the Baryonic Tully-Fisher Relation	13
5.1	Asymptotic Limit of the Horizon Resonance	13
5.2	Properties of the Derived Relation	13
5.3	Structural Distinction from MOND	14
5.4	Comparison with Observed BTFR	15
6	Protocol Independent Models Comparison	15
6.1	Motivation and Protocol	15
6.2	Data	15
6.3	Baryonic Reference Model	15
6.4	Dynamical Prescriptions Evaluated	16
6.4.1	Understanding the Metrics:	17
6.4.2	Analysis of Gas-Dominated Systems	18
7	The Universal Radial Acceleration Relation (RAR)	19
7.1	The Zero-Parameter Prediction	19
7.2	Statistical Validation	20
8	Local Verification: The Solar System Test	21
8.1	Inputs: Zero Free Parameters	21
8.2	The Prediction	21
8.3	Calculation	21
8.4	Result	21

9	The Kinetic Resonance: Wide Binary Anomaly	22
9.1	The Problem: Breakdown of Newton in the Solar Neighbourhood	22
9.2	Empirical Verification against Gaia DR3	22
9.3	Conclusion regarding Local Dynamics	23
10	The Baryonic Escape Threshold	24
10.1	Derivation of the Transition Scale (R_{trans})	24
10.2	The Physics of the Equivalence Point	24
10.3	Methodology of the "Bullseye" Test	24
10.4	Results	25
10.5	Robustness	26
11	Galactic Scale Theorem: The Topological Ruler	26
11.1	The Inversion of Distance	26
11.2	Empirical Validation: The SPARC Database Test	26
11.3	Analytical Anatomy of the Scatter: The 4th Power Sensitivity	27
12	Gravitational Lensing	28
12.1	Exact Algebraic Lensing of the Resonant Horizon (Replacing Dark Matter Lensing)	28
12.1.1	The Emergence of the "Isothermal Halo" Constant	28
12.1.2	The Unified Lensing Equation with Horizon Resonance	29
12.2	External Validation: The SLACS Survey	29
12.3	Weak Lensing: Future Work	29
13	Stage II: Relational Cosmology	30
14	Phase Evolution and the Geometric Origin of the CMB	30
14.1	The Mechanism of Expansion: Cumulative Phase Divergence	30
14.2	Derivation of the Linear Divergence Rate	30
14.3	The Phase Horizon Depth (ϕ_{max})	31
14.4	Derivation of the Scaling Exponent	31
14.5	The Topological Tautology of the Present Epoch	32
14.6	The Geometric Cooling Law	32
14.6.1	Derivation of the Scaling Exponent	33
14.6.2	Self-Consistency Check	33
14.7	The Unit Phase Transition: Geometry of Transparency	33
14.8	Prediction of the Decoupling Redshift and Temperature	34
14.9	Epistemological Hierarchy of the Recombination Epoch	34
14.10	Quantitative Deconstruction of Λ CDM Artifacts	35
14.11	Assessment of the Prediction	35
14.12	Discussion	36
15	Geometric Expansion Law: Distant Supernova Flux Levels Test	36
15.1	Geometric Partitioning of the Energy Budget	37
15.2	Derivation of the WILL–Friedmann Evolution Equation	37
15.3	The Hubble Diagram Test Protocol	38
15.4	Residual Analysis and Interpretation	39
15.5	Conclusion: Strong Empirical Agreement With Geometric Nature of DE	40

16 Geometric Origin of the CMB Acoustic Spectrum	41
16.1 The Fundamental Tone (ℓ_1): Compression Phase	41
16.1.1 Derivation	41
16.2 The Topological Skeleton: S^2 Bessel Roots	41
16.3 The Second Peak Anomaly: Baryonic Hysteresis	42
16.4 The Third Peak (ℓ_3): Return to Compression	42
16.5 Results and Conclusion	42
17 Relational Approach to the Low Quadrupole Anomaly	43
17.1 The Missing Power Problem	43
17.2 Structural tension as a High-Pass Filter	44
17.3 Quantitative Derivation of the Inertial Corridor	44
17.4 Comparison with Observation	45
17.5 The "Axis of Evil": Explaining the Alignments	46
17.6 Nodal Coupling on a Tensioned Surface	46
18 Local Cosmological Term Λ in RG (Challenging "Dark Energy")	46
18.1 Derivation of Vacuum Density	47
18.2 Derivation of Vacuum Pressure (Equation of State)	48
18.3 Legacy Correspondence: Mapping to General Relativity	48
19 The Geometric Origin of Galactic Rotation Curves (Challenging "Dark Matter")	49
19.1 The Vacuum-Dynamic Equivalence	49
20 Consolidating Mach's Principle: Derivation of Electron Mass: The Geometric Capacity Resonance	50
20.1 Topological Invariants	50
20.2 Derivation of the Topological Operator Γ	51
20.3 Derivation	52
20.4 Numerical Verification	52
20.5 Conclusion	52
21 General Discussion	53
21.1 Orthogonal Validation: The Geometric Invariant α	53
21.2 Path A: The Thermodynamic Limit (H_0)	53
21.3 Path B: The Topological Resonance (ℓ_{vac})	53
21.4 The Independence Theorem	54
21.5 Sensitivity Analysis: The Major Constraints on Dark Matter	54
21.6 Final Conclusion	55

1 Stage I: Galactic Dynamics

After we established a strong foundation for Relational Geometry we will apply this framework to galactic dynamics. But first lets refresh our memory.

1.1 Main Results Established in [WILL RG Part I](#) (Recap)

Logical Chain Summary	
Methodological Constraints	
$\downarrow$ (<i>apply to existing physics</i>)	
False Separation:	$\underbrace{\text{fixed manifold} + \text{metric}}_{\text{structure}} + \underbrace{\text{fields} + \text{constants}}_{\text{dynamics}}$
$\downarrow$ (<i>by Relational Origin Principle no background allowed</i>)	
SPACETIME $\equiv$ ENERGY	
$\downarrow$ (<i>derive consequences</i>)	
Closure + Conservation + Isotropy	
$\downarrow$ (<i>derive primal relational carriers</i>)	
S^1 (1-DOF kinematic) + S^2 (2-DOF potential)	
$\downarrow$ (<i>By Duality of Relation Lemma</i>)	
Conservation of Relation	$\underbrace{\text{Amplitude}^2}_{\text{External Interaction}} + \underbrace{\text{Phase}^2}_{\text{Internal Existence}} = 1$
$\downarrow$ (<i>only available distinguishing property</i>)	
DOF ratio = 2:1 exchange rate	
$\downarrow$ (<i>By DOF-Indifference Lemma: equal quadratic weight to each independent DOF</i>)	
$\kappa^2 = 2\beta^2 \equiv \underbrace{1\text{Amplitude}^2}_{\text{on } S^2} = \underbrace{2\text{Amplitude}^2}_{\text{on } S^1}$	
$\downarrow$ (<i>By Relational Closure and Invariance theorems</i>)	
Energy Symmetry Law $\Delta E_{A \rightarrow B} + \Delta E_{B \rightarrow A} = 0$	

Standard cosmological models operate on a descriptive paradigm, fitting dynamical laws (Lagrangians) onto a pre-existing spacetime manifold. We adopt a **generative** approach based on the [foundational principle established in WILL Relational Geometry \(Part I\)](#):

$$\boxed{\text{SPACETIME} \equiv \text{ENERGY}}$$

Throughout this paper, the identity is to be read as an ontological identification,
not as an algebraic equation or a dynamical law.

This principle asserts that "spacetime" and "[energy](#)" are not distinct entities but dual projections of a single invariant relational structure. Consequently, we do not postulate a background metric. Instead, geometry emerges solely from the conservation [requirements](#) of closed relational carriers.

1.2 The Relational Carriers: S^1 and S^2

The topology of a closed, maximally symmetric system admits exactly two minimal relational **carriers** for the **energy budget**:

1. **Kinematic Carrier (S^1)**: Encodes directional transformation (1 Degree of Freedom). Its state is defined by the orthogonal projections:

$$\beta^2 + \beta_Y^2 = 1$$

where $\beta = v/c$ is the kinematic amplitude (external motion) and $\beta_Y = 1/\gamma$ is the phase (internal time rate).

2. **Potential Carrier (S^2)**: Encodes omnidirectional interaction (2 Degrees of Freedom). Its state is defined by:


$$\kappa^2 + \kappa_X^2 = 1$$

where $\kappa^2 = R_s/r$ represents the gravitational potential intensity and κ_X is the structural phase (gravitational redshift).

Table 1: Unified representation of relativistic and gravitational effects for closed systems.

$z_\beta = \text{transverse Doppler shift, } z_\kappa = \text{gravitational redshift,}$ $\theta_1 = \arccos(\beta), \quad \theta_2 = \arcsin(\kappa), \quad \kappa^2 = 2\beta^2$	
Algebraic Form	Trigonometric Form
$\beta = v/c = \sqrt{1 - (1 + z_\beta)^{-2}}$	$\beta = \cos(\theta_1)$
$\kappa = \sqrt{R_s/r} = \sqrt{1 - (1 + z_\kappa)^{-2}}$	$\kappa = \sin(\theta_2)$
$\beta_Y = \sqrt{1 - \beta^2}$	$\beta_Y = \sin(\theta_1) = \sin(\arccos(\beta))$
$\kappa_X = \sqrt{1 - \kappa^2}$	$\kappa_X = \cos(\theta_2) = \cos(\arcsin(\kappa))$
$p = E_0/c \cdot \beta/\beta_Y$	$p = E_0/c \cdot \cot(\theta_1)$
$p_g = E_0/c \cdot \kappa/\kappa_X$	$p_g = E_0/c \cdot \tan(\theta_2)$
$\tau = \beta_Y \kappa_X$	$\tau = \sin(\theta_1) \cos(\theta_2)$
$Q = \sqrt{\kappa^2 + \beta^2} = \sqrt{3}\beta$	$Q = \sqrt{3} \cos(\theta_1)$

Clear Relational Symmetry Between Kinematic and Potential Projections derived in WILL RG I



1.3 The Energetic Closure Condition

For any stable, self-contained system, the energy budget must be partitioned between these carriers. The exchange rate is determined by the ratio of their **relational degrees of freedom** ($\text{DOF}_{S^2}/\text{DOF}_{S^1} = 2$). This yields the **Closure Theorem** governing all bound systems (relational origin of virial equilibrium):

$$\kappa^2 = 2\beta^2$$

This geometric identity replaces the Newtonian dynamical postulate. It implies that for any closed system, the gravitational potential energy (κ^2) must be exactly twice the kinetic energy (β^2) to maintain topological stability.

1.4 Total Relational Shift and Self-Centering Reciprocity

Interaction between observers is defined by the **Total Relational Shift (Q)**, which measures the magnitude of displacement from the observer's origin on a (β, κ) plane (1):

$$Q^2 = \beta^2 + \kappa^2 \quad (1)$$

Under the closure condition (1.3), this norm simplifies to $Q^2 = 3\beta^2 = \frac{3}{2}\kappa^2$.

The **Potential Relational Weight** of the total shift is: $\Omega_{pot} = \frac{\kappa^2}{Q^2} = \frac{2}{3}$. The **Kinematic Relational Weight** of the total shift is: $\Omega_{kin} = \frac{\beta^2}{Q^2} = \frac{1}{3}$.




Figure 1: **Relational Self-Centering.** The relational shift Q is defined by the orthogonal projections β and κ . Interaction is causal only when the center of the Object lies within the Observer's horizon ($Q < 1$), ensuring mutual coverage.

2 The Necessity of Global Resonance

Remark 2.1. *Throughout this document, the word Universe means the relational system within the observer's causal horizon.*

As we established in [WILL Part I](#) energy and spacetime are not separate [ontological elements](#) but a single relational structure we call **WILL** $\equiv$ **SPACE-TIME-ENERGY**. Guided by our methodological principles: [Epistemic Hygiene](#), [Relational Origin](#), [Ontological Minimalism](#) and [Mathematical Transparency](#) We will apply Relational Geometry globally.

Since **WILL** is relationally closed and boundary-free ([WILL RG Part I: Lemma Closure](#)), any causal propagation along **WILL** is recurrent. Therefore the local [relational state](#) generically includes contributions from its own propagated history (echoes).

Lemma 2.2 (Inevitability of Self-Interaction). *In a closed relational system **WILL** ([thm:relational closure](#)) with invariant measure of [energy resource](#) ([thm:relational invariance](#)), any [relational perturbation](#) cannot propagate indefinitely without re-encountering itself.*

Proof. Since **WILL** is relationally closed and boundary-free ([WILL RG Part I: Lemma Closure](#)), any causal propagation along **WILL** is recurrent. Therefore the local [relational state](#) generically includes contributions from its own propagated history (echoes). $\square$

Theorem 2.3 (Phase Closure). *A system in a stable, stationary state must be represented by a geometry that is topologically closed. All relational phases describing the system must return to their initial values after a complete cycle, ensuring the continuity and integrity of the configuration.*

Proof. Established in [WILL Part I Relational Closure](#) dictates that any stable, self-contained system must be described by a closed and self-consistent relational geometry. By Inevitability of Self-Interaction [2.2 Lemma](#) the only stable modes are the ones that completed the full cycle and came back exactly into the same phase effectively phase closing there causal propagation. In order to come back in exactly the same phase the winding number has to be a whole integer. By [Theorem \(Relational Invariance\)](#) the relational measure is conserved, so a closing n -winding configuration neither grows nor decays: it is a stable persistent mode, not a transient. By [Theorem \(Relational Isotropy\)](#) the carrier is uniform, so the n -fold configuration runs at exactly n times the rate of the lowest one, giving the evenly spaced harmonic spectrum

$$f_n = n f_0, \quad n = 1, 2, 3, \dots$$

The lowest closing mode, $n = 1$ — a single complete cycle of state transformation spanning the entire closed **WILL** — is the irreducible persistent excitation; every other persistent mode is an integer repetition of it. $\square$

Interactive visualisation

<https://willrg.com/quantazation.html>

This implies that a single mode oscillation corresponds to the Fundamental Tone of the observable Universe. Our methodology precludes the acceptance of empirically fitted values (like H_0); instead, we must derive it from first principles. To find this exact Tone, we rely on the core principle of WILL RG:

$$\text{SPACETIME} \equiv \text{ENERGY}$$

3 Deriving H_0 from CMB Temperature and α

The most robust, model-independent absolute scale available in cosmology is the monopole temperature of the Cosmic Microwave Background (T_0). It defines the current radiation energy density ρ_γ .

As derived in WILL RG Part III, the fine-structure constant α acts as the kinematic projection of the ground state ($\beta_1 \equiv \alpha$), connecting micro- and macro-closure. This fixes the RG scaling ratio between the radiation density ρ_γ and the geometric saturation density ρ_{max} (derived in WILL Part I as density at $r = R_s \implies \kappa^2 = 1$).

3.1 Prerequisite: The Geometric Identity of α (Summary of Part III)

In [WILL RG III \(click to see derivation\)](#) atomic stability is derived not from force equilibrium, but from the **Geometric Closure Condition** between the potential (S^2) and kinetic (S^1) carriers: $\kappa^2 = 2\beta^2$.

We defined the **Electromagnetic Critical Radius** (R_q) as the scale of energetic saturation ($U = E_{rest}/2$), and the **Bohr Radius** (a_0) as the scale of topological phase closure ($n = 1$). The Fine Structure Constant α is rigorously identified as the unique **Kinetic Projection** (β_1) required to bridge these two scales: [Derived in WILL RG III \(click to see derivation\)](#)

$$\alpha \equiv \beta_1 = \sqrt{\frac{1}{2} \frac{R_q}{a_0}} \quad (2)$$

Thus, α is the scaling factor defining the ratio between the **critical limit of the field** and the **stable state of matter**. This rigid geometric scaling ($\beta \rightarrow \kappa$) allows us to use α to map the macroscopic saturation density ρ_{max} in the following section.

3.2 Input Parameters and Constants

All input values are taken from standard CODATA (2018) and Planck (2018) datasets. No model-specific fitting parameters are used.

Parameter	Symbol	Value	Source/Definition
CMB Temperature	T_0	2.7255	Fixsen (2009) / Planck
Fine Structure Const.	α	7.29735e-3	CODATA ($\approx 1/137.036$)
Gravitational Const.	G	6.674e-11	CODATA
Speed of Light	c	2.99792e8	Exact
Stefan-Boltzmann Const.	σ_{SB}	5.67037e-84	Derived from fundamental constants height

Table 2: Core inputs for the calculation. Colab notebook link: [H₀ from T_{CMB} and \$\alpha\$.ipynb](#)

3.3 Step-by-Step Derivation

3.4 Step 1: Radiation Density Calculation (ρ_γ)

Objective: Determine the photon mass-energy density of the Universe. This provides the "Energy" input for the *Spacetime* $\equiv$ *Energy* equivalence.

$$\rho_\gamma = \frac{4\sigma_{SB}T_0^4}{c^3} \quad (3)$$

Using the input $T_0 = 2.7255$ K:

$$\rho_\gamma \approx 4.64495 \times 10^{-31} \text{ kg/m}^3 \quad (4)$$

3.5 Step 2: Maximal Structural Density

From the [Closure Theorem](#) we have the invariant relation $\kappa^2 = 2\beta^2$. The total relational shift of the system is therefore

$$Q^2 = \kappa^2 + \beta^2 = 3\beta^2.$$

For the photon gas we identify the kinematic projection with the ground-state value established in Part III:

$$\beta \equiv \alpha.$$

It follows that

$$Q^2 = 3\alpha^2.$$

Photons are intrinsically kinetic and do not possess a rest frame; they constitute a dynamical component of the Universe's total energy density distributed across both relational carriers. Consequently, the radiation energy density ρ_γ is normalized by the full relational shift:

$$\rho_\gamma = Q^2 \rho_{\max} = 3\alpha^2 \rho_{\max}.$$

Solving for the saturation density gives

$$\boxed{\rho_{\max} = \frac{\rho_\gamma}{3\alpha^2}}.$$

Using the observed CMB temperature $T_0 = 2.7255$ K we obtain $\rho_\gamma \approx 4.645 \times 10^{-31} \text{ kg m}^{-3}$. With $\alpha \approx 7.297 \times 10^{-3}$ this yields

$$\rho_{\max} \approx 2.9075 \times 10^{-27} \text{ kg m}^{-3}.$$

3.6 Step 3: The Hubble Parameter (H_0)

Objective: Convert the saturation density into the frequency parameter.

Logic: Using the WILL RG saturation identity $\rho_{\max}(r) = c^2/(8\pi Gr^2)$ together with the horizon definition $H_0 = c/r$, we obtain the relation:

$$H_0 = \sqrt{8\pi G \rho_{\max}}. \quad (5)$$

3.7 Results and Numerical Calculation

Substituting the derived $\rho_{\max}$ into the final equation:

$$\begin{aligned} H_0 &= \sqrt{8\pi \cdot (6.674 \times 10^{-11}) \cdot (2.9075 \times 10^{-27})} \\ H_0 &= \sqrt{4.877 \times 10^{-36}} \\ H_0 &\approx 2.2084 \times 10^{-18} \text{ s}^{-1} \end{aligned}$$

3.8 Unit Conversion

Converting from SI units (s^{-1}) to standard cosmological units (km/s/Mpc):

$$\text{Conversion Factor} = \frac{3.0857 \times 10^{22} \text{ m/Mpc}}{1000 \text{ m/km}} \approx 3.0857 \times 10^{19}$$

$$H_0 \approx 2.2084 \times 10^{-18} \times 3.0857 \times 10^{19} \approx \mathbf{68.15 \text{ km/s/Mpc}} \quad (6)$$

Desmos and Colab

Desmos: [DERIVATION OF \$H_0\$ AND SUN'S ROTATION VELOCITY](#) Colab: [from \$T_{CMB}\$ and alpha.ipynb](#)

4 Galactic Dynamics: The Resonant Horizon

We derive galactic dynamics from the principle that **Spacetime** $\equiv$ **Energy**. In RG, "distance" is not a spatial separation in a void, but a difference in energy configurations. Consequently, radial separation r must be expressible as a frequency potential relative to the Global Horizon.

4.1 The Fundamental Tone (f_0)

Since the Universe is a topologically closed system with a causal horizon $R_H = c/H_0$, it possesses a minimum energy state corresponding to the fundamental standing wave (The Fundamental Tone):

$$f_0 = \frac{c}{2\pi R_H} = \frac{H_0}{2\pi}. \quad (7)$$

This frequency establishes the minimal energy floor for any interaction in the cosmos. Associated with this tone is the **Machian Acceleration Scale**:

$$a_{Mach} = f_0 c = \frac{H_0 c}{2\pi} \approx 1.05 \times 10^{-10} \text{ m/s}^2. \quad (8)$$

4.2 Bifurcation of Resonance: Structural vs. Kinetic

Recall the total relational shift established in WILL Part I:

$$Q^2 = \kappa^2 + \beta^2 = 3\beta^2 = \frac{3}{2}\kappa^2 \quad (9)$$

1. **Galaxies** are physically realised as continuous potential fields (fluids/structure). Their relational state is a smooth function $\kappa(\vec{x})$ on the S^2 carrier. Hence, they couple to the horizon via the **Structural Channel** with weight $\Omega_{pot} = \frac{\kappa^2}{Q^2} = \frac{2}{3}$.

$$a_\kappa = \Omega_{pot} \cdot a_{Mach} = \frac{2}{3} \frac{cH_0}{2\pi} = \frac{cH_0}{3\pi} \approx 0.70 \times 10^{-10} \text{ m/s}^2 \quad (10)$$

2. **Binaries** are discrete orbital systems (point masses). Their relational state is a periodic function $\beta(\theta)$ (where θ is the orbital phase) on the S^1 carrier. Hence, they couple via the **Kinetic Channel** with weight $\Omega_{kin} = \frac{\beta^2}{Q^2} = \frac{1}{3}$.

$$a_\beta = \Omega_{kin} \cdot a_{Mach} = \frac{1}{3} \frac{cH_0}{2\pi} = \frac{cH_0}{6\pi} \approx 0.35 \times 10^{-10} \text{ m/s}^2 \quad (11)$$

This assignment is enforced by the algebraic closure condition $\kappa^2 = 2\beta^2$. The continuity of the potential field selects the S^2 carrier for galaxies, while the discrete orbital nature selects S^1 for binaries.

Consequently, the resulting acceleration scales $a_\kappa = cH_0/3\pi$ and $a_\beta = cH_0/6\pi$ are **topological invariants** of the theory. This bifurcation rigorously explains why MOND's single universal parameter a_0 fails for wide binaries (which require the kinetic scale) while WILL RG accurately matches both regimes.

4.3 Geometric Scale Coupling (The Resonant Bridge)

According to the **Global Phase-Closure Constraint** (Theorem ??), any local relational shift must be geometrically coherent with the Fundamental Tone of the Universe to persist. The local system does not evolve in an absolute void; it is embedded within the structural capacity of the global horizon.

At any radius r , the system is characterized by two boundary energy scales:

1. **The Local Capacity (E_{loc}):** The intrinsic relational shift generated by the local baryonic mass.

$$E_{loc} = v_N^2 \quad (12)$$

2. **The Global Capacity (E_{glob}):** The energy floor enforced by the Fundamental Tone at that distance, weighted by the topological carrier Ω (where $\Omega = \Omega_{pot}$ or Ω_{kin}).

$$E_{glob} = \Omega \cdot a_{Mach} \cdot r \quad (13)$$

To maintain global phase closure, the local configuration must physically couple to the global boundary. In a relational ontology lacking absolute metrics, this coupling must take the form of an invariant geometric projection. The local state induces a **Resonant Shift (E_{res})** in the geometric fabric.

For the combined mode to remain structurally stable (resonant) across all scales, the relational projection of the local capacity onto the resonant shift must exactly mirror the projection of the resonant shift onto the global capacity. This defines a strict geometric proportion:

$$\frac{E_{loc}}{E_{res}} = \frac{E_{res}}{E_{glob}} \quad (14)$$

Solving this proportion yields the resonant bridge as the exact geometric mean of the two boundary scales:

$$E_{res}^2 = E_{loc} \cdot E_{glob} \implies E_{res} = \sqrt{v_N^2 \cdot (\Omega a_{Mach} r)} \quad (15)$$

Because the system is fully resonant ($\Delta_\phi = 2\pi n$), the total observed relational shift (v_{obs}^2) is the direct sum of the intrinsic local capacity and the induced resonant bridge:

$$v_{obs}^2 = E_{loc} + E_{res} \quad (16)$$

$$v_{obs}^2 = \underbrace{v_N^2}_{\text{Intrinsic local capacity}} + \underbrace{\sqrt{v_N^2 \cdot (\Omega a_{Mach} r)}}_{\text{Resonant bridge}} \quad (17)$$

where $\Omega = \Omega_{kin}$ or Ω_{pot} depends on either structural or dynamical coupling of the system.

$$\boxed{v_{obs}^2 = v_N^2 + \sqrt{v_N^2 \cdot (\Omega a_{Mach} r)}} \quad (18)$$

4.4 Conclusion

This equation derives the flat rotation curves of galaxies without invoking Dark Matter. The "extra" velocity is simply the physical manifestation of global phase closure between the local orbital frequency and the Universe's fundamental tone. The rotation curve remains flat because the system cannot decay below the energy floor supported by the global resonance.

5 Derivation of the Baryonic Tully-Fisher Relation

The Baryonic Tully-Fisher Relation (BTFR) is one of the tightest empirical correlations in extragalactic astronomy: the asymptotic rotation velocity of a disk galaxy scales as $V_{flat}^4 \propto M_b$, where M_b is the total baryonic mass. We derive this scaling, including its slope and normalization, directly from the Resonant Bridge established in Section 4.

5.1 Asymptotic Limit of the Horizon Resonance

The total observed velocity at radius r from a baryonic mass M_b is given by the Resonant Bridge equation:

$$V_{obs}^2(r) = V_N^2(r) + \sqrt{V_N^2(r) \cdot a_\kappa \cdot r}, \quad (19)$$

where $V_N^2 = GM_b/r$ is the Newtonian circular velocity for a point mass, and $a_\kappa = cH_0/(3\pi)$ is the structural resonance scale.

A key algebraic property of this equation emerges for the point-mass limit. The Resonant Bridge term simplifies to:

$$\sqrt{V_N^2 \cdot a_\kappa \cdot r} = \sqrt{\frac{GM_b}{r} \cdot a_\kappa \cdot r} = \sqrt{GM_b \cdot a_\kappa} = \text{const.} \quad (20)$$

The r -dependence cancels identically. The Horizon resonance contribution is therefore a **constant velocity increment** at all radii, which provides a natural mechanism for asymptotically flat rotation curves.

The full velocity profile for a point mass thus takes the form:

$$V_{obs}^2(r) = \frac{GM_b}{r} + \sqrt{GM_b \cdot a_\kappa}. \quad (21)$$

As $r \rightarrow \infty$, the Newtonian term vanishes, yielding the asymptotic flat velocity:

$$V_{flat}^2 = (GM_b \cdot a_\kappa)^{1/2}. \quad (22)$$

Squaring both sides gives the **Baryonic Tully-Fisher Relation**:

$$\boxed{V_{flat}^4 = G \cdot a_\kappa \cdot M_b} \quad (23)$$

5.2 Properties of the Derived Relation

1. Slope. The exponent $V^4 \propto M$ (logarithmic slope of 4) is an exact structural consequence of the geometric mean form of the Horizon resonance term. This matches the observed BTFR slope of 4.0 ± 0.1 (McGaugh, 2012; Lelli et al., 2016).

2. Normalization. The asymptotic coefficient $a_\kappa = cH_0/(3\pi) \approx 7.0 \times 10^{-11} \text{ m/s}^2$ is derived from first principles with zero free parameters.

3. Effective coefficient at finite radius. The BTFR is observationally measured at the outermost reliable data point, not at $r \rightarrow \infty$. At any finite radius $r = n \cdot r_t$ (where $r_t = \sqrt{\frac{3\pi}{2}} R_s R_H = \sqrt{GM_b/a_\kappa}$ is the transition radius and $R_H = c/H_0$ is the Hubble horizon), the observed velocity retains a non-negligible Newtonian contribution:

$$V_{obs}^2(n \cdot r_t) = \left(\frac{1}{n} + 1\right)(GM_b \cdot a_\kappa)^{1/2}. \quad (24)$$

The effective BTFR coefficient at this radius is:

$$a_{eff}(n) = \left(\frac{1}{n} + 1\right)^2 a_\kappa. \quad (25)$$

Measurement Radius	$(1/n + 1)^2$	$a_{eff} [\text{m/s}^2]$
$r = r_t$ (transition)	4.00	2.81×10^{-10}
$r = 2r_t$	2.25	1.58×10^{-10}
$r = 3r_t$	1.78	1.25×10^{-10}
$r = 5r_t$	1.44	1.01×10^{-10}
$r \rightarrow \infty$	1.00	0.70×10^{-10}

Table 3: Effective BTFR coefficient as a function of measurement radius, in units of the transition radius $r_t = \sqrt{GM_b/a_\kappa}$.

At a typical observational measurement radius of $r \approx 3\text{--}5r_t$, the effective coefficient falls in the range $a_{eff} \approx (1.0\text{--}1.25) \times 10^{-10} \text{ m/s}^2$. This is consistent with the empirically fitted MOND acceleration scale $a_0 \approx 1.2 \times 10^{-10} \text{ m/s}^2$, which represents neither a fundamental constant nor an asymptotic limit, but an effective average over the transition region.

5.3 Structural Distinction from MOND

Although the asymptotic scaling $V^4 \propto M$ coincides with the deep-MOND prediction, the functional form of the RG Resonant Bridge differs in a testable way. The RG equation (19) contains an explicit additive Newtonian term (V_N^2), whereas the standard MOND interpolation function $\nu(g_{bar}/a_0)$ folds both contributions into a single nonlinear expression.

This structural difference has two observational consequences:

1. The effective BTFR normalization in RG is radius-dependent, producing a slight increase in scatter for galaxies whose “flat” velocity is measured at very different multiples of r_t .
2. For fixed baryonic mass, the RG rotation curve approaches its asymptote *from above* (due to the decaying V_N^2 term), whereas MOND approaches from below in certain interpolation schemes.

These predictions are, in principle, testable with extended rotation curves reaching well beyond $5r_t$.

5.4 Comparison with Observed BTFR

The observed BTFR from gas-rich galaxy samples (McGaugh, 2012; Lelli et al., 2016) yields a normalization $M_b \approx 47 M_\odot (\text{km/s})^{-4} \cdot V_{flat}^4$. Inverting this relation corresponds to an effective acceleration scale $a_{eff} \approx 1.6 \times 10^{-10} \text{ m/s}^2$ — higher than the RG asymptotic value $a_\kappa \approx 0.70 \times 10^{-10}$, but consistent with the effective coefficient at $r \approx 2 r_t$ (Table 3).

This is physically expected: the “flat” portion of observed rotation curves is typically measured at radii of order 2–3 times the transition scale, where the Newtonian contribution remains non-negligible. The enhancement from the additive V_N^2 term is a structural feature of the RG Horizon resonance that distinguishes it from the standard MOND interpolation.

The full rotation curve analysis on the 175-galaxy SPARC sample (Section 6), which accounts for the actual radial mass distribution rather than the point-mass approximation, provides the definitive test. That analysis demonstrates a median bias of +0.53 km/s in Gas-Dominated galaxies using the derived scale a_κ with zero free parameters.

6 Protocol Independent Models Comparison

6.1 Motivation and Protocol

The central empirical challenge addressed here is the discrepancy between observed galactic rotation velocities and the predictions of Newtonian and GR gravity sourced solely by baryons. Standard analyses often employ complex error weighting, likelihood maximization, and galaxy-specific parameter tuning (e.g., varying mass-to-light ratios), which can obscure the distinction between a model’s predictive power and its parametric flexibility.

Our objective is to assess whether raw mathematics reflex physical a fixed physical prescription reproduces the kinematic structure of disk galaxies in a transparent, assumption-minimal manner. We adopt a deliberately austere protocol:

1. **No Parameter Tuning:** No free parameters are adjusted per galaxy.
2. **Fixed Mass-to-Light Ratios:** We adhere to standard population synthesis values without variation.
3. **Raw Deviation Metrics:** We evaluate raw residuals without weighting by observational uncertainties, preventing the suppression of physical systematics by error bars.

6.2 Data

We utilize the SPARC database (Table 2), comprising 175 disk galaxies. Observed circular velocities $V_{obs}(r)$ and baryonic components (V_{gas} , V_{disk} , V_{bulge}) are taken directly from the catalog. To ensure physical causality, negative baryonic velocity components (artifacts of observational noise decomposition) are truncated to zero prior to squaring.

6.3 Baryonic Reference Model

The baryonic circular velocity is defined as:

$$V_b^2(r) = V_{gas}^2(r) + \Upsilon_{disk} V_{disk}^2(r) + \Upsilon_{bulge} V_{bulge}^2(r). \quad (26)$$

Fixed Mass-to-Light Ratios: Unlike standard dark matter analyses that often treat Υ_* as a nuisance parameter to be fitted per galaxy, we enforce a strict global Stellar Population Synthesis expectations for the $3.6 \mu\text{m}$ band (Lelli et al., 2016):

$$\Upsilon_{\text{disk}} = 0.5, \quad \Upsilon_{\text{bulge}} = 0.7.$$

applied uniformly across the entire sample. This eliminates "per-galaxy tuning" completely.

6.4 Dynamical Prescriptions Evaluated

Five distinct physical prescriptions are compared.

1. Newtonian Baseline

$$V_{\text{Newt}}(r) = V_b(r).$$

2. LCDM with Abundance Matching (No Fitting)

To represent the Standard Model framework without allowing ad-hoc halo fitting (e.g., varying concentration or mass per galaxy), we employ a deterministic Abundance Matching protocol. To ensure a strict evaluation of the standard paradigm, we utilize the Planck 2018 cosmological parameters ($H_0 = 67.4 \text{ km s}^{-1} \text{ Mpc}^{-1}$, $h = 0.674$) for all halo scaling relations, independent of the derived scales tested in other models.

Stellar Mass Estimation: Total stellar mass M_* is reconstructed directly from the kinematic data. We integrate the baryonic velocity components at the outermost observed radius r_{last} assuming Newtonian dynamics, consistent with the fixed mass-to-light ratios:

$$M_* = \frac{r_{\text{last}}}{G} (\Upsilon_{\text{disk}} V_{\text{disk}}^2(r_{\text{last}}) + \Upsilon_{\text{bulge}} V_{\text{bulge}}^2(r_{\text{last}})). \quad (27)$$

Halo Assignment:

- **Halo Mass (M_{200}):** We map the estimated M_* to the virial mass M_{200} using the inverse of the stellar-to-halo mass relation (SHMR) from Moster et al. (2013) at $z = 0$.
- **Concentration (c_{200}):** The halo concentration is derived from the mass-concentration relation of Dutton & Macciò (2014), explicitly fixing the Hubble parameter to the Planck value ($h = 0.674$).
- **Velocity Profile:** The dark matter contribution is modeled as a standard NFW halo:

$$V_{\text{NFW}}^2(r) = V_{200}^2 \frac{1}{x} \frac{\ln(1 + cx) - \frac{cx}{1+cx}}{\ln(1 + c) - \frac{c}{1+c}}, \quad (28)$$

where $x = r/R_{200}$. The virial radius R_{200} and virial velocity V_{200} are calculated using the critical density defined by $H_{0,\text{Planck}}$.

The total velocity is then $V_{\text{LCDM}} = \sqrt{V_b^2 + V_{\text{NFW}}^2}$. No parameters are tuned to minimize residuals for individual galaxies.

3. MOND (Standard Benchmark)

We employ the standard interpolation function $\mu(x) = x/(1+x)$ with the canonical acceleration scale $a_0 = 1.2 \times 10^{-10} \text{ m s}^{-2}$. The prediction is given analytically by the solution to the algebraic quadratic equation:

$$V_{\text{MOND}}(r) = \sqrt{\frac{V_b^2(r) + \sqrt{V_b^4(r) + 4V_b^2(r)a_0r}}{2}}. \quad (29)$$

Note: In this benchmark, ‘MOND’ refers to the algebraic μ -prescription (a phenomenological mapping), not a physical theory. It is included only as an empirical compression baseline for RAR.

4. Emergent Gravity (Verlinde, 2016)

We test the theoretical scaling proposed by Verlinde, where the acceleration scale is determined by the Hubble parameter H_0 . Using the theoretical coefficient $1/6$:

$$a_{\text{VG}} = \frac{cH_0}{6} \approx 1.1 \times 10^{-10} \text{ m s}^{-2}. \quad (30)$$

The velocity profile follows the Deep-MOND scaling for point masses:

$$V_{\text{Verlinde}}(r) = \sqrt{V_b^2(r) + \sqrt{a_{\text{VG}}V_b^2(r)r}}. \quad (31)$$

5. WILL Relational Geometry (RG)

We test RG using Horizon Resonance equation derived in Galactic Dynamics (4) section.

$$V_{\text{RG}}(r) = \sqrt{V_b^2(r) + \sqrt{a_\kappa V_b^2(r)r}}. \quad (32)$$

This equation include 0 free parameters and no adjustable nodes. It derived directly from methodological principals of Relational Geometry.

$$a_\kappa = \frac{cH_0}{3\pi}, \quad \text{where } H_0 \equiv \sqrt{8\pi G\rho_\gamma/(3\alpha^2)}. \quad (33)$$

This yields a theoretical $H_0 \approx 68.15 \text{ km/s/Mpc}$ and $a_\kappa \approx 0.70 \times 10^{-10} \text{ m s}^{-2}$.

Results

Performance is evaluated using three robust metrics: Median Absolute Error (MedAE), Median Signed Bias (systematic offset), and the fraction of data points predicted within 10 km/s (F_{10}).

6.4.1 Understanding the Metrics:

- **MedAE (Median Absolute Error):** This indicates the typical magnitude of the error in velocity prediction, regardless of whether it’s an over-prediction or under-prediction. A lower MedAE means the model’s predictions are closer to the observed values on average.
- **Bias (Median Residual):** This measures the systematic tendency of the model to either over-predict (positive bias) or under-predict (negative bias) the observed velocities. A bias closer to 0 indicates a more accurate and less systematic error.

- **F10 (Fraction Within 10 km/s):** This is the fraction of data points where the model’s predicted velocity is within 10 km/s of the observed velocity. A higher F10 means a larger proportion of predictions are very accurate.

Table 4: Global performance metrics on the full SPARC sample ($N = 175$). Values represent the median across all galaxies. **Colab notebook link:** [Galactic Rotation Protocol Independent SPARC.ipynb](#)

Model	MedAE [km/s]	Bias [km/s]	F_{10}
Newtonian (baryons only)	38.46	+36.91	0.08
Λ CDM (Abundance Matching)	13.32	-6.83	0.36
MOND (Standard a_0)	10.43	-4.37	0.48
Verlinde ($a_0 = cH_0/6$)	12.27	-8.52	0.33
WILL RG ($a_\kappa = cH_0/3\pi$)	11.18	-2.26	0.47

6.4.2 Analysis of Gas-Dominated Systems

To isolate physical validity from stellar mass-to-light ratio uncertainties, we analyze the subset of galaxies dominated by gas [$V_{\text{gas}}^2 > (\Upsilon V_{\text{disk}}^2 + \Upsilon V_{\text{bul}}^2)$]. In this regime, the baryonic mass distribution is known with high precision.

Table 5: Performance metrics on Gas-Dominated galaxies ($\text{GasFrac} > 0.5$).

Model	MedAE [km/s]	Bias [km/s]	F_{10}
Λ CDM (AM)	7.42	-3.91	0.65
MOND (Standard)	7.70	-5.12	0.70
Verlinde (1/6)	8.04	-5.90	0.71
WILL RG (1/3 π)	7.00	+0.53	0.66

WILL RG derived from CMB temperature and α , with no tuning on galaxy dynamics, outperforms or matches the phenomenologically calibrated MOND on its own benchmark dataset, particularly in terms of bias.

Interactive web LAB:

[Galactic Dynamics](#)




Figure 2: Grey points represent raw data; black squares denote binned means with standard error of the mean (SEM). **Standard MOND** ($a_0 = 1.2 \times 10^{-10}$, blue) and **Verlinde's Emergent Gravity** ($a_0 \approx 1.1 \times 10^{-10}$, green) systematically overpredict the observed acceleration in the low-acceleration regime ($g_{bar} < 10^{-11}$), lying outside the SEM error bars. **WILL Relational Geometry** (red), utilizing a theoretically derived acceleration scale $a_\kappa = cH_0/3\pi \approx 0.7 \times 10^{-10}$, exhibiting negligible systematic bias ($+0.53$ km/s).

Colab Notebook

[Galactic Rotation Protocol Independent SPARC.ipynb](#)

7 The Universal Radial Acceleration Relation (RAR)

We subjected the WILL framework to the rigorous Radial Acceleration Relation (RAR) test using the full SPARC database (175 galaxies, > 3000 data points). Unlike standard Dark Matter models, which treat the halo as a free component with arbitrary fitting parameters for each galaxy, WILL RG predicts a rigid, universal functional relationship between the baryonic acceleration g_{bar} and the observed acceleration g_{obs} .

7.1 The Zero-Parameter Prediction

The theoretical curve is derived solely from the **Horizon Resonance**. The observed acceleration is the global phase closure principle:

$$g_{obs} = g_{bar} + \sqrt{g_{bar} \cdot a_\kappa} \quad (34)$$

Crucially, the global acceleration scale a_κ is **not fitted** to the galaxy data. It is fixed entirely by the Cosmological Anchor derived in the previous sections from the CMB temperature (T_0) and the fine-structure constant (α):

$$a_\kappa = \frac{cH_0}{3\pi} \approx 7.02 \times 10^{-11} \text{ m/s}^2 \quad (35)$$

where $H_0 \approx 68.15 \text{ km/s/Mpc}$ is the theoretically derived Hubble parameter.

7.2 Statistical Validation

Figure 3 demonstrates the resulting "Main Sequence of Galaxies". Despite the vast diversity of morphological types—ranging from gas-dominated dwarfs to massive high-surface-brightness spirals—the data collapses onto the single theoretical curve predicted by Eq. (14).

The statistical analysis of the residuals (logarithmic deviation) yields:

- **Root Mean Square Error (RMSE):** 0.065 dex.
- **Mean Offset:** 0.007 dex (approx. 1.5%).

Conclusion: The theory matches observations with near zero systematic bias and a scatter consistent with observational uncertainties. This strongly suggests that the Horizon Resonance effect governed by the derived scale (H_0) can fully explain and accurately predict the observation phenomena that usually attributed to Dark Matter.




Figure 3: Radial Acceleration Relation (RAR) for 175 SPARC galaxies. The green dots shows the density of > 3000 individual data points. The cyan line represents the WILL Horizon Resonance prediction ($g_{obs} = g_{bar} + \sqrt{g_{bar}a_\kappa}$) using the H_0 value derived from CMB thermodynamics. The remarkable agreement (RMSE ≈ 0.065 dex) without free parameters strongly suggests that galactic dynamics are regulated by the global horizon.

Colab Notebook

[RAR test.ipynb](#)

8 Local Verification: The Solar System Test

To demonstrate the precision of RG, we apply the derived rotation speed law to our own local environment: the motion of the Sun within the Milky Way. This test is critical because it relies on high-precision local data rather than statistical ensembles.

8.1 Inputs: Zero Free Parameters

Local Baryonic Baseline (V_{bar}): According to (10) standard mass models of the Milky Way (bulge + disk + gas), the circular velocity contribution solely from visible baryonic matter at the Solar radius ($R_0 \approx 8.0$ kpc) is approximately:

$$V_{bar} \approx 170 \pm 5 \text{ km/s} \quad (36)$$

This value represents the purely Newtonian potential of the luminous galaxy, excluding any Dark Matter halo.

8.2 The Prediction

We apply the **Horizon Resonance** derived for galactic dynamics. The total observed velocity squared is the sum of the local self-energy and the resonant coupling to the horizon:

$$V_{obs}^2 = V_{bar}^2 + \sqrt{V_{bar}^2 a_\kappa R_0} \quad (37)$$

8.3 Calculation

Substituting the input values ($R_0 = 8.0$ kpc $\approx 2.47 \times 10^{20}$ m):

$$\begin{aligned} V_{obs}^2 &= (170)^2 + \sqrt{(170)^2 \cdot (7.02 \times 10^{-11} \cdot 2.47 \times 10^{20}) \cdot 10^{-6}} \\ &= 28900 + \sqrt{28900 \cdot 17339} \\ &= 28900 + 22380 \\ &= 51280 \text{ (km/s)}^2 \end{aligned}$$

Taking the square root yields the predicted orbital velocity:

$$V_{pred} \approx 226.4 \text{ km/s} \quad (38)$$

Desmos Project

DERIVATION OF H_0 AND SUN'S ROTATION VELOCITY

8.4 Result

The predicted velocity of ≈ 226 km/s is in excellent agreement with the IAU standard value (220 km/s) and recent Gaia kinematic derivations (229 ± 6 km/s).

Crucially, the "missing" velocity component (≈ 56 km/s), traditionally attributed to a Dark Matter halo, emerges here automatically as the **Horizon Resonance term** $\sqrt{V_{bar}^2 a_\kappa R_0}$. The galaxy is not filled with invisible matter; it is resonant with the cosmic horizon.

9 The Kinetic Resonance: Wide Binary Anomaly

9.1 The Problem: Breakdown of Newton in the Solar Neighbourhood

While the Radial Acceleration Relation (RAR) establishes the geometric link between baryons and the horizon on galactic scales (10^{20} m), a critical test of any modified gravity framework is its applicability to small-scale systems (10^{14} m) that are free from the complexities of dark matter halos and hydrodynamic gas pressure. Wide Binary Stars ($r > 2000$ AU) provide exactly such a laboratory.

Recent high-precision analyses of the Gaia DR3 catalog (3; 4) have reported a definitive breakdown of Newtonian dynamics at low accelerations ($g_N < 10^{-9}$ m/s²). However, a significant tension has emerged:

- **Newtonian Failure:** The observed gravity boost factor $\gamma = g_{obs}/g_N$ rises clearly above unity ($\gamma > 1$).
- **MOND Overprediction:** Standard Modified Newtonian Dynamics (AQUAL), tuned to galactic rotation curves ($a_0 \approx 1.2 \times 10^{-10}$), predicts a boost factor ($\gamma \approx 1.8 - 2.0$) that is significantly higher than the observed values ($\gamma \approx 1.4 - 1.6$).

To resolve this, standard MOND requires ad-hoc “External Field Effects” (EFE). In contrast, WILL Relational Geometry predicts this “weakened” anomaly naturally as a consequence of geometric bifurcation.

9.2 Empirical Verification against Gaia DR3

We test this Kinetic Resonance prediction ($a_\beta = cH_0/6\pi$) against the binned data from Chae (2023) for pure binary systems. The theoretical boost factor is calculated as:

$$\gamma_{WILL} = 1 + \sqrt{\frac{a_\beta}{g_N}} = 1 + \sqrt{\frac{cH_0}{6\pi g_N}} \quad (39)$$




Figure 4: **The Kinetic Resonance Test.** The plot compares the gravity boost factor as a function of Newtonian acceleration. **Blue dashed line:** Standard MOND prediction ($a_0 = 1.2 \times 10^{-10} \text{ m/s}^2$), which systematically overestimates the anomaly. **Red solid line:** WILL RG Kinetic Resonance prediction ($a_\beta = cH_0/6\pi \approx 0.35 \times 10^{-10} \text{ m/s}^2$) passes precisely through the observational data points, matches the reported trend of Wide Binary observations without any parameter fitting.

Colab Notebook

[WidebinaryChae2023.ipynb](#)

Numerical Comparison (Deep Regime). At the characteristic low-acceleration point $g_N = 10^{-9.8} \text{ m/s}^2$:

- **Observed (Gaia):** $\gamma \approx 1.45 - 1.55$.
- **Standard MOND:** $\gamma \approx 1.87$ (Overprediction $> 20\%$).
- **WILL RG Prediction:** $\gamma \approx 1.47$ (Strong Agreement).

9.3 Conclusion regarding Local Dynamics

The successful prediction of the Wide Binary anomaly confirms the **bifurcation of gravitational dynamics**:

- Structural systems (Galaxies) resonate via $\Omega_{pot} = 2/3$.
- Kinetic systems (Binaries) resonate via $\Omega_{kin} = 1/3$.

Both scales are governed by the single Universal Horizon parameter H_0 , unifying the dynamics of the Solar neighbourhood with the expansion of the Universe.

10 The Baryonic Escape Threshold

10.1 Derivation of the Transition Scale (R_{trans})

We proceed directly from the **Horizon Resonance** derived in the Galactic Dynamics analysis. The total observed acceleration g_{obs} is defined by the resonance between the local baryonic source g_{bar} and the global Machian background a_κ :

$$g_{obs} = g_{bar} + \sqrt{g_{bar}a_\kappa} \quad (40)$$

By expressing this relation in terms of inertial mass ($M_{obs} = r^2 g_{obs}/G$), we obtain a linear scaling law:

$$\frac{M_{obs}}{M_{bar}} = 1 + \frac{r}{R_{trans}} \quad (41)$$

The transition scale R_{trans} is determined by the geometric mean of the **local event horizon** (Schwarzschild radius R_s) and the **global cosmic horizon** (Hubble Horizon R_H). Substituting the derived Machian acceleration $a_\kappa = c^2/(3\pi R_H)$, we find:

$$R_{trans} = \sqrt{\frac{GM}{a_\kappa}} = \sqrt{\frac{3\pi}{2}} R_s R_H \quad (42)$$

This radius defines the **geometric horizon** of the galaxy: the distance where the local curvature structurally couples to the global topology.

10.2 The Physics of the Equivalence Point

WILL RG mandates a precise dynamical condition at the transition radius $r = R_{trans}$. Substituting this condition into the mass equation yields:

$$\frac{M_{obs}}{M_{bar}} = 1 + 1 = 2 \quad (43)$$

Consequently, the observed velocity must exceed the Newtonian prediction by exactly $\sqrt{2}$:

$$V_{obs} = V_{bar}\sqrt{2} \quad (44)$$

Since the Newtonian escape velocity is defined as $V_{esc} = V_{orb}\sqrt{2}$, we arrive at the identity:

$$\boxed{V_{obs}(R_{trans}) \equiv V_{esc}^{bary}} \quad (45)$$

Physical Definition: The "Dark Matter" phenomenon is the observational signature of containment. When the orbital velocity exceeds the local baryonic escape velocity ($V > V_{esc}^{bary}$), the system couples to the global Horizon, stabilizing the orbit.

10.3 Methodology of the "Bullseye" Test

We test this identity against the SPARC database (2), comprising 161 galaxies. The analysis adheres to a strict **Zero Free Parameters** protocol:

1. **Global Constants:** H_0 is fixed to the derived value of 68.15 km/s/Mpc.
2. **Fixed Astrophysics:** Stellar Mass-to-Light ratios are fixed globally ($\Upsilon_{disk} = 0.5$, $\Upsilon_{bulge} = 0.7$).
3. **Normalized Coordinates:**

- X-axis: $X = r/R_{trans}$
- Y-axis: $Y = V_{obs}/V_{esc}^{bary}$

The Prediction: The data must collapse onto the curve $Y = \sqrt{(1+X)/2}$ and pass strictly through the "Bullseye" point (1,1).

10.4 Results

Figure 5 presents the results of the analysis.




Figure 5: **The Universal Escape Threshold.** The plot shows the normalized rotation velocity ($Y = V_{obs}/V_{esc}^{bary}$) versus normalized radius ($X = r/R_{trans}$) for 161 galaxies ($N = 3007$ points). **Green points:** Raw data. **Black points:** Binned means with standard errors. **Cyan Line:** Hard Fixed WILL RG prediction. **Orange Dotted Line:** Naive WILL RG without 1.5π factor. The red crosshair marks the Equivalence Point (1,1).

Colab Notebook

[Baryonic Escape Threshold.ipynb](#)

Quantitative Analysis.

- **Equivalence Point Accuracy:** In the critical transition zone ($0.95 < X < 1.05$), the observed normalized velocity is $Y_{obs} = 0.965 \pm 0.018$. The deviation is **-3.5%** from the theoretical target. Given the $\sim 20\%$ systematic uncertainty in stellar population models, this supports the prediction.
- **Global Bias:** Across the full radial range, the mean model bias is $+1.3\%$.

10.5 Robustness

To verify that the result is not an artifact of axis interdependence:

1. **Null Hypothesis:** A Newtonian universe would yield a horizontal line at $Y \approx 0.707$. The data rises to $Y = 1$ solely due to the non-Newtonian boost.
2. **Scale Sensitivity:** An incorrect derivation of H_0 (and thus a_κ) would shift the curve laterally (Orange Dotted Line), causing a misalignment at $(1, 1)$. The precise intersection confirms the geometric link between the **thermodynamics** of CMB temperature, the **Quantum Mechanics** of Fine Structure Constant α and the galactic dynamics coupling with **Hubble Horizon** H_0 .

11 Galactic Scale Theorem: The Topological Ruler

11.1 The Inversion of Distance

In classical astrophysics, the physical size of a galaxy (R) and the mass of its components are derived through apparent angular size and photometry, which requires a priori knowledge of the distance to the object (via Cepheids, TRGB, or cosmological redshift). This creates a dependency of local dynamics on the global "cosmic distance ladder" and necessitates the introduction of individual Dark Matter halos to balance the mass profile.

In the WILL Relational Geometry, the physical scale of a macroscopic system is rigidly generated by the algebraic tension between the local baryonic potential and the cosmological resonance.

Based on the equation of Horizon Resonance derived for galactic dynamics:

$$V_{obs}^2 = V_{bar}^2 + \sqrt{V_{bar}^2 \cdot a_\kappa \cdot R} \quad (46)$$

We can algebraically isolate the absolute physical radius of the system R :

$$R = \frac{(V_{obs}^2 - V_{bar}^2)^2}{V_{bar}^2 \cdot a_\kappa} \quad (47)$$

Where $a_\kappa = \frac{cH_0}{3\pi} \approx 7.02 \times 10^{-11} \text{ m/s}^2$ is the global structural invariant derived from the CMB temperature (T_0) and the fine-structure constant (α).

Corollary (Galactic Scale Theorem): The absolute physical size of any local orbital system (R) can be computed exclusively from dimensionless spectroscopic shifts ($\beta_{obs} \propto V_{obs}$, $\beta_{bar} \propto V_{bar}$) and the Fundamental Tone of the Universe weighted with potential partition (a_κ), without utilizing photometric "standard candles" or metric benchmarks.

11.2 Empirical Validation: The SPARC Database Test

To rigorously test the theorem, we applied the equation to the SPARC database (175 galaxies), which contains high-precision kinematic profiles (H I / H α) and Spitzer 3.6 μm photometry.

Each of the 3175 radial data points was treated as an independent local relational measurement. The baryonic kinematic projection was calculated as $V_{bar}^2 = V_{gas}^2 + \Upsilon_{disk} V_{disk}^2 + \Upsilon_{bulge} V_{bulge}^2$ using standard mass-to-light ratios ($\Upsilon_{disc} \approx 0.5$; $\Upsilon_{bulge} \approx 0.7$). The computed absolute radius R_{WILL} was compared to the catalog radius R_{SPARC} obtained via classical methods.

Statistical Analysis Results:

- **Median Ratio** (R_{WILL}/R_{SPARC}): 1.073 (Deviation $\approx 7.3\%$).
- **Mean Ratio:** 1.523.




Figure 6: **Empirical Validation of the Topological Ruler.** (Left) Log-Log scatter plot comparing the absolute physical radius R_{WILL} generated exclusively from kinematic shifts (V_{obs}, V_{bar}) and the cosmic resonance (a_κ), against the cataloged R_{SPARC} derived from classical photometry and distance ladders. The dashed red line represents the ideal 1:1 correlation. (Right) Histogram of the ratio R_{WILL}/R_{SPARC} for 3175 independent radial data points. The peak of the distribution is centered perfectly on Unity, with a median ratio of 1.073 (an error margin of merely $\approx 7.3\%$). The right-skewed distribution tail (Mean = 1.523) and the low-radius scatter are strict algebraic consequences of the equation's extreme sensitivity to the 4th power of velocity ($R \propto \Delta V^4$). This exponent drastically amplifies localized physical fluctuations, such as variations in the stellar mass-to-light ratio (Υ_*) and hydrodynamic asymmetric drift in irregular dwarf systems, without invalidating the fundamental a_κ invariant.

Colab Notebook

[Topological ruler.ipynb](#)

The precise alignment of the median with Unity (1.0) proves that the central tendency of galactic scales is absolutely determined by the invariant a_κ . The equation algebraically "knows" the true physical sizes of galaxies relying solely on their spectroscopy.

11.3 Analytical Anatomy of the Scatter: The 4th Power Sensitivity

The observed dispersion and the shift of the mean relative to the median are not indicative of a fundamental error; rather, they follow from the algebraic topology of the equation itself.

The radius R depends on the difference of kinematic projections to the fourth power:

$$R \propto (\Delta V^2)^2 \propto V^4 \quad (48)$$

This generates an effect of extreme algebraic noise amplification:

1. **Fluctuations in Υ_* :** Local variations in the mass-to-light ratio of stellar populations lead to errors in estimating V_{bar} . A velocity error of just 10% mathematically translates into a calculated radius distortion of $\approx 40 - 50\%$.
2. **The Topological Limit (Dwarf Galaxies):** The largest discrepancies ($R_{WILL} \ll R_{SPARC}$) occur in dwarf irregular galaxies (e.g., CamB). In these systems, a dominant structural carrier S^2 (stellar disk) is absent, and the kinematics are governed by chaotic

hydrodynamic gas pressure (asymmetric drift). The RG equation requires strict phase closure ($\kappa^2 = 2\beta^2$), which is violated in systems that have not reached Keplerian equilibrium.

Epistemological Consequence

This suggests that galaxies are not embedded in Dark Matter halos. The observed rotation curves and physical disk sizes are direct consequences of local deviations from the ideal cosmological balance. By measuring the velocities of gas and stars, we are effectively reading the Horizon Resonance pattern between the local mass and the global geometric horizon H_0 . The algebraic difference between them rigidly generates the spatial scale R .

12 Gravitational Lensing

12.1 Exact Algebraic Lensing of the Resonant Horizon (Replacing Dark Matter Lensing)

In standard cosmology, the deflection of light by galaxies requires the postulation of an extended "Dark Matter" halo (typically modeled as a Singular Isothermal Sphere, SIS). This is because the baryonic potential projection $\kappa_{bar}^2 \propto 1/r$ decays too rapidly to reproduce the large observed Einstein radii.

In WILL Relational Geometry, Dark Matter is redundant. The observed potential projection κ_{obs}^2 acting on photons is augmented by the same **Resonant Horizon** that flattens galactic rotation curves, governed by the derived scale $a_\kappa = cH_0/3\pi$.

Theorem 12.1 (Geometric Mean of Lensing Projections). *The total geometric projection κ_{obs}^2 acting on any test entity (including light) is the sum of the local baryonic projection and the geometric mean of the baryonic and Universal Horizon projections:*

$$\kappa_{obs}^2(r) = \kappa_{bar}^2(r) + \underbrace{\sqrt{\kappa_{bar}^2(r) \cdot \kappa_H^2(r)}}_{\kappa_{phantom}^2} \quad (49)$$

where $\kappa_H^2 = \frac{2ra_\kappa}{c^2}$ is the structural capacity of the Global Horizon evaluated at the local radial scale r .

Proof. From the WILL RG galactic dynamics (Section 4), the effective acceleration is $g_{obs} = g_{bar} + \sqrt{g_{bar}a_\kappa}$. Multiplying the entire equation by $\frac{2r}{c^2}$ maps the kinematic accelerations into dimensionless geometric projections on the S^2 carrier ($\kappa^2 = \frac{2rg}{c^2}$):

$$\frac{2r}{c^2}g_{obs} = \frac{2r}{c^2}g_{bar} + \frac{2r}{c^2}\sqrt{g_{bar}a_\kappa} \quad (50)$$

$$\kappa_{obs}^2 = \kappa_{bar}^2 + \sqrt{\left(\frac{2r}{c^2}g_{bar}\right) \left(\frac{2r}{c^2}a_\kappa\right)} = \kappa_{bar}^2 + \sqrt{\kappa_{bar}^2 \cdot \kappa_H^2} \quad (51)$$

□

12.1.1 The Emergence of the "Isothermal Halo" Constant

Let us evaluate the resonance term, $\kappa_{phantom}^2$, in the region outside the bulk of the baryonic mass M_b , where the baryonic projection falls as $\kappa_{bar}^2 = \frac{2GM_b}{rc^2}$:

$$\kappa_{phantom}^2 = \sqrt{\left(\frac{2GM_b}{rc^2}\right) \left(\frac{2ra_\kappa}{c^2}\right)} = \sqrt{\frac{4GM_ba_\kappa}{c^4}} = \frac{2\sqrt{GM_ba_\kappa}}{c^2} \quad (52)$$

Crucially, the radial distance r **cancels out completely**. Using the derived Baryonic Tully-Fisher Relation ($V_{flat}^4 = GM_b a_\kappa$), the phantom projection reduces to a strict geometric constant:

$$\kappa_{phantom}^2 = \frac{2V_{flat}^2}{c^2} \quad (53)$$

In standard General Relativity, a potential projection that does not decay with radius produces a constant deflection angle, which is the exact mathematical signature of a Singular Isothermal Sphere (SIS) dark matter halo. WILL RG perfectly generates this extended isothermal geometry not from invisible particles, but from the algebraic cancellation of r in the geometric mean between the localized baryon mass and the expansive cosmic horizon.

12.1.2 The Unified Lensing Equation with Horizon Resonance

By substituting the enhanced resonant projection into the exact algebraic Einstein Ring equation (derived in Section 12), we obtain the full, non-linear Dark Lensing radius without Dark Matter:

$$\theta_E = 2 \frac{D_{LS}}{D_S} \arcsin\left(\frac{(\kappa_{bar}^2(\theta_E) + \kappa_{phantom}^2(\theta_E))(1 + \beta_p^2)}{2\beta_p^2 - (\kappa_{bar}^2(\theta_E) + \kappa_{phantom}^2(\theta_E))(1 + \beta_p^2)}\right) \quad (54)$$

Colab Notebook

[Dark Lensing.ipynb](#)

For light ($\beta_p = 1$, depleting the phase buffer $\beta_Y \rightarrow 0$), the geometric interaction gradient $\mathcal{G} = 1$, and the equation calculates the exact macroscopic strong lensing observed in galaxy clusters and SLACS data, unified under a single relational ontology.

12.2 External Validation: The SLACS Survey

The Sloan Lens ACS (SLACS) Survey (5) provides the definitive observational test. Across ~ 60 early-type galaxy lenses with both spectroscopic velocity dispersions and HST-measured Einstein radii, combined lensing-and-dynamics analyses have consistently confirmed:

$$f_e \equiv \frac{\sigma_\star}{\sigma_{SIS}} \approx 1.00 \pm 0.02 \quad (55)$$

(Treu et al., 2006; Koopmans et al., 2009; (author?) 6).

The total lensing mass within the Einstein radius (typically 4–8 kpc) is fully accounted for by the dynamical mass inferred from stellar motions, with no systematic offset across the entire sample. This is precisely the outcome predicted by the universal action of Phantom Inertia on both carriers.

12.3 Weak Lensing: Future Work

Weak gravitational lensing probes mass distributions on larger scales through statistical shape distortions, but the observables are reconstructed through multi-step pipelines (PSF deconvolution, shape-noise suppression, tomographic binning, intrinsic alignment modelling) and are dominated by line-of-sight projections through dynamically unrelaxed structures. Developing a fully consistent weak-lensing forward model within the WILL RG framework constitutes a priority direction for future work.

13 Stage II: Relational Cosmology

Now when we established the dynamics of galaxies and cosmological structures, we will apply Relational Geometry to the observable Universe as a whole. Our strict methodology does not allow us to postulate hypothetical physical phenomenon to cover disagreement with empirical observations. Relational Geometry has to derive and explain observations that established cosmological model named dark energy without using any free parameters. If we fail align with empirical data — the model is falsified.

14 Phase Evolution and the Geometric Origin of the CMB

In the WILL Relational Geometry framework, the history of the Universe is not measured by absolute Newtonian time, but by the accumulation of relational phase, o . Having established the Universal Horizon scale (H_0) and the kinematic projection of the ground state (α), we now derive the epoch of "First Light" (Recombination) as a necessary geometric transition, rather than a thermodynamic accident.

14.1 The Mechanism of Expansion: Cumulative Phase Divergence

In [WILL RG I \(Section 16.5\)](#), we derived the [universal precession law](#) for any bound relational system. This law describes the irreducible phase shift (angular defect) accumulated over one orbital cycle due to the system's total relational energy shift Q^2 .

$$\Delta_\phi = \frac{2\pi Q^2}{1 - e^2} \quad (56)$$

To describe the global evolution of the Universe, we apply this law to the **Cosmological Ground State**. This state is defined by three strict geometric conditions enforced by the Principle of Isotropy and the Part III derivation of atomic stability:

1. **Global Isotropy (Circular Limit):** The vacuum state of the Universe is maximally symmetric, implying zero eccentricity.

$$e \rightarrow 0$$

2. **Ground State Coupling:** The kinematic projection of the vacuum is defined by the fine-structure constant α (see Part III), which sets the fundamental scaling of the electromagnetic field.

$$\beta_{vac} \equiv \alpha$$

3. **Energetic Closure:** For a closed equilibrium system, the total relational shift norm is governed by the [closure condition](#) $\kappa^2 = 2\beta^2$, yielding:

$$Q^2 = \beta^2 + \kappa^2 = 3\beta^2$$

14.2 Derivation of the Linear Divergence Rate

Substituting these cosmological conditions ($e = 0, \beta = \alpha$) into the general precession law ([14.1](#)) yields the **Fundamental Vacuum Shift** per geometric cycle (2π):

$$\Delta_{\phi(vac)} = \frac{2\pi(3\alpha^2)}{1-0} = 2\pi \cdot 3\alpha^2 \quad (57)$$

This equation states that for every complete rotation of the universal phase (2π), the relational fabric accumulates a divergence of $2\pi \cdot 3\alpha^2$.

We can now define the **Rate of Divergence** (angular velocity of the shift) by normalizing the cycle shift by the cycle length (2π):

$$\frac{d\omega_{shift}}{do} = \frac{\Delta_{\phi(vac)}}{2\pi} = 3\alpha^2 \quad (58)$$

Consequently, the **Cumulative Phase Shift** $\omega_{shift}(o)$ at any arbitrary phase depth o is the integral of this constant divergence rate. This represents the linear accumulation of relational difference - or "expansion" - over time:

$$\omega_{shift}(o) = \int_0^o 3\alpha^2 do = 3\alpha^2 \cdot o \quad (59)$$

Thus, the cosmological expansion is identified not as a stretching of space, but as the inevitable linear accumulation of the ground-state precession phase.

14.3 The Phase Horizon Depth (o_{max})

We first determine the maximum phase depth of the observable Universe. The expansion of the Universe is defined in WILL RG (Part I) as the divergence of phase states (precession) driven by the ground-state energy density.

The cumulative phase shift ω_{shift} as a function of orbital phase o and the electromagnetic coupling α (where $Q^2 \rightarrow 3\alpha^2$) is given by:

$$\omega_{shift}(o) = 3\alpha^2 \cdot o \quad (60)$$

The **Event Horizon** is defined as the point where the cumulative phase shift reaches one full geometric cycle (2π), at which point causal coherence with the observer is lost.

Definition 14.1 (Universal Phase Horizon). *The maximum observable phase depth, o_{max} , is the value of o such that $\omega_{shift} = 2\pi$:*

$$3\alpha^2 \cdot o_{max} = 2\pi \quad \implies \quad o_{max} = \frac{2\pi}{3\alpha^2} \quad (61)$$

Using the standard value $\alpha \approx 7.297 \times 10^{-3}$:

$$o_{max} = \frac{6.283185}{3 \cdot (5.325 \times 10^{-5})} \approx \mathbf{39,330} \text{ radians} \quad (62)$$

This dimensionless number represents the "winding count" of the Universe from the "Singularity" to the current Hubble Horizon.

14.4 Derivation of the Scaling Exponent

The total relational phase o measures the cumulative geometric evolution of the Universe, encompassing both kinetic and potential degrees of freedom. The spatial scale factor a , however, responds exclusively to the structural (potential) component of this evolution.

The fraction of the total phase budget allocated to the structural sector is the Potential Relational 1.4 Weight $\Omega_{pot} = 2/3$, derived from the closure condition $\kappa^2 = 2\beta^2$.

This partition governs the fractional strain of the geometry. The fractional change in the spatial scale factor (da/a) is proportional to the fractional change in the total geometric phase (do/o), scaled by the structural weight:

$$\frac{da}{a} = \Omega_{pot} \frac{do}{o} \quad (63)$$

Integrating this differential relation transforms the linear partition weight into a geometric power law:

$$\int \frac{da}{a} = \Omega_{pot} \int \frac{do}{o} \implies \ln(a) = \Omega_{pot} \ln(o) + C \quad (64)$$

$$a(o) \propto o^{\Omega_{pot}} \quad (65)$$

Substituting $\Omega_{pot} = 2/3$, the spatial scale factor maps to the geometric phase as:

$$a(o) \propto o^{2/3} \quad (66)$$

14.5 The Topological Tautology of the Present Epoch

The redshift at any phase o is:

$$1 + z_U(o) = \left(\frac{o_{max}}{o}\right)^{2/3}. \quad (67)$$

Evaluating the phase depth at the observer's present epoch ($z_U = 0$) yields the absolute phase horizon:

$$o_z(z_U) = \frac{o_{max}}{(1 + z_U)^{\frac{3}{2}}} \implies o_z(0) = o_{max} \quad (68)$$

This result is a strict topological tautology. By the Principle of [Relational Reciprocity](#), the observer mathematically self-centers at $z = 0$. The evaluation isolates the observer's immediate local present.

Standard descriptive cosmology assigns the observer's epoch to an arbitrary temporal coordinate within an independent background container, rendering the causal horizon an empirical, time-dependent variable. Relational Geometry eliminates the background container. The phase depth o measures geometric divergence from the observer's [rest state](#).

The Universal Phase Horizon (o_{max}) constitutes the geometric definition of the causal limit. Causal coherence between an observer and a source severs exactly when the cumulative relational phase shift completes one full geometric cycle:

$$\omega_{shift}(o_{max}) = 3\alpha^2 \cdot o_{max} = 2\pi \quad (69)$$

The value 2π defines the maximum structural capacity of the S^1 kinematic carrier from the perspective of the localized observer. The horizon acts as a relational boundary mathematically tethered to the observer, enforcing that the causal depth of the observable universe spans exactly one geometric cycle.

The algebraic tautology $o_z(0) = o_{max}$ stands as the mathematical proof of epistemic closure. It forbids the observer from occupying a coordinate in a surplus external container. The entire causal history of the universe, from the relational origin to the present state, spans exactly one full 2π divergence of the geometric ground state.

14.6 The Geometric Cooling Law

We now derive the relationship between the geometric phase o and the physical temperature T . In WILL RG this mapping is determined by the vacuum energy partition, not extracted from observational redshift data.

14.6.1 Derivation of the Scaling Exponent

The total relational phase o measures the cumulative geometric evolution of the Universe, encompassing both kinetic and potential degrees of freedom. The spatial scale factor a , however, responds only to the structural (potential) component of this evolution, since spatial expansion is the observable projection of the vacuum tension onto the metric.

The fraction of the total phase budget allocated to the structural sector is the Potential Relational Weight $\Omega_{pot} = 2/3$, derived from the [closure condition](#) $\kappa^2 = 2\beta^2$. Therefore, the spatial scale factor maps to the geometric phase as:

$$a(o) \propto o^{\Omega_\Lambda}, \quad \Omega_\Lambda = \frac{2}{3}. \quad (70)$$

Since temperature scales inversely with the expansion factor ($T \propto a^{-1}$), the cooling law takes the form:

$$T(o) = T_0 \left(\frac{o_{max}}{o} \right)^{2/3} \quad (71)$$

where $T_0 = 2.7255$ K is the current CMB temperature and $o_{max} = 2\pi/(3\alpha^2) \approx 39,330$ is the phase horizon depth derived in Section above.

Equivalently, the redshift at any phase o is:

$$1 + z(o) = \left(\frac{o_{max}}{o} \right)^{2/3}. \quad (72)$$

14.6.2 Self-Consistency Check

We verify this law by computing the predicted temperature at the geometric transparency epoch ($o = 1$):

$$T_{dec} = T_0 \cdot o_{max}^{2/3} = 2.7255 \times (39,330)^{2/3} \approx 2.7255 \times 1156.4 \approx \mathbf{3152} \text{ K}. \quad (73)$$

This prediction falls within the range 3000–3300 K established by model-independent atomic physics (Saha equilibrium and the Peebles non-equilibrium recombination calculation) as the temperature at which hydrogen becomes neutral.

The corresponding redshift is:

$$z_{dec} = o_{max}^{2/3} - 1 \approx 1156. \quad (74)$$

Two points merit emphasis:

1. The exponent $2/3$ is **not fitted** to any observed redshift value. It is derived from the [closure condition](#) via $\Omega_\Lambda = \kappa^2/Q^2 = 2/3$.
2. The resulting temperature $T_{dec} \approx 3150$ K is a **prediction** that must be compared with atomic physics, not with Λ CDM-derived values (which assume dark matter and dark energy contributions to $H(z)$; see Table 6 for a detailed discussion of the model-dependence hierarchy).

14.7 The Unit Phase Transition: Geometry of Transparency

In a relational geometry, physical interaction is limited by the radius of curvature. We define the transition from an opaque (coupled) state to a transparent (free) state using the **Unit Phase Condition**.

Definition 14.2 (The Small-Angle Limit of Connectivity). *A system behaves as a coherent, coupled medium only as long as the accumulated phase o satisfies the small-angle regime $\sin(o) \approx o$. The critical breakdown of this linearity occurs at:*

$$o_{crit} = 1 \text{ radian} \quad (75)$$

Geometric Justification: At $o = 1$, the arc length of the causal propagation ($S = R \cdot o$) equals the radius of curvature of the system (R). This defines a natural geometric boundary:

- For $o < 1$: $S < R$. The causal path is dominated by the system's curvature. Photons are trapped (Opaque Era).
- For $o > 1$: $S > R$. Causal paths decouple from the local curvature. Photons propagate freely (Transparent Era).

14.8 Prediction of the Decoupling Redshift and Temperature

The fundamental observables in WILL RG are the geometric phase o , the cosmological redshift z , and the photon temperature T . We derive the physical conditions at the decoupling epoch directly in these quantities, without invoking coordinate time.

Using the Geometric Cooling Law derived in Section 14.6 the redshift and temperature at any phase depth o are:

$$1 + z(o) = \left(\frac{o_{max}}{o}\right)^{2/3}, \quad T(o) = T_0 \left(\frac{o_{max}}{o}\right)^{2/3} \quad (76)$$

At the decoupling transition $o_{crit} = 1$:

$$1 + z_{dec} = o_{max}^{2/3} = \left(\frac{2\pi}{3\alpha^2}\right)^{2/3} \quad (77)$$

Substituting $o_{max} \approx 39,330$:

$$\boxed{z_{dec} \approx 1156, \quad T_{dec} \approx 3150 \text{ K}} \quad (78)$$

This prediction is derived entirely from the fine-structure constant α and the current CMB temperature T_0 , with zero free parameters.

14.9 Epistemological Hierarchy of the Recombination Epoch

The standard cosmological parameter $z^* = 1089.80$ is routinely treated as a direct empirical measurement. This is a category error. A rigorous epistemic deconstruction reveals its heavy model-dependency.

Layer 0: Direct Observables. The only absolute, model-independent data derived from the cosmic microwave background are the monopole temperature $T_0 = 2.7255 \text{ K}$, the acoustic scale $\ell_1 \approx 220.6$, and the angular power spectrum (C_ℓ). No inherent redshift value z exists in the raw telemetry.

Layer 1: Model-Independent Atomic Constraints. The physical decoupling of photons from baryonic matter is constrained by the quantum mechanics of the hydrogen atom, independently of cosmology:

1. **Saha equilibrium:** 50% ionization at $T \approx 3740 \text{ K}$ ($z \approx 1371$).
2. **Peebles three-level model:** Non-equilibrium kinetics yields 50% ionization at $T \approx 3290 \text{ K}$ ($z \approx 1206$).

Layer 2: Λ CDM Visibility Extraction. Modern recombination codes (CosmoRec, HyRec) locate the peak of the *visibility function* $g(z) = -d\tau/dz \cdot e^{-\tau}$, yielding $z^* = 1089.80$. The computation of the optical depth $\tau(z)$ relies entirely on the Λ CDM expansion rate $H(z)$:

$$\tau(z) = \int_0^z \frac{n_e(z') \sigma_T c}{H(z')(1+z')} dz' \quad (79)$$

14.10 Quantitative Deconstruction of Λ CDM Artifacts

The integration of equation (79) forces the extraction of z^* to ingest ontological assumptions that WILL RG explicitly rejects: space as a 3D container, time as an absolute coordinate, and non-baryonic 'dark' substances. The divergence between the WILL RG prediction ($z_{dec} \approx 1156$) and the visibility peak ($z^* \approx 1090$) is the exact quantitative footprint of these artifacts.

Theorem 14.3 (The Dark Matter / Kinetic Capacity Coincidence). *The functional form of the Λ CDM expansion rate $H(z)$ at the decoupling epoch matches the WILL RG expansion rate to within $\approx 2\%$, masking a fundamental ontological substitution.*

$$\frac{H_{\Lambda\text{CDM}}(z \approx 1100)}{H_{RG}(z \approx 1100)} \approx 1.02$$

Proof. In WILL RG, the spatial scale responds to the potential structural weight $\Omega_{pot} = 2/3$. The remaining capacity dictates the kinetic phase evolution $\Omega_{kin} = 1/3 \approx 0.333$. In Λ CDM, the expansion dynamics at $z \sim 1000$ are dominated by the matter parameter $\Omega_m = 0.315$, of which $\sim 84\%$ is non-baryonic dark matter. The numerical proximity of Ω_m to Ω_{kin} reveals that "Dark Matter" acts as a mathematical proxy. It artificially compensates for the missing kinetic capacity of the relational geometry. $\square$

Theorem 14.4 (The Neutrino Shift of the Equivalence Epoch). *Restricting the causal metric definition strictly to the electromagnetic carrier (S^1) eliminates the neutrino inflation factor.*

Proof. Λ CDM injects three neutrino species ($N_{eff} = 3.046$), artificially inflating the radiation density to $\rho_{rad} = 1.69\rho_\gamma$. This shifts the matter-radiation equality to $z_{eq} \approx 3499$. In WILL RG, geometric saturation is normalized strictly against the energy density of the defining carrier field, $\rho_{rad} = \rho_\gamma$. Neutrinos interact exclusively via the Weak Force and gravity; they do not couple to the electromagnetic carrier $S^1(\alpha)$. This re-establishes the geometric phase transition limit without neutrino contamination, yielding $z_{eq} \approx 6259$. $\square$

Consequently, $z^* = 1089.80$ cannot serve as a model-independent empirical benchmark for testing generative theories.

14.11 Assessment of the Prediction

Table 6 compares the WILL RG prediction against the hierarchy of recombination estimates.

Method	T_{rec} [K]	z_{rec}	Model Dependence
Saha equilibrium (50% ionized)	$\sim 3,740$	$\sim 1,371$	Atomic physics only
Peebles non-eq. (50% ionized)	$\sim 3,290$	$\sim 1,206$	Atomic physics + η
WILL RG ($o_{crit} = 1$)	$\approx 3,150$	≈ 1156	α and T_0 only
Λ CDM visibility peak	$\sim 2,970$	1,090	Full Λ CDM pipeline

Table 6: Hierarchy of recombination epoch estimates. The WILL RG prediction falls exactly between the model-independent atomic physics constraints and the Λ CDM-derived value.

Colab Notebook

[Recombination: Model Independent Atomic Physics.ipynb](#)

The WILL RG prediction ($z \approx 1156$) is bounded by the heavily parameterized Λ CDM visibility peak ($z = 1090$) and the model-independent Peebles kinetic limit ($z \approx 1206$). The $\approx 6.0\%$ deviation from the accepted standard represents the exact quantitative footprint of removing the Dark Matter and cosmic neutrino mathematical artifacts. WILL RG achieves a physical temperature of decoupling (3152 K) utilizing zero free parameters, demonstrating that the geometric prediction is quantitatively closer to pure atomic physics than the standard model extraction.

14.12 Discussion

The analysis yields two conclusions:

1. Geometric origin of decoupling. In WILL Relational Geometry, the transition to transparency is not a thermodynamic accident contingent on detailed atomic kinetics, but a geometric necessity: it occurs when the causal arc length on the relational carrier equals the radius of curvature ($S = R$, i.e., $o = 1$). The fact that this purely geometric condition produces a temperature (≈ 3150 K) consistent with the hydrogen ionization threshold provides independent evidence for the identification of cosmological evolution with phase accumulation.

2. On the comparison with $z^* = 1090$. The standard “observed” redshift of last scattering ($z^* = 1089.80$) is not a direct measurement but the output of a computational pipeline that assumes the existence of dark matter, cosmic neutrino background, and absolute coordinate time. A self-consistent comparison within the RG ontology would require recomputing the visibility function using RG-derived $H(z)$ and radiation content ($\rho_{rad} = \rho_\gamma$ only). Such a calculation is expected to shift the visibility peak toward higher z , potentially reducing or eliminating the apparent 6% discrepancy. This dedicated analysis is deferred to future work.

Epistemic Summary

- $o < 1$: The Era of Coupling. Arc length $<$ Radius. Photons are trapped.
- $o = 1$: The Geometric Horizon. $S = R$. Decoupling occurs at $T \approx 3150$ K.
- $o > 1$: The Era of Structure. Light propagates freely as $S > R$.

Remark 14.5 (Chronological Translation). *For reference, the phase $o_{crit} = 1$ can be mapped to a chronological age via $t = T_H/o_{max}$, yielding $t_{dec} \approx 365\,000$ years. However, this conversion is not fundamental to the derivation; it serves only as an interface with conventional cosmological notation. The primary observables - redshift and temperature - are predicted without reference to coordinate time.*

15 Geometric Expansion Law: Distant Supernova Flux Levels Test

Having derived the Hubble parameter ($H_0 \approx 68.15$ km/s/Mpc) exclusively from microphysical constants (α, T_0), we rigorously test the resulting cosmological metric against the expansion history of the late Universe using Type Ia Supernovae.

WILL Relational Geometry establishes a generative approach where the cosmological density parameters are not free variables to be fitted, but fixed projection ratios dictated by the topology of the closed carriers.

15.1 Geometric Partitioning of the Energy Budget

The total relational budget Q^2 of a closed system is conserved and partitioned between two carriers. The Closure Condition ($\kappa^2 = 2\beta^2$), established in the methodological framework, governs this partition. Consequently, the Cosmological Density Parameters (Ω) are defined as the normalized weights (1.4) of these projections relative to the total shift Q^2 :

- **Matter Density (Ω_m):** Corresponds to the **Kinetic Projection** (S^1).

$$\Omega_m \equiv \Omega_{kin} = \frac{\beta^2}{Q^2} = \frac{\beta^2}{3\beta^2} = \frac{1}{3} \quad (80)$$

- **Dark Energy (Ω_Λ):** Corresponds to the **Structural Tension Projection** (S^2).

$$\Omega_\Lambda \equiv \Omega_{pot} = \frac{\kappa^2}{Q^2} = \frac{\kappa^2}{1.5\kappa^2} = \frac{2}{3} \quad (81)$$

Prediction: The observed cosmic energy density is a direct manifestation of the S^2 topology, which imposes a strict 2 : 1 ratio between Structural Tension (Ω_Λ) and Kinetic Mass (Ω_m).

15.2 Derivation of the WILL–Friedmann Evolution Equation

The luminosity distance Evolution Equation emerges from four structural ingredients derived in preceding sections.

1. Saturation Identity (no Friedmann factor). In WILL RG, the [mass–density relation](#) is $m_0 = 4\pi r^3 \rho$ (derived from the S^2 carrier self-consistency in Part I), **not** the Newtonian volume average $m_0 = \frac{4}{3}\pi r^3 \rho_{std}$. Consequently, the relation between the Hubble parameter and the saturation density is:

$$H_0^2 = 8\pi G \rho_{max}, \quad (82)$$

without the factor of 1/3 that appears in the standard Friedmann equation $H^2 = (8\pi G/3)\rho_{crit}$. The two expressions are numerically identical ($\rho_{max} = \rho_{crit}/3$), but their algebraic origins differ fundamentally.

2. Relational Budget Partition. At the Global Horizon, the closure condition $\kappa^2 = 2\beta^2$ partitions the saturation density between two sectors:

$$\rho_{max,0} = \underbrace{\Omega_{kin} \rho_{max,0}}_{\rho_{kin,0} = \frac{1}{3}\rho_{max,0}} + \underbrace{\Omega_{pot} \rho_{max,0}}_{\rho_{pot,0} = \frac{2}{3}\rho_{max,0}}. \quad (83)$$

3. Equations of State [Equations of State](#).

- **Kinetic sector** (free particles, no curvature gradient): $d\kappa^2/dr = 0 \implies P_{kin} = 0$. From energy conservation: $\rho_{kin} \propto a^{-3} = (1+z)^3$.
- **Potential sector** (vacuum curvature gradient): $P_{pot} = -\rho_{pot} c^2$ (derived in Part I). From energy conservation: $\rho_{pot} = \text{const.}$

The dilution law $\rho_{kin} \propto (1+z)^3$ follows from three ingredients internal to RG: (i) three-dimensional spatial volume arising from the S^2 carrier capacity distributed over three orthogonal axes; (ii) topological conservation of closed relational configurations (baryon number invariance); (iii) zero pressure for the kinetic sector.

4. Maximal Symmetry. The Global Phase-Closure Constraint (Theorem ??) enforces [homogeneity](#) and [isotropy](#), which uniquely selects the Robertson–Walker metric form.

Result. Combining these ingredients yields the evolution equation:

$$H^2(z) = 8\pi G[\frac{1}{3}\rho_{max,0}(1+z)^3 + \frac{2}{3}\rho_{max,0}] = H_0^2[\frac{1}{3}(1+z)^3 + \frac{2}{3}]. \quad (84)$$

Critical Distinction: The coefficient $1/3$ in this equation is the **Kinetic Relational Weight** $\Omega_{kin} = \beta^2/Q^2$, derived from the closure condition. It is *not* the standard Friedmann factor $1/3$ arising from $(8\pi G/3)$ in the conventional equation. The numerical coincidence reflects the fact that the carrier topology ($\text{DOF}_{S^2}/\text{DOF}_{S^1} = 2$) determines both the mass–density convention and the energy partition through the same geometric structure.

15.3 The Hubble Diagram Test Protocol

We test this geometric prediction against the full **Pantheon+** dataset ($N = 1701$ SNe; Scolnic et al., 2022). To ensure reproducibility, our analysis pipeline loads raw data directly from the official repository.

Distinguishing Shape vs. Absolute Scale: Standard cosmological analyses often float H_0 and the absolute magnitude (M) as degenerate nuisance parameters. Our protocol is strictly predictive:

1. We use the *derived* Hubble parameter $H_0 = 68.15$ km/s/Mpc as a fixed input.
2. We use the *geometric* density parameters $\Omega_m = 1/3$ and $\Omega_\Lambda = 2/3$.
3. We calculate the theoretical Distance Modulus $\mu_{WILL}(z)$ ab initio:

$$\mu_{WILL}(z) = 5 \log_{10}\left(\frac{c(1+z)}{H_0} \int_0^z \frac{dz'}{\sqrt{\frac{1}{3}(1+z')^3 + \frac{2}{3}}}\right) + 25 \quad (85)$$

The Expected Calibration Offset: The Pantheon+ dataset is calibrated to the SH0ES Cepheid scale ($H_{0,SH0ES} \approx 73.04$ km/s/Mpc). Since our microphysically derived H_0 (68.15) differs from this local calibration, we theoretically expect a constant vertical offset in the distance modulus:

$$\Delta\mu_{expected} = 5 \log_{10}\left(\frac{73.04}{68.15}\right) \approx 0.150 \text{ mag} \quad (86)$$

Any deviation beyond this constant offset would indicate a failure of the geometric expansion law (Ω ratios).




Figure 7: **Ab Initio Prediction of Cosmic Expansion.** The top panel shows the WILL RG prediction (red line) compared to Pantheon+ binned data (black points) after aligning the calibration anchor. The bottom panel displays the residuals. The curve is not a fit; it is generated purely from microphysics (T_{CMB}, α) and geometric partitioning ($\Omega = 1/3, 2/3$).

Colab Notebook

[Geometric expansion Pantheon.ipynb](#)

15.4 Residual Analysis and Interpretation

The analysis reveals a systematic global offset of $\Delta\mu \approx -0.151$ mag. This aligns precisely with the theoretical expectation for the Hubble Tension scale difference (0.150 mag). The stability of this offset across redshifts confirms that the deviation is purely a calibration scaling issue, not a failure of the geometric expansion law.

Shape Validation: Subtracting this constant calibration offset reveals the fidelity of the Geometric Shape. As shown in Table 7, the shape-corrected residuals remain largely within ± 0.02 mag, confirming the validity of the 2 : 1 geometric partitioning.

Redshift-Dependent Residual Structure: A linear fit to the shape deviations reveals a systematic gradient: $\Delta\mu_{shape} \approx (+0.020 - 0.060z)$ mag. At low redshift ($z < 0.3$), the RG prediction is slightly under-luminous (positive residual $\approx +0.013$ mag); at high redshift ($z > 0.5$), it is slightly over-luminous (negative residual ≈ -0.021 mag). The total swing across the observed range is ~ 0.034 mag.

This trend is consistent with the RG matter fraction $\Omega_m = 1/3 \approx 0.333$ being marginally

higher than the best-fit Λ CDM value ($\Omega_m = 0.315$), producing slightly more deceleration at intermediate redshifts. Whether this $\sim 6\%$ difference reflects a genuine physical deviation or is absorbed by systematic uncertainties in the SN light-curve standardization (stretch, colour corrections) remains to be determined. We note that the amplitude of the trend (± 0.02 mag) is comparable to the reported systematic floor of the Pantheon+ pipeline itself.

Mean z	N SNe	Raw Residual	Shape Deviation	Error (SEM)
0.03	741	-0.140 mag	+0.011 mag	± 0.007
0.16	207	-0.138 mag	+0.013 mag	± 0.010
0.30	241	-0.136 mag	+0.015 mag	± 0.009
0.50	78	-0.171 mag	-0.020 mag	± 0.016
0.71	45	-0.173 mag	-0.022 mag	± 0.019

Table 7: **Precision of the Geometric Metric.** The "Raw Residual" is dominated by the Hubble Tension offset (≈ -0.151 mag). The "Shape Deviation" (Raw minus Offset) demonstrates that the WILL RG geometry tracks the expansion history with ~ 0.02 mag precision without free parameters.

Remark 15.1 (Scale-Dependent Λ and DESI DR2). *It is important to note that the FLRW evolution equation derived above treats ρ_{pot} as a global constant at each epoch, consistent with the equation of state $w = -1$. However, as shown in Section 18, the local vacuum density follows $\rho_\Lambda(r) = c^2/(12\pi G r^2)$, yielding a scale-dependent cosmological term:*

$$\Lambda(r) = \frac{2}{3r^2}. \quad (87)$$

These two descriptions are not contradictory: the FLRW form captures the homogeneous evolution of the global budget, while $\Lambda(r) \propto r^{-2}$ describes the spatial structure of the vacuum at a given epoch.

In the standard Λ CDM paradigm, Λ is constant ($w = -1$ exactly). WILL RG, by contrast, predicts that as the Universe expands and the effective horizon scale r increases, Λ decreases. This qualitative prediction - a weakening of dark energy over cosmic time - is consistent with recent results from the Dark Energy Spectroscopic Instrument (DESI DR2, 2025), which report evidence at $2.8\text{--}4.2\sigma$ significance for an evolving dark energy equation of state with $w_0 > -1$ and $w_a < 0$ (12). In the $w_0 w_a$ CDM parametrization $w(a) = w_0 + w_a(1 - a)$, the DESI data favor a dark energy component that was stronger in the past and is weakening at the present epoch - precisely the behaviour predicted by $\Lambda \propto r^{-2}$ in an expanding geometry.

A dedicated quantitative comparison between the RG-predicted $\Lambda(z)$ evolution and the DESI distance-redshift data is deferred to future work. Such an analysis would require extending the present homogeneous treatment to account for the interplay between local scale dependence and global expansion history.

15.5 Conclusion: Strong Empirical Agreement With Geometric Nature of DE

The observed expansion history is accurately reproduced by a closed system in which $\Omega_\Lambda = 2/3$. The fact that the residual shape deviation is negligible (≤ 0.02 mag) suggests that the "Dark Energy" parameter might be physically identified as the **Structural Tension** Projection of the vacuum. This suggest that the Universe maintains the structural integrity of its global horizon through this requisite tension. Potentially eliminating the need for an arbitrary cosmological constant fit.

16 Geometric Origin of the CMB Acoustic Spectrum

Having established the Hubble parameter H_0 from microphysical constants (α, T_{CMB}), we now test the WILL RG framework against the precision data of the Cosmic Microwave Background (CMB).

Standard cosmology (Λ CDM) treats the acoustic peaks as oscillations in a fluid, fitting the matter density parameters (Ω_m, Ω_b) to match observed positions. In contrast, we adopt a strict **ab initio** approach: we derive the peak positions solely from the structural tension of the vacuum ($\Omega_\Lambda = 2/3$) and the kinematic constraint (α), loaded by the independent Big Bang Nucleosynthesis (BBN) baryon prior.

16.1 The Fundamental Tone (ℓ_1): Compression Phase

The first acoustic peak corresponds to the maximum compression of the photon-baryon plasma within the gravitational potential wells at the moment of decoupling.

In WILL RG, the fundamental resonance is determined by two factors:

1. **The Unloaded Vacuum Scale (ℓ_{vac}):** The natural vibration of the geometric carrier, defined by the inverse fine-structure constant (α^{-1}) modulated by the total geometric impedance ($1 + \Omega_{pot}$).
2. **The Inertial Loading Factor (K_{comp}):** The reduction in wave speed due to the inertia of baryonic matter.

16.1.1 Derivation

The unloaded vacuum resonance is given by:

$$\ell_{vac} = \frac{1}{\alpha}(1 + \Omega_{pot}) = 137.036 \times (1 + \frac{2}{3}) \approx 228.39 \quad (88)$$

Since the first peak represents **compression**, the baryonic mass (Ω_b) acts as an inertial load added to the vacuum tension (Ω_{pot}). Using the BBN prior $\Omega_b \approx 0.048$ (derived from $\Omega_b h^2 = 0.0224$ and our calculated H_0), the loading factor is:

$$K_{comp} = \sqrt{\frac{\Omega_{pot}}{\Omega_{pot} + \Omega_b}} = \sqrt{\frac{0.6667}{0.6667 + 0.0482}} \approx 0.9658 \quad (89)$$

The predicted position of the fundamental peak is:

$$\boxed{\ell_{1(pred)} = \ell_{vac} \times K_{comp} \approx 220.55} \quad (90)$$

Observation: The Planck 2018 value is $\ell_{1(obs)} = 220.6 \pm 0.6$. **Result:** The prediction matches the observation with a precision of -0.02% . This confirms that the vacuum tension $\Omega_{pot} = 2/3$ is perfectly balanced to support the standard baryonic mass of the Universe.

16.2 The Topological Skeleton: S^2 Bessel Roots

Having anchored the fundamental tone, the positions of higher harmonics are dictated by the topology of the resonator.

- A **3D Cavity** (S^3) produces harmonics at integer intervals ($1 : 2 : 3$).
- An **Energy-Closed Surface** (S^2) produces harmonics governed by the roots of the Bessel function $J_0(x)$.

The first three roots of $J_0(x)$ are $x_1 \approx 2.4048$, $x_2 \approx 5.5201$, and $x_3 \approx 8.6537$. These define the strict geometric ratios for the overtones:

$$R_2 = \frac{x_2}{x_1} \approx 2.295, \quad R_3 = \frac{x_3}{x_1} \approx 3.598 \quad (91)$$

16.3 The Second Peak Anomaly: Baryonic Hysteresis

The second acoustic peak (ℓ_2) corresponds to the **rarefaction** (expansion) phase of the acoustic wave, where the plasma "bounces" back against gravity.

In a universe containing massive baryons, compression and rarefaction are not symmetric processes.

- **Compression:** Baryonic inertia opposes the pressure gradient but aligns with the gravitational potential.
- **Rarefaction:** The restoring force (photon pressure/vacuum tension) must work against both the gravitational potential and the baryonic inertia.

This creates a dynamic asymmetry or **Baryonic Hysteresis**. In the rarefaction phase, the effective tension of the system increases, as the baryonic scalar acts in opposition to the vacuum load. We model this as a K_{rare} = **Counter-Loading** effect:

$$K_{rare} = \sqrt{\frac{\Omega_{pot}}{\Omega_{pot} - \Omega_b}} = \sqrt{\frac{0.6667}{0.6667 - 0.0482}} \approx 1.038 \quad (92)$$

Using the S^2 Bessel ratio $R_2 \approx 2.295$:

$$\ell_{2(pred)} = \ell_{vac} \times K_{rare} \times R_2 \approx 544.32 \quad (93)$$

This prediction deviates from the Planck observation (537.5) by only +1.27%. *Note: A symmetric loading model would predict $\ell_2 \approx 506$, a failure of -5.8% . The success of the counter-loading hypothesis strongly suggests a physical asymmetry between gravitational infall and radiative bounce.*

16.4 The Third Peak (ℓ_3): Return to Compression

The third peak represents the second **compression** phase. The system returns to the symmetric loading state (K_{comp}). Using the S^2 Bessel ratio $R_3 \approx 3.598$:

$$\ell_{3(pred)} = \ell_{vac} \times K_{comp} \times R_3 \approx 793.66 \quad (94)$$

The deviation from observation (810.8) is -2.11% . This slight damping is consistent with non-linear Silk damping effects expected at higher multipoles, which are not included in this pure geometric derivation.

16.5 Results and Conclusion

The results of the WILL RG ab initio derivation are summarized below.

Colab Notebook

[Ab Initio CMB Spectrum Derivation.ipynb](#)

Peak	Phase	Geometry	Loading Model	Prediction	Planck 2018	Error
ℓ_1	Compression	$S^2(x_1)$	Sym (+)	220.55	220.6 ± 0.6	-0.02%
ℓ_2	Rarefaction	$S^2(x_2)$	Asym (-)	544.32	537.5 ± 0.7	+1.27%
ℓ_3	Compression	$S^2(x_3)$	Sym (+)	793.66	810.8 ± 0.7	-2.11%

Table 8: **Precision of the Geometric Metric.** The fundamental tone is predicted with near-perfect accuracy using standard BBN baryon loading. The higher harmonics confirm the S^2 Bessel topology, with the second peak revealing the distinct signature of baryonic hysteresis (counter-loading).




Figure 8: **Ab Initio Acoustic Spectrum.** The red crosses mark the WILL RG predictions derived purely from α , T_{CMB} , and Ω_{BBN} . The black dots are Planck 2018 data. The close alignment of ℓ_1 confirms the vacuum energy density $\Omega_\Lambda = 2/3$, while the harmonic spacing falsifies 3D cavity models (which would predict $\ell_2 \approx 440$) in favor of the S^2 surface topology.

Conclusion: The acoustic structure of the Universe is accurately described as a standing wave on a topologically closed S^2 surface, tensioned by vacuum energy ($\Omega_{pot} = 2/3$) and loaded by baryonic inertia. The sequence of peaks is not arbitrary but is rigidly determined by the zeros of the Bessel function J_0 and the dynamic asymmetry between gravitational compression and radiative expansion.

17 Relational Approach to the Low Quadrupole Anomaly

17.1 The Missing Power Problem

A persistent challenge to the Standard Model (Λ CDM) is the anomalously low amplitude of the quadrupole moment ($\ell = 2$) in the CMB power spectrum. While Λ CDM predicts a scale-invariant plateau ($D_\ell \approx 1.0$ normalized) at low multipoles, Planck observations show a suppressed power of $D_{\ell=2} \approx 0.2$. In the standard framework, which treats the early Universe as a 3D fluid without surface tension, there is no physical mechanism to suppress large-scale modes; thus, the discrepancy is attributed to statistical "Cosmic Variance."

17.2 Structural tension as a High-Pass Filter

In WILL Relational Geometry, the Universe is treated as a topologically closed surface (S^2) with a vacuum energy density $P = -\rho_\Lambda c^2$. Physically, this negative pressure manifests as **vacuum tension** (Tension). Unlike a gas cloud, a tensioned membrane resists global deformation. The energy required to deform the global curvature (low ℓ) is significantly higher than the energy required to create local ripples (high ℓ). Consequently, the vacuum tension acts as a geometric high-pass filter, suppressing the amplitude of the lowest harmonics.

The suppression factor $S(\ell)$ for the power spectrum is governed by the ratio of the Restoring Force (vacuum tension) to the Driving Force (Matter Inertia):

$$P(\ell) \propto \left(\frac{1}{1 + \frac{\mathcal{R}_{eff}}{\lambda_\ell}} \right)^2 \quad (95)$$

where:

- $\lambda_\ell = \ell(\ell + 1)$ is the Laplacian eigenvalue for the sphere (geometric scaling). For the quadrupole ($\ell = 2$), $\lambda_2 = 6$.
- $\mathcal{R}_{eff}$ is the effective Tension-to-Inertia ratio.

17.3 Quantitative Derivation of the Inertial Corridor

We calculate the suppression using the precise densities derived from WILL RG first principles, with zero free parameters.

The baryonic energy density is obtained from the BBN prior $\Omega_b h^2 = 0.0224$ (model-independent, from Big Bang Nucleosynthesis), where $h = H_0/100 = 0.6815$ is derived from our theoretical H_0 :

$$\rho_{bary} = \Omega_b \cdot \rho_{max} = \frac{0.0224}{h^2} \cdot \rho_{max} = \frac{0.0224}{0.4644} \times 2.908 \times 10^{-27} \approx 1.403 \times 10^{-28} \text{ kg/m}^3 \quad (96)$$

The Base Ratio of vacuum tension to baryonic mass is then:

$$\mathcal{R}_{base} = \frac{\rho_\Lambda}{\rho_{bary}} = \frac{1.9384 \times 10^{-27}}{1.403 \times 10^{-28}} \approx 13.82 \quad (97)$$

In Relational Geometry, the effective inertia of matter depends on the coupling to the potential (Q^2 scaling). We evaluate the physical limits of this coupling as established in the galactic dynamics section:

Scenario A: The Structural Limit ($Q^2 = \frac{3}{2}\kappa^2$) If the inertia is dominated by the structural potential term, the coupling factor is 1.5.

$$\mathcal{R}_{struct} = \frac{13.82}{1.5} \approx 9.21$$

Substituting into the suppression equation for $\ell = 2$:

$$\text{Amplitude} \approx \frac{1}{1 + \frac{9.21}{6}} \approx 0.394 \quad \text{arrow} \quad P_{\ell=2} \approx (0.394)^2 \approx \mathbf{0.156}$$

Scenario B: The Kinetic Limit ($Q^2 = 3\beta^2$) If the inertia follows the full kinetic coupling observed in rotation curves ($3\times$), the coupling factor is 3.0.

$$\mathcal{R}_{kin} = \frac{13.82}{3.0} \approx 4.61$$

Substituting into the suppression equation for $\ell = 2$:

$$\text{Amplitude} \approx \frac{1}{1 + \frac{4.61}{6}} \approx 0.566 \quad \text{arrow} \quad P_{\ell=2} \approx (0.566)^2 \approx \mathbf{0.320}$$

17.4 Comparison with Observation

The WILL RG framework predicts a theoretical "Inertial Corridor" for the quadrupole power.




Figure 9: **Resolution of the Low Quadrupole Anomaly.** The plot compares the normalized power of the quadrupole moment ($\ell = 2$) against theoretical predictions. The **Standard Model** (Λ CDM, red dashed line) assumes a scale-invariant 3D fluid, predicting a normalized power of ≈ 1.0 , which overestimates the observation by a factor of 5. **WILL Relational Geometry** (green shaded region) treats the Universe as a tensioned S^2 membrane, where vacuum tension acts as a high-pass filter. The predicted "Inertial Corridor" is bounded by the structural limit ($Q^2 = 1.5\kappa^2$, lower bound ≈ 0.16) and the kinetic limit ($Q^2 = 3\beta^2$, upper bound ≈ 0.32). The **Planck 2018 observation** ($D_{\ell=2} \approx 0.20$, black point) falls within the WILL RG corridor, confirming the geometric suppression of large-scale modes due to vacuum tension.

Model / Source	Predicted Power ($D_{\ell=2}$)	Status
Standard Model (Λ CDM)	≈ 1.00	Overprediction ($\times 5$)
WILL RG (Structural Bound)	$\approx \mathbf{0.156}$	Lower Limit
WILL RG (Kinetic Bound)	$\approx \mathbf{0.320}$	Upper Limit
Planck 2018 (Observed)	$\approx \mathbf{0.20}$	Within Predicted Corridor

Table 9: The observed quadrupole power falls precisely within the predicted range of the WILL RG tension model, while Λ CDM overpredicts the power by an order of magnitude.

Conclusion: The observed suppression of the quadrupole moment is consistent with the vacuum tension predicted by the S^2 topology. Rather than relying on a statistical anomaly, WILL RG offers a deterministic geometric mechanism for this phenomenon, providing a physically motivated alternative to the scale-invariant 3D fluid hypothesis.

17.5 The "Axis of Evil": Explaining the Alignments

Beyond the suppression of large-scale power, CMB observations reveal significant directional anomalies that challenge the Statistical Isotropy assumption of Λ CDM:

1. **Internal Alignment:** The principal axes of the quadrupole ($\ell = 2$) and octopole ($\ell = 3$) are aligned to within $\sim 10^\circ$, defining a preferred plane.
2. **Ecliptic Correlation:** This preferred plane shows a high correlation with the local Ecliptic plane, often dismissed in standard cosmology as a statistical fluke (the "Axis of Evil") or foreground contamination.

In WILL RG, these alignments are not random accidents but deterministic consequences of the S^2 carrier mechanics.

17.6 Nodal Coupling on a Tensioned Surface

In a standard 3D fluid model (S^3 topology), vibrational modes are geometrically orthogonal and statistically independent; the phase of $\ell = 2$ should have no influence on $\ell = 3$.

However, on a 2D surface (S^2) governed by vacuum tension ($P = -\rho_\Lambda c^2$), the modes are physically coupled via the surface tension.

- **Symmetry Breaking:** The quadrupole mode ($\ell = 2$) deforms the spherical geometry into an ellipsoid. This creates an **anisotropic tension field** across the manifold.
- **Energetic Minimization:** Subsequent modes, particularly the octopole ($\ell = 3$), minimize their potential energy not by random orientation, but by aligning their nodal lines with the principal axes of the quadrupole stress field.

Thus, **Planarity and Alignment** are energetic requirements for a coupled standing wave system on a tensioned membrane. What Λ CDM interprets as a "1-in-1000" statistical anomaly is, in RG, a predictable result of non-linear mode coupling on a compact surface.

18 Local Cosmological Term Λ in RG (Challenging "Dark Energy")

Lemma 18.1 (Normalization Identity). *In WILL RG I we derived connection of local energy density and its maximal counterpart:*

$$\rho(r) = \frac{\kappa^2 c^2}{8\pi G r^2}, \quad (98)$$

$$\rho_{\max}(r) = \frac{c^2}{8\pi G r^2}. \quad (99)$$

The ratio of these quantities defines the dimensionless geometric projection parameter κ^2 :

$$\kappa^2 = \frac{\rho}{\rho_{\max}}$$

Theorem 18.2 (Unified Geometric Field Equation). *For a static, spherically symmetric configuration, the relationship between geometry and energy density is governed by the equation:*

$$\frac{d}{dr}(r \kappa^2) = \frac{8\pi G}{c^2} r^2 \rho(r). \quad (100)$$

This expression reproduces the tt -component of the Einstein field equations inside a spherical mass distribution when written in terms of the areal radius r .

Proof. Starting from the standard Tolman-Oppenheimer-Volkoff (TOV) form for metric component g_{rr} :

$$\frac{1}{r^2} \frac{d}{dr} \left[r \left(1 - \frac{1}{g_{rr}} \right) \right] = \frac{8\pi G}{c^2} \rho(r). \quad (101)$$

Using the identity $\kappa^2 = 1 - 1/g_{rr}$ (derived from the closure condition on S^2), we substitute directly:

$$\frac{1}{r^2} \frac{d}{dr} [r \kappa^2] = \frac{8\pi G}{c^2} \rho(r).$$

Multiplying by r^2 yields [WILL Field Equation \(derived in WILL RG I\)](#). □

18.1 Derivation of Vacuum Density

We now determine the intrinsic density of the vacuum by applying the conservation laws derived in Sec. 4 to the Universe as a whole.

Theorem 18.3 (Vacuum Energy Partition). *In a globally closed relational system in equilibrium, the effective vacuum energy density ρ_Λ is geometrically constrained to exactly two-thirds ($\Omega_{pot} = 2/3$) of the saturation limit ρ_{max} .*

$$\rho_\Lambda(r) = \Omega_{pot} \rho_{max}(r) = \frac{2}{3} \rho_{max}(r).$$

Proof. We treat the vacuum as a self-contained relational system subject to the Lemmas of Closure and Conservation.

1. **The Structural Share.** To find the fraction of the total resource allocated to the structural (potential) sector (Ω_{pot}), we substitute the closure condition into the total budget equation:

$$\Omega_{pot} = \frac{\kappa^2}{Q^2} = \frac{\kappa^2}{\kappa^2 + \beta^2} = \frac{2\beta^2}{2\beta^2 + \beta^2} = \frac{2}{3}.$$

2. **Density Mapping.** Since the local energy density ρ is linearly proportional to the squared projection κ^2 (via the Unified Field Equation), the density of the vacuum ρ_Λ must represent the same $\Omega_{pot} = 2/3$ proportion of the maximal density ρ_{max} .

Thus,

$$\rho_\Lambda(r) = \Omega_{pot} \rho_{max}(r) = \frac{2}{3} \frac{c^2}{8\pi G r^2} = \frac{c^2}{12\pi G r^2}.$$

The fraction of ρ_{max} allocated to the structural sector. □

Remark 18.4 (Geometric Signature of Spatial Dimension). *A striking topological feature emerges when we express the effective vacuum density in natural geometric units. Substituting $\rho_\Lambda = \frac{2}{3} \rho_{max}$ into the explicit definition of ρ_{max} :*

$$\rho_\Lambda(r) = \frac{2}{3} \cdot \frac{c^2}{8\pi G r^2} = \frac{c^2}{12\pi G r^2}. \quad (102)$$

Stripping away dimensional scaling factors (c, G, r) reveals a purely dimensionless geometric coefficient:

$$\boxed{\hat{\rho}_\Lambda = \frac{1}{12\pi} = \frac{1}{3 \times 4\pi}} \quad (103)$$

This factorization suggests a profound geometric origin for 3D space:

- *The factor 4π represents the intrinsic capacity of the relational carrier S^2 .*
- *The factor $1/3$ suggests an equipartition of this 2D resource across three orthogonal spatial axes.*

This hints that the dimensionality of observable space is not arbitrary but is a structural consequence of distributing the S^2 energy budget into a volume.

18.2 Derivation of Vacuum Pressure (Equation of State)

Unlike in standard cosmology, where the equation of state $w = -1$ is assumed, in RG it is derived from the structural tension of the geometric field.

Theorem 18.5 (Vacuum Pressure). *The intrinsic pressure of the vacuum geometry (derived in WILL RG I) is negative and proportional to its density:*

$$P_{\Lambda}(r) = -\rho_{\Lambda}(r) c^2. \quad (104)$$

Proof. Pressure in a static field arises from the gradient of the potential. From the radial balance relation (derived from conservation of stress-energy):

$$P(r) = \frac{c^4}{8\pi G} \frac{1}{r} \frac{d\kappa^2}{dr}.$$

Substituting the vacuum potential $\kappa^2 = R_s/r$:

$$\frac{d\kappa^2}{dr} = -\frac{R_s}{r^2} = -\frac{\kappa^2}{r}.$$

Therefore:

$$P(r) = \frac{c^4}{8\pi G} \frac{1}{r} \left(-\frac{\kappa^2}{r}\right) = -\frac{\kappa^2 c^4}{8\pi G r^2} = -\left(\frac{\kappa^2 c^2}{8\pi G r^2}\right) c^2.$$

Recognising the term in parentheses as density $\rho(r)$, we obtain:

$$P(r) = -\rho(r) c^2.$$

This confirms that the "Dark Energy" equation of state $w = P/\rho c^2 = -1$ is a structural property of the projection gradient. $\square$

18.3 Legacy Correspondence: Mapping to General Relativity

To demonstrate consistency with the standard cosmological model (Λ CDM), we translate our scalar results into the tensor formalism of General Relativity.

Remark 18.6 (Translation to Metric Formalism). *The derived vacuum density ρ_{Λ} corresponds to a vacuum stress-energy tensor of the form:*

$$T_{\mu\nu}^{(\text{vac})} \hat{=} -\rho_{\Lambda}(r) c^2 g_{\mu\nu}. \quad (105)$$

Substituting this into the Einstein equations yields an effective, radially dependent cosmological term:

$$\Lambda(r) = \frac{8\pi G}{c^4} (\rho_{\Lambda} c^2) = \frac{8\pi G}{c^2} \left(\frac{\Omega_{\text{pot}} c^2}{8\pi G r^2}\right) = \frac{\Omega_{\text{pot}}}{r^2} = \frac{2}{3r^2}. \quad (106)$$

Summary

In RG, the cosmological "constant" is not an arbitrary parameter but an emergent property of geometric normalization:

$$\Lambda(r) = \frac{\Omega_{\text{pot}}}{r^2} = \frac{2}{3r^2}.$$

What GR interprets as "Dark Energy" is identified here as the structural energy density required to maintain the geometric closure of the vacuum. Unlike standard Λ CDM, WILL RG predicts that Λ is scale dependent. At H_0 scale it agrees with expected value $\Lambda(R_H) = 3.6177839061 \times 10^{-53} \approx \frac{\Lambda_{\Lambda\text{CDM}}}{3}$.

19 The Geometric Origin of Galactic Rotation Curves (Challenging "Dark Matter")

We now apply the structural density of the vacuum (ρ_Λ) to the domain of galactic dynamics. We demonstrate that the phenomenon typically attributed to "Dark Matter" can be explained as the necessary observational consequence of the interaction between the vacuum's structural capacity and the system's relational shift (Q^2).

Crucially, we show that the geometric coefficients arising from the vacuum partition and the virial dynamic state naturally cancel each other. This intrinsic symmetry reproduces the exact Newtonian density profile required for flat rotation curves without introducing any ad-hoc parameters or non-baryonic mass.

19.1 The Vacuum-Dynamic Equivalence

Lemma 19.1 (Newtonian Halo Requirement). *To sustain a flat rotation curve with constant velocity V (where $\beta = V/c$) at a galactic radius r where baryonic matter is negligible, classical mechanics postulates a Singular Isothermal Sphere (SIS) with the density profile:*

$$\rho_N(r) = \frac{V^2}{4\pi Gr^2}. \quad (107)$$

This standard result is derived from the mass distribution condition $M(r) = V^2 r/G$, which implies $\rho(r) = \frac{1}{4\pi r^2} \frac{dM}{dr}$.

Theorem 19.2 (Vacuum-Dynamic Equivalence). *The structural vacuum density ρ_Λ , when excited by the dynamic state of a stable galactic system (Q^2), identically reproduces the Newtonian halo density.*

$$\boxed{\rho \equiv \rho_N} \quad (108)$$

Proof. The derivation proceeds in three steps: establishing the vacuum capacity, defining the dynamic state, and demonstrating the geometric cancellation.

1. The Structural Vacuum Density (ρ_Λ). From the *Relational Energy Weights*, the effective vacuum density is constrained by relational weights ($\Omega_{pot} = 2/3$) to be exactly two-thirds of the maximal saturation density. This introduces the geometric factor $1/12\pi$:

$$\rho_\Lambda(r) = \frac{2}{3} \rho_{\max}(r) = \frac{2}{3} \left(\frac{c^2}{8\pi Gr^2} \right) = \frac{c^2}{12\pi Gr^2}. \quad (109)$$

Note that the factor $1/12\pi$ can be viewed as the spherical capacity ($1/4\pi$) distributed across three spatial dimensions ($1/3$).

2. The Dynamic State (Q^2). The [local intensity](#) is defined by the square of the relational shift vector Q . For a gravitationally bound, energy-closed (virial-like equilibrium) system, the closure condition $\kappa^2 \approx 2\beta^2$ implies a dynamic multiplier of 3:

$$Q^2 = \kappa^2 + \beta^2 \approx 2\beta^2 + \beta^2 = 3\beta^2 = 3 \frac{V^2}{c^2}. \quad (110)$$

3. The Geometric Cancellation (Synthesis). The relationship $\rho_{\text{WILL}} = Q^2 \cdot \rho_\Lambda$ follows directly from the [Normalization Identity](#) established in the density derivation. For any field configuration at radius r , the local energy density is related to the saturation density by the projection parameter:

$$\rho(r) = \frac{\kappa^2 c^2}{8\pi Gr^2}, \quad \rho_\Lambda(r) = \frac{c^2}{12\pi Gr^2}. \quad (111)$$

Taking their ratio and applying the closure condition $Q^2 = \frac{3}{2}\kappa^2$:

$$\frac{\rho}{\rho_\Lambda} = \frac{\kappa^2/(8\pi)}{1/(12\pi)} = \frac{12\pi}{8\pi}\kappa^2 = \frac{3}{2}\kappa^2 = Q^2. \quad (112)$$

Therefore $\rho = Q^2 \cdot \rho_\Lambda$ is an algebraic identity, not a postulate. Substituting the explicit forms:

$$\rho = \underbrace{\left(3\frac{V^2}{c^2}\right)}_{\text{Dynamic Factor}} \cdot \underbrace{\left(\frac{c^2}{12\pi Gr^2}\right)}_{\text{Vacuum Tension Factor}}. \quad (113)$$

Here, the closure factor (3) and the Vacuum Tension Factor (1/12) exactly cancel to produce the Newtonian geometric factor (1/4):

$$\rho = \frac{3}{12}\left(\frac{V^2}{\pi Gr^2}\right) = \frac{1}{4}\left(\frac{V^2}{\pi Gr^2}\right) = \frac{V^2}{4\pi Gr^2}. \quad (114)$$

Conclusion. Comparing this result with the standard requirement Eq. (107), we find an exact identity:

$$\boxed{\rho(r) \equiv \rho_N(r)}$$

The "Dark Matter" halo is thus identified not as a substance, but as the tension of the vacuum structure itself ($Q^2\rho_\Lambda$). $\square$

Corollary 19.3 (Redundancy of Dark Matter Hypotheses). *Since ρ provides 100% of the required dynamical mass density purely through geometric relations, the hypothesis of Dark Matter is redundant.*

20 Consolidating Mach's Principle: Derivation of Electron Mass: The Geometric Capacity Resonance

20.1 Topological Invariants

We derive the electron mass directly from the resonance between the internal geometry of the particle and the global geometry of the Universe. We define two dimensionless topological invariants describing the system state:

1. **The Internal Capacity Ratio ($\mathcal{R}_{int}$):** The ratio of the particle's [Electromagnetic Critical Radius \(\$R_q\$ \)](#) to its [Gravitational Schwarzschild Radius \(\$R_s\$ \)](#). This describes the intrinsic *intensity* of the particle's curvature.

$$\mathcal{R}_{int} = \frac{R_q}{R_s} \quad (115)$$

Using the definitions derived in Part I ($R_s = \kappa^2 r = 2Gm/c^2$) and Part III ($R_q = \kappa_q^2 r = 2\alpha^2 a_0 = 2e^2/4\pi\epsilon_0 mc^2$):

$$\mathcal{R}_{int} = \frac{\left(\frac{2e^2}{4\pi\epsilon_0 m_e c^2}\right)}{\left(\frac{2Gm_e}{c^2}\right)} = \frac{e^2}{4\pi\epsilon_0 Gm_e^2} \quad (116)$$

2. **The External Horizon Ratio ($\mathcal{R}_{ext}$):** The ratio of the Universal Hubble Horizon (R_H) to the particle's Unitary Energy Radius (r_e). This describes the particle's *scale* relative to the cosmos.

$$\mathcal{R}_{ext} = \frac{R_H}{r_e} = \frac{c/H_0}{\alpha\hbar/m_e c} \quad (117)$$

Note: We use the Unitary Energy Radius r_e (Classical Radius) here because it defines the localization scale of the rest energy E_0 , consistent with the inertial definition of mass.

20.2 Derivation of the Topological Operator Γ

A direct equality $\mathcal{R}_{int} = \mathcal{R}_{ext}$ is geometrically invalid because it equates a **Volumetric Potential Capacity** (scaling with mass density) with a **Linear Metric Distance**.

The topological operator Γ governs the dimensional reduction from the closed 2D potential surface of the particle (S^2) to the 1D linear macroscopic horizon of the Universe (R^1). This transformation is the composition of two orthogonal geometric projections.

1. Energetic Projection (Γ_E). The first operation maps the static capacity of the gravitational carrier (S^2) onto the dynamic kinematics of the quantum oscillator (S^1). From the [Closure Theorem](#), a fully balanced structural state is governed by the invariant geometric identity:

$$\kappa^2 = 2\beta^2 \quad (118)$$

The projection of the spherical potential amplitude onto the circular kinematic amplitude yields the energetic scaling factor:

$$\Gamma_E = \frac{\beta}{\kappa} = \frac{1}{\sqrt{2}} \quad (119)$$

2. Diametric Projection (Γ_T). The second operation maps the internal quantum phase topology onto the external macroscopic geometry. The internal structure is defined by its closed phase trajectory, a topological loop (S^1) with circumference $C = 2\pi r$. Conversely, the external macroscopic horizon (R_H) is measured as a linear radial axis, topologically equivalent to a 1D segment (R^1).

The geometric unrolling of a closed structural loop onto an open linear measurement axis requires the ratio of its equivalent linear span (diameter $D = 2r$) to its full phase length (circumference C):

$$\Gamma_T = \frac{D}{C} = \frac{2r}{2\pi r} = \frac{1}{\pi} \quad (120)$$

3. Total Holographic Operator (Γ). Since the energetic projection (Γ_E) and the spatial topological unwrapping (Γ_T) are independent, orthogonal operations, the complete holographic operator is their direct product:

$$\Gamma = \Gamma_E \cdot \Gamma_T = \frac{1}{\pi\sqrt{2}} \quad (121)$$

4. The Geometric Mach Equation and Dimensional Resolution (Γ^3). To map the internal curvature intensity onto the linear universal axis, we formulate the **Geometric Mach Equation**:

$$\mathcal{R}_{int} \cdot \Gamma^3 = \mathcal{R}_{ext} \quad (122)$$

In Equation (20.2), the Internal Capacity Ratio $\mathcal{R}_{int}$ incorporates the mass term squared in the denominator (m_e^{-2}). As established in [Part I section density](#) (Translating RG to Conventional Density), mass is not a fundamental scalar but a 3D translation artifact implicitly encoding a volumetric dependence ($1/r^3$).

Because $\mathcal{R}_{int}$ operates as a volumetric capacity metric, mapping it onto the 1D linear horizon ratio ($\mathcal{R}_{ext}$) creates a dimensional mismatch. To enforce geometric equilibrium, the 1D filter Γ must be applied across all three spatial dimensions embedded within the mass artifact. The operator is cubed (Γ^3), successfully unpacking the 3D-mass variable into a 1D relational length without empirical fitting parameters.

20.3 Derivation

Substituting the definitions into Eq. (20.2):

$$\left(\frac{e^2}{4\pi\epsilon_0 G m_e^2} \right) \cdot \Gamma^3 = \frac{m_e c^2}{H_0 \alpha \hbar}$$

Using the definition of the fine structure constant $\alpha = \frac{e^2}{4\pi\epsilon_0 \hbar c}$, we simplify the Left Hand Side (LHS):

$$\text{LHS} = \frac{\alpha \hbar c}{G m_e^2} \cdot \Gamma^3$$

Now equate to RHS:

$$\frac{\alpha \hbar c}{G m_e^2} \cdot \Gamma^3 = \frac{m_e c^2}{H_0 \alpha \hbar}$$

Isolating the mass term (m_e^3) by multiplying both sides by m_e^2 and rearranging:

$$m_e^3 = \Gamma^3 \cdot \frac{(\alpha \hbar c)(H_0 \alpha \hbar)}{G c^2}$$

$$m_e^3 = \Gamma^3 \cdot \frac{H_0 (\alpha \hbar c)^2}{G c^3}$$

Taking the cube root of both sides:

$$m_e = \Gamma \left(\frac{H_0 (\alpha \hbar c)^2}{G c^3} \right)^{1/3} \quad (123)$$

Substituting the explicit geometric projection $\Gamma = \frac{1}{\pi\sqrt{2}}$:

$$\boxed{m_{e(WILL)} = \frac{1}{\pi\sqrt{2}} \left(\frac{H_0 (\alpha \hbar c)^2}{G c^3} \right)^{1/3}} \quad (124)$$

20.4 Numerical Verification

Using the value of H_0 derived in Part II from the CMB temperature ($H_0 \approx 2.208 \times 10^{-18} \text{ s}^{-1}$):

- **Calculated Mass:** $m_{e(WILL)} \approx 9.064 \times 10^{-31} \text{ kg}$
- **Observed Mass:** $m_{e(CODATA)} \approx 9.109 \times 10^{-31} \text{ kg}$
- **Precision:** The deviation is $\approx 0.49\%$.

20.5 Conclusion

This result confirms that the electron's mass is the consequence of a holographic equilibrium. The intrinsic curvature intensity of the particle, when projected from its closed topology onto the linear universe via Γ^3 , exactly balances the scale of the cosmic horizon.

This completes the Theoretical Ouroboros:

1. **Micro $\rightarrow$ Macro:** $\alpha \rightarrow H_0$ (Part II).
2. **Macro $\rightarrow$ Micro:** $H_0 \rightarrow m_e$ (Part II Conclusion).

Provided evidence strongly suggests that the Universe is a single, resonant, self-defining geometric relational unity.

21 General Discussion

The analysis presented in this work yields three structurally significant results that challenge the foundations of the Standard Model (Λ CDM) and standard Modified Gravity (MOND):

The Failure of "Dark" Parameters. Λ CDM, when constrained by global scaling laws rather than individual halo fitting, fails to reproduce galactic rotation curves (systematic bias -6.83 km/s). Furthermore, the requirement of $\approx 26\%$ Dark Matter for CMB acoustics is shown to be redundant: the acoustic peaks are accurately recovered by a pure baryonic load ($\approx 4.2\%$) on a tensioned S^2 topology.

The Resolution of the Gravity Boost Tension. The systematic failure of standard MOND to predict the Wide Binary boost factor ($\gamma \approx 1.8$ vs observed 1.4) illustrates the limitations of treating gravity as a modified force field with a universal acceleration scale. WILL RG reveals that the coupling to the Horizon is topology-dependent. Phenomenological models fail because they lack the geometric ontology to distinguish the different topological coupling weights (Ω) connecting local systems to the Cosmic Fundamental Tone ($f_0 = \frac{H_0}{2\pi}$).

The Thermodynamic Origin of Dynamics. Unlike phenomenological models that fit acceleration scales (a_0) to minimize residuals, WILL RG derives the acceleration scale $a_{Mach} = cH_0/2\pi$ entirely from the CMB temperature and α . The fact that this thermodynamically derived scale eliminates the bias in rotation curves ($+0.53$ km/s vs > 5 km/s for MOND) serves as strong evidence that gravity is not an isolated force, but a holographic response to the global energy state.

21.1 Orthogonal Validation: The Geometric Invariant α

Epistemological test of any physical theory is the independence of its observables. In standard Λ CDM cosmology, the Hubble parameter H_0 and the acoustic scale ℓ_{peak} are coupled variables; they are derived from a multi-parameter fit where altering H_0 shifts the angular diameter distance, forcing a recalibration of matter densities (Ω_m, Ω_Λ) to maintain consistency with the CMB spectrum.

In WILL Relational Geometry, we observe a strict **Orthogonal Decoupling** of these parameters. The derivations presented in this paper proceed along two mathematically disjoint paths, yet converge on the same observational reality (Planck 2018).

21.2 Path A: The Thermodynamic Limit (H_0)

The expansion rate H_0 is derived (3) as a limit of **Energy Saturation**. It relies on thermodynamic input:

$$H_0 = \sqrt{8\pi G \cdot \rho_{max}} \quad \text{where} \quad \rho_{max} = \frac{\rho_\gamma(T_0)}{3\alpha^2} \quad (125)$$

Input: T_{CMB} (Temperature) and α .

Result: $H_0 \approx 68.15$ km/s/Mpc.

21.3 Path B: The Topological Resonance (ℓ_{vac})

The acoustic scale is derived (16) as a limit of **Resonant Structure**. It relies exclusively on the kinematic scale and geometric tension:

$$\ell_{vac} = \frac{1}{\alpha} \underbrace{(1 + \Omega_\Lambda)}_{\text{Impedance}} \quad (126)$$

Input: α and $\Omega_\Lambda = 2/3$. (Independent of T_{CMB} or ρ_γ).

Result: $\ell_{vac} \approx 228.39$ (unloaded).

21.4 The Independence Theorem

Crucially, the calculation of the acoustic peak does not require knowledge of the expansion history (H_0) or the age of the Universe. Conversely, the calculation of H_0 does not require knowledge of the vacuum tension. The correlation between these macroscopic observables is not a result of parameter fitting, but a consequence of their common origin in the geometric invariant α :

$$\boxed{\frac{\partial H_0}{\partial \ell_{peak}} = 0} \quad (127)$$

The fact that two independent derivation paths - one thermodynamic and one topological - yield results matching high-precision empirical data ($\delta \approx 1.0\%$ for H_0 , $\delta \approx 0.004\%$ for ℓ) constitutes a strong argument for the generative WILL RG.

21.5 Sensitivity Analysis: The Major Constraints on Dark Matter

WILL RG allows us to perform a precise sensitivity analysis regarding the mass content of the Universe. The acoustic peak position is determined by the inertial loading of the vacuum tension:

$$\ell_{observed} = \ell_{vac} \cdot \sqrt{\frac{\Omega_\Lambda}{\Omega_\Lambda + \Omega_{load}}} \quad (128)$$

We compare two scenarios: one containing only BBN-derived Baryons (WILL RG), and one including the standard Dark Matter halo (Λ CDM).

Scenario	Load	Peak	Deviation
	(Ω_{load})	(ℓ)	from Planck
WILL RG (Pure Baryons)	0.048	220.55	-0.02%
Standard Model (Λ CDM + DM)	0.308	188.89	-14.37%

Table 10: **Baryonic Consistency Test.** The vacuum tension $\Omega_\Lambda = 2/3$ accurately reproduces the first acoustic peak ($\ell \approx 220.6$) only when loaded with standard baryonic matter ($\Omega_b \approx 0.048$ From Nuclear Physics of the Yearly Universe). The addition of Dark Matter ($\Omega_{dm} \approx 0.26$) creates excessive inertial drag, shifting the peak to $\ell \approx 189$, which is in tension with observational data.

Conclusion: The high tension of the structural projection ($\Omega_\Lambda = 2/3$) suggests that Dark Matter hypothesis is unnecessary. The Universe operates as a Baryonic oscillator on a tense relational geometry.

Parameter or Observable	Derived Theoretical Value	Empirical Comparison Value	System or Dataset	Deviation or Accuracy	Physical Formulation
1. Hubble Constant (H_0)	68.15 km/s/Mpc	67.4 ± 0.5 km/s/Mpc	Planck 2018	+1.0%	Geometric saturation density derived from CMB temperature and α
2. CMB First Acoustic Peak (l_1)	220.55	220.60	Planck 2018	$\approx 0.02\%$	Resonant harmonics of an S^2 topology loaded by 4.2% baryonic mass
3. CMB Quadrupole Power ($D_{l=2}$)	0.156–0.320 (Corridor)	≈ 0.20	Planck 2018	Within predicted corridor	Vacuum tension acting as a high-pass filter on a tensioned S^2 membrane
4. Galactic Rotation Curves Bias	0.70×10^{-10} m/s ² (a_k)	-2.26 km/s (Bias)	SPARC (175 galaxies)	RMSE ≈ 0.065 dex	Structural Projection Resonance with Universal Fundamental Tone
5. Solar Orbital Velocity	226.4 km/s	229 ± 6 km/s	Gaia DR3 / Milky Way	Excellent agreement	Projection Resonance between local potential and global horizon
6. Wide Binary Gravity Boost (γ)	≈ 1.47	$\approx 1.45\text{--}1.55$	Gaia DR3 / Chae 2023	Exact Agreement	Kinetic Resonance Scale (S^1 carrier coupling weight 1/3)
7. Type Ia Supernova Distance Modulus	Offset expected ≈ 0.150 mag	Raw residual ≈ -0.151 mag	Pantheon+	Shape deviation ≤ 0.02 mag	Geometric Energy Budget Partitioning (2:1 ratio of S^2 tension to kinetic mass)
8. Strong Lensing Einstein Radius	1.46''	$1.43 \pm 0.01''$	SDSSJ0946+1006 (SLACS)	$\approx 2\%$	Phantom Inertia (Q^2) acting as universal refractive medium
9. Recombination Epoch	$\approx 364,860$ years	$\approx 378,000$ years	Standard Cosmological Dating	$\approx 3.5\%$	Unit Phase Condition ($\Omega_{crit} = 1$ radian) where arc length equals radius of curvature
10. Electron Mass (m_e)	9.064×10^{-31} kg	9.109×10^{-31} kg	CODATA	$\approx 0.49\%$	Holographic Projection Principle / Geometric Capacity Resonance

21.6 Final Conclusion

We have demonstrated that the "Dark Matter" phenomenon is the observational signature of scale-invariant geometric closure. By replacing the dark sector with the rigid geometry of the Global Horizon, we achieve a unification of Baryonic physics across 20 orders of magnitude.

Just as the electron must satisfy the standing wave condition to exist as a bound state within the atom, the galaxy must satisfy the frequency resonance condition to exist as a bound state within the Universe. The precision of these predictions, achieved without any free parameters, strongly suggests that the paradigm of "Dark" phenomenology is becoming obsolete, superseded by a transparent **Relational Geometric Ontology**.

Code and data are fully open-source at: <https://willrg.com>

References

- [1] Planck Collaboration. (2020). Planck 2018 results. VI. Cosmological parameters. *Astronomy & Astrophysics*, 641, A6.
- [2] Lelli, F., McGaugh, S. S., & Schombert, J. M. (2016). SPARC: Mass Models for 175 Disk Galaxies with Spitzer Photometry and Accurate Rotation Curves. *The Astronomical Journal*, 152(6), 157.
- [3] Chae, K. H. (2023). Breakdown of the Newton–Einstein Standard Gravity at Low Acceleration in Internal Dynamics of Wide Binary Stars. *The Astrophysical Journal*, 952(2), 128.
- [4] Hernandez, X., et al. (2023). Internal kinematics of Gaia DR3 wide binaries. *Monthly Notices of the Royal Astronomical Society*, 525(2), 2615.
- [5] Bolton, A. S., et al. (2008). The Sloan Lens ACS Survey. V. The Full Sample of 70 Lens Candidates and Strong Lensing Mass Models. *The Astrophysical Journal*, 682(2), 964.
- [6] Auger, M. W., et al. (2009). The Sloan Lens ACS Survey. IX. Colors, Lensing and Stellar Masses of Early-Type Galaxies. *The Astrophysical Journal*, 705(2), 1099.
- [7] Li, P., Lelli, F., McGaugh, S. S., & Schombert, J. M. (2020). A Comprehensive Catalog of Dark Matter Halo Models for SPARC Galaxies. *The Astrophysical Journal Supplement Series*, 247(1), 31.
- [8] Wang, D.-C., Xu, F., & Luo, X. (2020). Comparison of Modeling SPARC spiral galaxies' rotation curves with different dark matter and MOND models. *arXiv preprint arXiv:2008.04795*.
- [9] Milgrom, M. (2001). MOND - A Pedagogical Review. *NED Level 5 Review*.
- [10] F. Iocco, M. Pato, and G. Bertone, “Evidence for dark matter in the inner Milky Way,” *Nature Physics*, vol. 11, no. 3, pp. 245–248, 2015.
- [11] Peebles, P. J. E. (1968). Recombination of the Primeval Plasma. *The Astrophysical Journal*, 153, 1.
- [12] DESI Collaboration (2025). DESI DR2 Results II: Measurements of Baryon Acoustic Oscillations and Cosmological Constraints. *Physical Review D*, 112, 083515.
- [13] Treu, T., Koopmans, L. V. E., Bolton, A. S., Burles, S., & Moustakas, L. A. (2006). The Sloan Lens ACS Survey. II. Stellar Populations and Internal Structure of Early-Type Lens Galaxies. *The Astrophysical Journal*, 640(2), 662.
- [14] Koopmans, L. V. E., et al. (2009). The Structure and Dynamics of Massive Early-Type Galaxies: On Homology, Isothermality, and Isotropy Inside One Effective Radius. *The Astrophysical Journal Letters*, 703(1), L51.