

WILL Part I

Relational Geometry

Anton Rize
antonrize@willrg.com *

October 2025

Abstract

This paper, the first in the **WILL Trilogy**:

[Part I: Relational Geometry](#);

[Part II: Relational Cosmology](#);

[Part III: Relational Quantum Geometry](#).

Applying extreme methodological constraints and establishes *Relational Geometry (RG)*: a foundational framework where spacetime is an emergent property of relational energy transformations. This shift establishes an ontological transition from *descriptive* to *generative* physics: instead of introducing laws to model observations, it derives them as necessary consequences of Relational Geometry itself - turning physics from a catalogue of phenomena into the logical unfolding of inevitable geometrical constraints on closed relational carriers S^1 (directional) and S^2 (omnidirectional).

Without metrics, tensors, or free parameters, it reproduces Lorentz factors, the energy-momentum relation, Schwarzschild and Einstein field equations via the dimensionless projections β (kinematic) and κ (potential). All known GR critical surfaces (photon sphere, ISCO, horizons) emerge as simple fractions of (κ, β) from the single closure law $\kappa^2 = 2\beta^2$ (topologically derived, virial-like theorem (14.3)).

WILL Part I offers new perspective to several long-standing problems, including:

- **Resolution of the Classical $M \sin i$ Degeneracy** (by introducing independent invariants²⁵),
- **Resolution of GR singularities** (via naturally bounded $\rho_{\max} = \frac{c^2}{8\pi G r^2}$ ³⁰),
- **Derivation of the equivalence principle** (from the common channel of rest-invariant state (13.3) scaling^{13.2}),
- **Removal of local energy density ambiguity** $\rho = \frac{\kappa^2 c^2}{8\pi G r^2}$ (28.1),
- **Revelation of a clear relational symmetry between kinematic and potential projections** (12),
- **Developed closed system of equations for Relational Orbital Mechanics (ROM)** without G, mass and differential formalism (18).
- **Establishment of a computationally simpler and ontologically consistent foundation** for subsequent papers on cosmology ([Part II](#)) and quantum mechanics ([Part III](#)).

A Direct Note to the Reader: Why should you read a 60-page independent paper?

The fact that you opened this file and reading these lines already separates you from the majority, and for that, you have my respect. However, sixty pages of unknown research is a substantial commitment, and you should not engage it blindly.

I recommend the following: Be honest with yourself. Define the exact criteria this paper must satisfy to justify the investment of your time. Then, ask [WILL-AI](#) if this work meets your standards. The AI is trained strictly on this document and will give you an honest, unbiased answer based on its contents.

If you are still in doubt, browse [willrg.com](#), interact with the [Galactic Dynamics Lab](#), trace the derivation chain using the [Logos Map](#), review the [falsifiable predictions](#), and have a chat or few with [WILL-AI](#).

Regardless of your decision, I hope you find what you are seeking.

Anton Rize

*This work is archived on Zenodo: ,

Contents

1 Foundational Methodological Principles	5
1.1 What is Energy in Relational Framework?	5
2 Ontological Blind Spot In Modern Physics	6
2.1 Historical Pattern: breakthroughs delete, not add	6
2.2 The contemporary split: an unpaid ontological bill	6
2.3 Empirical bankruptcy of the separation	6
2.4 Consequence	7
3 Unifying Principle Removing the Hidden Assumption	7
3.1 False Separation	7
4 Deriving the WILL Structure	8
4.1 Derivation of the Relational Carriers	8
5 The Amplitude-Phase Duality	9
5.1 Consequence: Relativistic Effects	10
6 Energy as a Relation - What κ and β Actually Mean	10
6.1 Ontological Status of the Relational Carriers S^1 and S^2	11
7 Emergence of Spacetime	11
8 Kinetic Energy Projection on S^1	12
8.1 Geometric Nature of Rest Energy State	12
8.2 The Geometric and Historical Nature of Mass	12
8.3 Energy–Momentum Relation	13
9 Potential Energy Projection on S^2	13
9.1 Gravitational Meridional Section of S^2	14
9.2 Consequence: Gravitational Effects	14
9.3 Gravitational Tangent Formulation	14
10 Geometric composition of SR and GR factors	15
11 Total Relational Shift Q	15
11.1 Principle of Relational Reciprocity	16
12 Clear Relational Symmetry Between Kinematic and Potential Projections	16
13 Equivalence Principle as Derived Identity	17
14 Unification of Projections: The Geometric Exchange Rate	18
14.1 Derivation of the Energetic Closure Condition	19
14.2 Uniqueness of the Exchange Rate (No Hidden Weighting)	19
15 Energy-Symmetry Law	20
15.1 Causal Continuity and Energy Symmetry	20
15.2 The Specific Energy Transfer (ΔE):	21
15.3 Physical Meaning of the Factor $\frac{1}{2}$	21
15.4 Universal Speed Limit as a Consequence of Energy Symmetry	21
15.5 Single-Axis Energy Transfer and the Nature of Light	22
16 Classical Keplerian Energy and Minkowski Interval as Single-Point Limits of the Relational Energy-Symmetry Law	23
16.1 Classical Result with Surface Reference	23
16.2 Projection Parameters and Minkowski-like Form	23
16.3 Physical Interpretation	23

17 Lagrangian and Hamiltonian as Single-Point Limits of the Relational Energy-Symmetry Law	24
17.1 Definitions of Parameters	24
17.2 The Relational Lagrangian	24
17.3 First Ontological Collapse: The Newtonian Lagrangian	24
17.4 Second Ontological Collapse: The Hamiltonian	25
17.4.1 Interpretation	25
17.5 Third Ontological Collapse: Derivation of Newton's Third Law	26
18 Relational Orbital Mechanics (R.O.M.)	26
19 Operational Independence and the Role of Constants	27
19.1 Derivation of the Spectroscopic Shifts	27
19.2 Operational Measurability of Relational Projections	28
19.3 Derivation of Eccentricity	29
19.4 Orbital Precession as Relational Phase Accumulation	30
19.5 Kinematic Mechanism of the Dynamic Horizon	31
19.6 Algebraic Determination of Absolute System Scale	32
19.6.1 Method A: Differential (Two-Point Method)	32
19.6.2 Method B: Geometric Resonance (Balance Point Method)	33
19.6.3 Method C: Instantaneous (Arbitrary Phase Method)	33
19.7 The Role of G as Translation Constant	34
19.8 Summary	35
20 Relational Parameterization and the Fundamental Primitives	35
21 Rotational Systems (Kerr Without Metric)	37
21.1 Contextual Bounds	37
21.2 Event Horizon	37
21.3 Ergosphere	38
21.4 Ring Singularity	38
21.5 Naked Singularity	38
21.6 Chiral Asymmetry: The Algebraic Origin of Frame-Dragging	38
21.6.1 Composite Kinematic Projection	38
21.6.2 The Algebraic Symmetry Breaker (Cross-Term)	38
21.6.3 Orbital and Optical Splitting	39
21.7 The Relationship Between $\kappa > 1$ and Rotation	39
22 Time and Phase Evolution	39
22.1 Continuous Model and Closed-Form Evaluation	39
22.2 Irreducible Transcendence of the Inverse Problem	40
22.3 Operational Discrete Alternative	40
23 Strong Field Test (S2 Star)	40
24 Blind Prediction for S4716 (In Silico Experiment)	41
24.1 Operational Derivation from Observation	41
24.2 Geometric De-projection of Single-Epoch Velocity	41
24.3 Propagation to Periapsis	41
24.4 Prediction	42
25 Sidestepping the Classical $M \sin i$ Degeneracy	42
25.1 The 130-Year Limitation	42
25.2 The Interior Kinematic Projection	42
25.3 Non-Linear Relational Invariants	42
25.4 Empirical Validation: Bayesian MCMC Analysis on S0-2 Data	43
25.4.1 Ablation Study Results	43
25.4.2 Topology of the Posterior and the Degeneracy Ridge	43
25.4.3 Independent Validation via Background Drift (v_{z0})	44
25.5 Independent Validation via Synthetic 1PN Data	45
25.5.1 Data Generation and Observational Strategy	45
25.5.2 Three-Stage Algebraic Extraction	45
25.5.3 Blind Test Results	45

25.6 Analytical Extraction of the Inclination Angle: The Decryption Invariant	46
25.6.1 The Observer Equations at Extrema	46
25.6.2 Algebraic Decoupling of β and i	46
25.6.3 Analytical Extraction of $\sin i$	46
25.6.4 Numerical Verification: S0-2 Extrema Synthesis	47
25.7 Conclusion	47
26 Derivation of Gravitational Deflection and Lensing	47
26.1 Gravitational Deflection and Interaction Gradient	47
26.2 Gravitational Lensing and the Einstein Ring	49
27 Geometric Mechanism of Orbital Decay and Radiation	49
28 Derivation of Density, Mass, and Pressure	50
28.1 Derivation of Density	50
28.2 Self-Consistency Requirement	51
28.3 Pressure as Surface Curvature Gradient	51
29 Unified Geometric Field Equation	52
29.1 Field Equation and Matter Sources	52
30 No Singularities, No Hidden Regions	52
31 Theoretical Ouroboros	53
32 W_{ILL}: Unity of Relational Structure	53
32.1 Interpretive Note: The Name "WILL"	55
33 Ontological Shift: From Descriptive to Generative Physics	55
34 Conclusion	57
35 References:	58
.	
.	
.	
.	
.	
.	

Following yet untamed thought

"There is no such thing as an empty space, i.e., a space without field. . . . Space-time does not claim existence on its own, but only as a structural quality of the field."

— *Albert Einstein, Relativity: The Special and the General Theory* (Appendix V: "Relativity and the Problem of Space"), 1952 edition, Methuen (London), p. 155; based on earlier 1920 additions.

IMPORTANT:

This document must be read **literally**. All terms are defined within the relational framework of WILL Relational Geometry. Any attempt to reinterpret them through conventional notions (*absolute energies, external backgrounds, hidden containers*) will produce distortions and misreadings. Just like responsibility of formulating lies with the author, the responsibility of interpretation lies with the reader: take the words as written, not as filtered through prior formalisms.

1 Foundational Methodological Principles

Every logical step is subject to empirical audit: if any consequence of this chain contradicts observation, the chain fails.

As humans we often project ourselves on observed phenomenon, obscuring the true nature of reality behind our anthropic tendency's. In order to prevent such misleading practises, we have to establish extreme methodological constrains. This constrains are not statement about the nature of reality, but the rule of logical purity for constructing a theory. Without strict methodology any theoretical construct is risking that excessive freedom of interpretations will conclude with the framework that tells us more about ourselves then about nature of the Universe. The philosophical reasoning behind this particular methodological constrains might deserve its own dedicated publication and can be challenged, but this research main domain lays within physical reality that can be empirically tested, there for empirical alignment is the ultimate test of the methods presented.

Principle 1.1 (Epistemic Hygiene). Epistemic Hygiene as Refusal to Import Unjustified Assumptions. *This line of reasoning derive physics by **removing hidden assumptions**, rather than introducing new postulates. This construction is deliberate and contains zero free parameters. This is not a simplification - it is a deliberate epistemic constraint. No assumptions are introduced and no constructs are retained unless they are geometrically or energetically necessary.*

Principle 1.2 (Ontological Minimalism). *Any fundamental theory must proceed from the minimum possible number of ontological assumptions. The burden of proof lies with any assertion that introduces additional complexity or new entities.*

Principle 1.3 (Relational Origin). All physical quantities must be defined by their relations. *Any introduction of absolute properties risks reintroducing metaphysical artefacts and contradicts the foundational insight of relationalism.*

Principle 1.4 (Simplicity). **Everything** must be expressed in the **simplest** form possible. *Any unjustified complexity risks reintroducing metaphysical artefacts and contradicts the foundational insight of Epistemic Hygiene.*

Principle 1.5 (Mathematical Transparency). 1. *Every mathematical phrase, operational choice, or identity carries its own ontological statement.*

2. *Each mathematical object must correspond to explicitly identifiable relation between observers with transparent ontological origin.*
3. *Every symbol must be anchored to unique physical idea.*
4. *Introducing symbols without explicit necessity constitutes **Over-parameterization**: the proliferation of symbols without corresponding physical meaning.*
5. *Number of symbols = Number of independent physical ideas.*

Mathematics is a language, not a world. Its symbols must never outnumber the physical meanings they encode.

1.1 What is Energy in Relational Framework?

Across all domains of physics, one empirical fact persists: in every closed system there exists a quantity that never disappears or arises spontaneously, but only transforms in form. This invariant is observed under many guises — kinetic, potential, thermal, chemical — yet all are interchangeable, pointing to a single underlying structure.

Crucially, this quantity is never observed directly, but only through *differences between states*: a change of velocity, a shift in configuration, a transition of phase. Its value is relational, not absolute: it depends on the chosen frame or comparison, never on an object in isolation.

Moreover, this quantity provides continuity of causality. If it changes in one part of the system, a complementary change must occur elsewhere, ensuring the unbroken chain of transitions. Thus it is the bookkeeping of causality itself.

From these empirical and relational facts the definition follows unavoidably:

Energy :

Definition 1.6 (Energy).

*Energy is the invariant relational measure
of state transformation
within a topologically closed system.*

2 Ontological Blind Spot In Modern Physics

The standard formulation of General Relativity often relies on the concept of a pre-existing, container-like spacetime that then gets “filled” with fields and matter. This is in direct tension with the Relational Principle 1.3.

The standard derivation of the Einstein field equations begins with the Einstein-Hilbert action, which is built upon the metric as the fundamental variable. This metric is defined on a smooth manifold that is assumed to exist *a priori*. This manifold, even without a metric, carries topological and differential structure - an absolute scaffold. This is an ontological extra. It violates Principle 1.2 (Ontological Minimalism) because it introduces an entity (the manifold) that is not derived from relational observations.

In the standard formulation, the stress-energy tensor is derived from the variation of the matter Lagrangian with respect to the metric. This assumes that energy is a property of matter fields that can be localized in spacetime. However, this localization is frame-dependent (via the equivalence principle) and leads to well-known problems such as the non-uniqueness of the gravitational energy-momentum pseudotensor. This is a direct violation of the Relational Origin principle 1.3: energy is treated as an absolute property of matter rather than a relational measure.

We proceed from strict epistemic minimalism, disallowing all background structures, even hidden or asymptotic ones.

2.1 Historical Pattern: breakthroughs delete, not add

- **Copernicus** eliminated the Earth/cosmos separation.
- **Newton** eliminated the terrestrial/celestial law separation.
- **Einstein** eliminated the space/time separation.
- **Maxwell** eliminated the electricity/magnetism separation.

Each step widened the relational circle and reduced the number of unexplained absolutes. The spacetime–energy split is the only survivor of this pruning sequence.

2.2 The contemporary split: an unpaid ontological bill

All present-day theories (SR, GR, QFT, CDM, Standard Model) are built with a *bi-variable* syntax:

$$\underbrace{\text{fixed manifold} + \text{metric}}_{\text{structure}} + \underbrace{\text{fields} + \text{constants}}_{\text{dynamics}}.$$

No observation demands this duplication; it is retained purely because the resulting Lagrangians are empirically adequate *inside* the split. The split is therefore *not* an empirical discovery but an unpaid ontological debt.

2.3 Empirical bankruptcy of the separation

- **Local energy conservation** verified only *after* the metric is declared fixed; no experiment varies the *volume* of flat space and checks calorimetry.
- **Universality of free fall** tests $m_i = m_g$ numerically, not the claim that inertia resides *in* the object rather than in a geometric scaling relation.

- **Gravitational-wave polarisations** test spin content, not ontology; extra modes can still be called “matter on spacetime”.
- **Casimir/Lamb shift** measure *differences* of vacuum energy between two geometries; the absolute bulk term is explicitly subtracted, leaving the split intact.

In short, every “test” is an *internal consistency check* of a formalism that already presupposes two substances. None constitute *positive evidence* for the split.

2.4 Consequence

Until an experiment varies the amount of space while holding everything else fixed, the spacetime–energy separation remains an *un-evidenced metaphysical postulate*—the last geocentric epicycle in physics.

Summary

Any attempt to treat “*spacetime structure*” as separate from “*dynamics*” smuggles in a background container that is not justified by the phenomena. This violates epistemic hygiene: it introduces an ontological artifact without necessity. Eliminating this separation compels the identification of structure and dynamics as two aspects of a single entity.

Ontological Minimalism:

If no empirical or logical ground justifies the distinction between (structure) and (dynamics), the distinction must be dissolved.

$$\text{SPACETIME} \equiv \text{ENERGY}$$

*This equivalence is not algebraic but ontological;
spacetime and energy are two descriptive projections of a single invariant entity we call:*

WILL

3 Unifying Principle Removing the Hidden Assumption

3.1 False Separation

Lemma 3.1 (False Separation). *Any model that treats processes as unfolding within an independent background necessarily assigns to that background structural features (metric, orientation, or frame) not derivable from the relations among the processes themselves. Such a background constitutes an extraneous absolute.*

Proof. Suppose an independent background exists. Then at least one of its structural attributes - metric relations, a preferred orientation, or a class of inertial frames - remains fixed regardless of inter-process data. This attribute is not relationally inferred but posited a priori. It thereby violates the relational closure principle: it introduces a non-relational absolute external to the system. Hence the separation is illicit. $\square$

Corollary 3.2 (Structure–Dynamics Coincidence). *To avoid the artifact of Lemma 3.1, the structural arena and the dynamical content must be identified: geometry is energy, and energy is geometry.*

Principle 3.3 (Unifying Ontological Principle: Removing the Hidden Assumption).

$$\text{SPACETIME} \equiv \text{ENERGY}$$

This is not introduced as a new ontological entity but as a Principle with negative ontological weight: it removes the hidden unjustified separation between “structure” and “dynamics.” Spacetime is not a “container” but emergent relational energy structure.

Remark 3.4 (Auditability). *Principle 3.3 is foundational but testable: it is subject to (i) geometric audit (internal logical consequences) and (ii) empirical audit (agreement with empirical data).*

Definition 3.5 (WILL). **WILL** $\equiv$ **SPACE-TIME-ENERGY** is the technical term we use for unified relational structure determined by 3.3. All physically meaningful quantities are relational features of WILL; no external container is permitted.

Summary:

This Principle does not add, it subtracts: it removes the hidden assumption. Structure and dynamics are two aspects of a single entity that we call - WILL.

4 Deriving the WILL Structure

Having established our Principle 3.3 by removing the illicit separation of structure and dynamics, we now proceed to derive its necessary geometric and physical consequences. We will demonstrate that this single principle is sufficient to enforce the closure, conservation, and isotropy of the relational structure, leading to a unique set of geometric carriers for energy.

Lemma 4.1 (Closure). *Under 3.3, WILL is self-contained: there is no external reservoir into or from which the relational resource can flow.*

Proof. If WILL were not self-contained, there would exist an external structure mediating exchange. That external structure would then serve as a background distinct from the dynamics, contradicting Corollary 3.2. $\square$

Lemma 4.2 (Conservation). *Within WILL, the total relational “transformation resource” (energy) is conserved.*

Proof. By Lemma 4.1, no external fluxes exist. Any change in one part of WILL must be balanced by complementary change elsewhere. Hence a conserved global quantity is enforced at the relational level. $\square$

Lemma 4.3 (Isotropy from Background-Free Relationality). *If no external background is allowed (Cor. 3.2), then no direction can be a priori privileged. Thus the admissible relational geometry of WILL must be maximally symmetric (isotropic and homogeneous) at the level at which it encodes the conserved resource.*

Proof. Any alleged privilege that cannot be constructed from relations among participants is unobservable (pure gauge). Therefore the carrier used to encode the conserved WILL resource must have no intrinsic privileged direction or point. Therefore the encoding geometry must be maximally symmetric. $\square$

4.1 Derivation of the Relational Carriers

The lemmas of Closure, Conservation, and Isotropy (Lemmas 4.1–4.3–4.2) establish the necessary properties of geometric carriers of the relational resource (energy).

Relational Carrier Conventions:

All references to “carriers” within WILL RG research are to be read in the strict relational sense:

- **Degree of freedom (DOF):** An n -dimensional relational carrier encoding the conserved transformation resource.
- **Direction:** An oriented relational ray. Opposite rays are physically distinct and not identified.
- **Closed carrier:** Compact and without boundary; resource cannot leak into an external reservoir.
- **No background:** No external embedding space. All geometric structure is reconstructible from relations between participants.
- **Maximal symmetry:** The carrier is homogeneous and isotropic with respect to oriented directions.
- **Minimal relational carrier:** A closed, maximally symmetric carrier using exactly the required DOF without arbitrary parameters.

Theorem 4.4 (Minimal Relational Carriers of the Conserved Energy Resource). *The minimal relational carriers satisfying Closure, Conservation, Maximal Symmetry, and Mathematical Transparency are:*

- (a) S^1 for directional (Kinematic) relational transformation;
- (b) S^2 for omnidirectional (Potential) relational transformation.

Proof. The proof classifies the minimal types of relations and applies the derived constraints:

- (a) **1-DOF (Kinematic) Relation:** The minimal non-trivial relation involves 1 degree of freedom (1-DOF): a sequence of state transformations.

By Lemma 4.1 (Closure), this sequence must form a loop to prevent resource leakage. A discrete sequence (a finite cyclic group C_n) requires assigning a specific integer value n to the number of states. Principle 1.5 (Mathematical Transparency) prohibits the introduction of arbitrary numerical parameters without unique physical justification. To achieve a parameter-free structure, the state cardinality must remain undefined, forcing the continuum limit.

By Lemma 4.3 (Isotropy), this continuous 1-DOF loop must be maximally symmetric. The unique connected, closed, parameter-free 1-carrier is the continuous S^1 (circle).

- (b) **2-DOF (Potential) Relation:** The next minimal relation involves 2 degrees of freedom (2-DOF): the omnidirectional distribution of the transformational resource from a primary state.

This requires a 2-dimensional carrier. By Principle 1.5, any discrete tiling of this distribution introduces arbitrary parameters (number of nodes or faces), strictly forcing a continuous 2-carrier.

By Lemma 4.1, this continuous 2-carrier must be closed. By Lemma 4.3, it must be maximally symmetric (isotropic distribution of potential).

The topological classification of closed 2-carrier under the requirement of maximal continuous symmetry eliminates anisotropic structures (e.g., the torus T^2 , which possesses preferred axes). The unique parameter-free, maximally symmetric 2-carrier is S^2 (2-sphere).

Ontological Minimalism and Mathematical Transparency uniquely enforce S^1 and S^2 as the minimal relational carriers without invoking metric geometry *a priori*. $\square$

Corollary 4.5 (Uniqueness). *Under Principle 3.3 with Closure, Conservation, Isotropy, and Mathematical Transparency, S^1 and S^2 are necessary relational carriers for directional and omnidirectional modes of energy transformation.*

Remark 4.6 (Non-spatial Reading). *S^1 and S^2 are not to be interpreted as spacetime geometries. They are relational carriers encoding the closure, conservation, and isotropy of the transformational resource. Spatial and temporal notions are emergent descriptors of patterns within WILL.*

Logical Chain Summary

Methodological Constraints
$\downarrow$ (apply to existing physics)
False Separation Identified
$\downarrow$ (dissolve)
SPACETIME $\equiv$ ENERGY
$\downarrow$ (derive consequences)
Closure + Conservation + Isotropy
$\downarrow$ (classify carriers)
S^1 (1-DOF kinematic) + S^2 (2-DOF potential)

5 The Amplitude-Phase Duality

The manifestation of any system is distributed between its internal (Phase) and external (Amplitude) aspects. This single geometric constraint gives rise to the core phenomena of modern physics:

Lemma 5.1 (Duality of Evolution). *The identification of spacetime with energy and its transformations necessitates two complementary relational measures:*

1. the **amplitude** of transformation (external shift), and
2. the **phase** of transformation (internal order).

Proof. Any complete description of transformation must specify both what changes and how that change is internally ordered. A single measure cannot capture both. Relational Carriers S^1 and S^2 both **provide** the minimal geometry enforcing such complementarity: **their** orthogonal projections furnish precisely two non-redundant coordinates. $\square$

The orthogonal decomposition of the relational carriers S^1 and S^2 reveals a functional duality. Every physical state is a superposition of two projections:

Definition 5.2 (The Amplitude Projection (External Interaction)). Denoted by β (kinematic) and κ (potential). This component measures the extent of external relation. It manifests as momentum or potential intensity. Physically, it represents the system's relational "shift" from the **Observer's relational Origin** (the rest frame).

$$\text{Amplitude} \rightarrow \text{External Power (Kinetic/Potential)}$$

Definition 5.3 (The Phase Projection (Internal Evolution)). Denoted by β_Y and κ_X . This component measures the internal ordering. It governs the intrinsic scale of proper time and proper length. A Phase of 1 represents maximal internal flow (rest/vacuum), while a Phase of 0 represents the cessation of internal causality (light-speed/horizon).

$$\text{Phase} \rightarrow \text{Internal Order (Time/Structure)}$$

Theorem 5.4 (Universal Conservation of Relation). For both kinematic (S^1) and gravitational (S^2) modes, the sum of the squared Amplitude and squared Phase is invariant:

$$\underbrace{\text{Amplitude}^2}_{\text{External Interaction}} + \underbrace{\text{Phase}^2}_{\text{Internal Existence}} = 1 \quad (1)$$

Proof. By geometric nature of the relational carriers S^1 ($\beta^2 + \beta_Y^2 = 1$) and S^2 ($\kappa^2 + \kappa_X^2 = 1$), they satisfy [$\underbrace{\text{Amplitude}^2}_{\text{External Interaction}} + \underbrace{\text{Phase}^2}_{\text{Internal Existence}} = 1$] which encodes the finite relational budget. $\square$

5.1 Consequence: Relativistic Effects

Proposition 5.5 (Physical Interpretation: Relativistic Effects). The conservation law of Theorem (Universal Conservation of Relation) implies that any redistribution between the orthogonal components $\underbrace{\text{Amplitude}^2}_{\text{External Interaction}} + \underbrace{\text{Phase}^2}_{\text{Internal Existence}} = 1$ manifests physically as the relativistic effects of time dilation and length contraction.

Proof. By Theorem (Universal Conservation of Relation) the components satisfy ($\beta^2 + \beta_Y^2 = 1$); ($\kappa^2 + \kappa_X^2 = 1$). An increase in Amplitude (β, κ) enforces a decrease in Phase measure (β_Y, κ_X). This reduction of β_Y, κ_X corresponds to dilation of proper time and contraction of proper length, while the growth of β, κ represents momentum/potential. Thus the relativistic and gravitational trade-off is the direct physical expression of the geometric closure of S^1 and S^2 relational carriers. $\square$

Summary:

Geometry of spacetime is dictated by the geometry of energy relations.

Remark 5.6. This identifies Relativistic and Gravitational Time Dilation not as a mysterious "slowing down" of clocks, but as a strict geometric **phase rotation**. As a system invests more of its existence into external Amplitude (β or κ), it necessarily withdraws from its internal Phase (β_Y or κ_X), changing the rate of its proper time evolution.

6 Energy as a Relation - What κ and β Actually Mean

Energy (1.6) is the measure of differences between states.

In relational framework:

- Physical parameters like energy, speed, and gravitational potential don't belong to objects.
- Instead, they represent how we, as observers, measure objects state differences from our own point of view.

In this view, your perspective is always the reference frame. You are always at the point of origin (0,0) on your (β, κ) plane. At this origin, your internal phases are maximal ($\beta_Y = 1, \kappa_X = 1$), ensuring the conservation equations hold for the observer's self-state.

Everything else is described by how it differs from your state:

- β is the measure of how much of the universal "transformation rate" you see as motion through space, relative to yourself.
- κ is the measure of how deeply an object sits in a potential well, as seen from your position. It's your personal "ruler" for potential depth.

Think of κ and β as your own relational measuring tools:

- β is how far along your “kinetic ruler” you project another object’s state.
- κ is how deep into your “potential well” you see another object’s state.

Thus relational understanding emerges naturally:

- Energy is the capacity to move between states.
- Saying “the object’s energy” always implicitly means “the object’s energy as measured from your perspective.”

Here’s a simple analogy:

Imagine standing on a train platform. A train passes by rapidly: to you, it has significant kinetic energy. But if you jump onto the train, it instantly becomes stationary relative to you. Its kinetic energy is now zero - because your frame of reference shifted. The energy didn’t vanish; your perspective changed.

Summary:

- The projections κ and β are your personal, relational measurements of energy difference.
- All measurements boils down to describing how things differ in relation to you.

6.1 Ontological Status of the Relational Carriers S^1 and S^2

A natural question arises regarding the ontological status of the circle S^1 and the sphere S^2 : What are they, and where do they "exist"?

The answer requires a shift in perspective. In WILL Relational Geometry, S^1 and S^2 are **not spatial entities** existing within a pre-defined container. They are the necessary **relational architectures** that implement the core identity $\text{SPACETIME} \equiv \text{ENERGY}$.

Energy as Relational Transformation Capacity Recall that energy is defined as the *relational measure of difference between possible states*. It is not an intrinsic property but a relational potential for change. It is never observed directly, only through transformations.

The Carriers as Protocols of Interaction The Carriers S^1 and S^2 are the minimal, unique mathematical structures capable of hosting this relational "bookkeeping" for directional and omnidirectional transformations, respectively. They enforce closure, conservation, and symmetry by their very topology.

Imagine two observers, A and B :

- Observer A is the center of their own relational framework. Observer B is a point on A ’s S^1 (for kinematic relations) and S^2 (for gravitational relations).
- Simultaneously, observer B is the center of their own framework. Observer A is a point on B ’s S^1 and S^2 .

There is no privileged "master" carrier. Each observable interaction is structured by these mutually-centered relational protocols. The parameters β and κ are the coordinates within these relational dimensions, and the conservation laws (e.g., $\beta^2 + \beta_Y^2 = 1$; $\kappa_X^2 + \kappa^2 = 1$) are the innate accounting rules of these protocols.

7 Emergence of Spacetime

In this construction, “space,” “time,” are not treated as separate, fundamental aspects of reality. Instead, they are shown to arise as necessary consequences of a single, underlying principle: the geometry of a closed, relational system.

Therefore, the question "Where are S^1 and S^2 ?" is a category error. They are not *in* space; they are the structures whose coordinated, multi-centered interactions **give rise to** the phenomenon we perceive as spacetime. Spacetime is the emergent, collective shadow cast by the dynamics of energy relations projected onto these architectures.

In essence, S^1 and S^2 are the ontological embodiment of the relational principle. They are derived as structures that can house the transformational resource (energy) in a closed, conserved, and isotropic system. Their status is that of a fundamental **relational geometry** from which physics is generated.

8 Kinetic Energy Projection on S^1

Since S^1 encodes one-dimensional shift, the total energy E of the system must project consistently onto both axes:

$$E^2 = E_\beta^2 + E_{\beta_Y}^2$$

where

$$E_X = E\beta, \quad E_Y = E\beta_Y.$$

Desmos project

Kinetic projection β on the S^1 carrier (Special Relativity)

8.1 Geometric Nature of Rest Energy State

To derive the energy transformation, we must first explicitly identify the physical meaning of the orthogonal projections.

Theorem 8.1 (Invariant Projection of Rest Energy). *For any state (β, β_Y) on the relational circle S^1 , the total energy E must scale such that its vertical projection remains constant and equal to the rest energy E_0 .*

$$E\beta_Y = E_0.$$

Proof. Let the total energy E be distributed on the relational carrier S^1 with projections β (kinematic amplitude) and β_Y (internal phase).

$$E_Y = E \cdot \beta_Y$$

By Principle 1.3 (Relational Origin), the internal structure of an object (its rest energy E_0) is defined solely by relations internal to itself. Therefore, it must be invariant under changes in its relation to an external observer. In relation to itself kinematic projection is always $(\beta = 0 \implies \beta_Y = 1)$. Therefore:

$$E_Y \equiv E_0$$

where E_0 is the energy measured in the frame where the system is at rest ($\beta = 0, \beta_Y = 1$). Thus, geometric consistency requires the vertical leg to be fixed:

$$E\beta_Y = E_0 \implies E = \frac{E_0}{\beta_Y}.$$

The "hypotenuse" (Total Energy E) is therefore not a fixed-length vector that rotates (which would reduce E_Y), but a scalable relational magnitude that grows to preserve the invariant vertical leg E_0 against the closure constraint $\beta_Y = \sqrt{1 - \beta^2}$. $\square$

Summary:

The historical Lorentz factor γ is the reciprocal of β_Y . $\gamma = 1/\beta_Y$

8.2 The Geometric and Historical Nature of Mass

Corollary 8.2 (Rest Energy and Mass Equivalence). *This geometry does not require kilograms as units or mass as a concept. Within the normalization $c = 1$, the invariant rest energy E_0 can be expressed as human convention driven parameter of mass in kilograms:*

$$E_0 = m.$$

Proof. From the invariant projection $E\beta_Y = E_0$ and closure of S^1 , no additional scaling parameter is required. Hence the conventional bookkeeping identities $E_0 = mc^2$ or $m = E_0/c^2$ reduce to tautologies. Mass is therefore not independent, but the rest-energy invariant itself. $\square$

Summary:

Mass is the invariant projection of total rest energy expressed in anthropocentric units of kilograms.

8.3 Energy–Momentum Relation

Proposition 8.3 (Horizontal Projection as Momentum). *On the relational circle, the unique linear geometric projection measure of external amplitude from rest is the horizontal projection $E_X = E\beta$; hence*

$$p \equiv E_X \equiv E\beta \quad (c = 1).$$

Proof. The rest state is $(\beta, \beta_Y) = (0, 1)$. A shift measure must (i) vanish at rest, (ii) grow monotonically with $|\beta|$, and (iii) flip sign under $\beta \mapsto -\beta$. The only relational candidate satisfying (i)-(iii) is the horizontal projection $E_X = E\beta$. Thus the identification is necessary and unique within methodological constraints. $\square$

Desmos project

Energy-momentum Triangle

Corollary 8.4 (Energy–Momentum Relation). *With p identified by Proposition 8.3 and $m = E_0$, the closure identity yields*

$$E^2 = p^2 + m^2 \quad (c = 1).$$

Equivalently, upon restoring c ,

$$E^2 = (pc)^2 + (mc^2)^2.$$

Proof. By closure, $(E\beta)^2 + (E\beta_Y)^2 = E^2$. Substituting $p = E\beta$ and $m = E_0$ proves the claim. Restoring c is dimensional bookkeeping: $p \mapsto pc$ and $m \mapsto mc^2$, while E remains E , yielding the standard form. $\square$

Remark 8.5 (Geometric Forms). *The same identity may be expressed explicitly in terms of circle coordinates:*

$$E^2 = \left(\frac{\beta}{\beta_Y} E_0\right)^2 + E_0^2 = (\cot(\theta_1) E_0)^2 + E_0^2.$$

These are equivalent renderings of the same geometric necessity.

Remark 8.6 (Units sanity check - bookkeeping). *Using $\beta = v/c$, the identification $p \equiv E\beta$ gives*

$$pc = E \frac{v}{c} \implies p = \frac{E v}{c^2}.$$

With $E = \frac{1}{\beta_Y} mc^2 = \gamma mc^2$ this reduces to $p = \frac{\beta}{\beta_Y} mc = \gamma mv$, the standard relativistic momentum. No new parameters are introduced.

$\beta = v/c \quad \theta_1 = \arccos(\beta)$	
Algebraic Form	Trigonometric Form
$\beta = v/c = \sqrt{1 - \beta_Y^2}$	$\beta = \cos(\theta_1) = \cos(\arccos(\beta))$
$\beta_Y = \sqrt{1 - (v/c)^2} = \sqrt{1 - \beta^2}$	$\beta_Y = \sin(\theta_1) = \sin(\arccos(\beta))$

Table 1: Geometric representation of relativistic effects.

Summary

The energy–momentum relation $E^2 = (pc)^2 + (mc^2)^2$ is geometric identity of S^1 .

9 Potential Energy Projection on S^2

IMPORTANT:

Throughout this work, S^1 and S^2 are not to be interpreted as spacetime geometries but purely as relational carriers encoding energy conservation. Any reading otherwise is a misinterpretation.

Analogous to S^1 the relational geometry of the sphere, S^2 , provides orthogonal projections, for two aspects of omnidirectional transformation. We define them as follows:

- **The Amplitude Component (κ):** This projection represents the *relational gravitational measure* between the object and the observer. It corresponds to the *extent* of transformation, which manifests physically as gravitational potential. A value of $\kappa = 1$ denotes *saturation*: the entire relational resource of the system has been allocated into the gravitational channel. No residual capacity remains for kinematic projection. This condition defines the relational horizon.
- **The Phase Component (κ_X):** This projection governs the intrinsic scale of its proper length and proper time units, corresponding to the *sequence* of its transformation.

These two components are not independent but are bound by the fundamental conservation law of the closed system, which acts as a finite “budget of transformation”:

$$\kappa_X^2 + \kappa^2 = 1$$

The manifestation of any system is distributed between its internal (Phase) and relational (Amplitude) aspects. This single geometric constraint gives rise to the core phenomena of modern physics.

Desmos project

Potential projection as κ on S^2 carrier

9.1 Gravitational Meridional Section of S^2

By isotropy the omnidirectional carrier is S^2 , but any radially symmetric exchange reduces to a great-circle meridional section. We therefore work on a unit great circle of S^2 with the parametrization $(\kappa_X, \kappa) = (\cos \theta_2, \sin \theta_2)$.

9.2 Consequence: Gravitational Effects

The redistribution of the budget between the Phase and Amplitude components directly produces the effects of General Relativity. An increase in the relational measure (κ , gravitational potential) necessarily requires a decrease in the measure of the internal structure (κ_X). This geometric trade-off is observed physically as gravitational length and time corrections. Thus, the geometry of spacetime is the shadow cast by the geometry of relations.

9.3 Gravitational Tangent Formulation

Just as the relativistic energy–momentum relation can be expressed in terms of the kinematic projection $\beta = v/c$, we may construct its gravitational analogue using the potential projection $\kappa = v_e/c$, where v_e is the escape velocity at radius r .

In the kinematic case, with $\beta = \cos \theta_1$, the energy relation can be written as

$$E^2 = (\cot \theta_1 E_0)^2 + E_0^2, \quad (2)$$

so that the relativistic momentum is expressed as

$$p = E_0/c \cot \theta_1. \quad (3)$$

In full symmetry, the gravitational case follows from $\kappa = \sin \theta_2$. We define the gravitational energy as

$$E_g = \frac{E_0}{\kappa_X}, \quad \kappa_X = \sqrt{1 - \kappa^2}, \quad (4)$$

and introduce the gravitational analogue of momentum:

$$p_g = E_0/c \tan \theta_2. \quad (5)$$

This yields the gravitational energy relation

$$E_g^2 = (p_g c)^2 + (m c^2)^2. \quad (6)$$

Summary:

$$\begin{aligned} \beta &= \cos \theta_1, & \kappa &= \sin \theta_2, \\ \beta &\longleftrightarrow \kappa, & \cot \theta_1 &\longleftrightarrow \tan \theta_2. \end{aligned}$$

Kinematic momentum p and gravitational momentum p_g are thus dual projections of the same relational circle, expressed through complementary trigonometric forms.

10 Geometric composition of SR and GR factors

On the unit kinematic circle (S^1) we parametrize

$$(\beta, \beta_Y) = (\cos \theta_1, \sin \theta_1),$$

so that the invariant vertical projection reads

$$E \beta_Y = E_0 \implies \boxed{E = \frac{E_0}{\beta_Y} = \frac{E_0}{\sin \theta_1}}, \quad p = \frac{E}{c} \beta = \frac{E_0 \beta}{\beta_Y} = E_0 \cot \theta_1,$$

and therefore $E^2 = (pc)^2 + E_0^2$.


On the gravitational circle (S^2) we parametrize

$$(\kappa_X, \kappa) = (\cos \theta_2, \sin \theta_2),$$

so that the invariant horizontal projection reads

$$E_g \kappa_X = E_0 \implies \boxed{E_g = \frac{E_0}{\kappa_X} = \frac{E_0}{\cos \theta_2}}, \quad p_g = E_g \kappa = \frac{E_0 \kappa}{\kappa_X} = E_0 \tan \theta_2,$$

and therefore $E_g^2 = p_g^2 + E_0^2$.



11 Total Relational Shift Q

Desmos Project

Q as Total Relational Shift

When an observer observes another system, they assign to it a Total Relational Shift norm Q :

$$\boxed{Q^2 = \beta^2 + \kappa^2} \tag{7}$$

Relational reciprocity is the invariance of this norm under the self-centering operation of each observer.

Each observer places itself at the relational origin

$$(\beta, \kappa) = (0, 0).$$

If the other system now looks back, it again self-centres at $(0, 0)$ and applies the same rule. It measures the observer's (β, κ) and again obtains

$$Q^2 = \beta^2 + \kappa^2.$$

Thus Q is the *norm* of Total Relational Shift, not a spatial distance. Geometrically, the observer is always at the centre of its own S^1 (or S^2) carrier, and any external system is a point (β, κ) on that plane. The scalar Q measures the total deviation from the observer's relational origin.

Remark 11.1 (Closure-specific simplification). *Under energetic closure $\kappa^2 = 2\beta^2$ (circular/periodic systems), the norm reduces to $Q^2 = 3\beta^2$ (proved uniquely in Chapter, 14.2).*

In general (open or elliptic) configurations, the full definition $Q^2 = \beta^2 + \kappa^2$ must be used.

11.1 Principle of Relational Reciprocity




Figure 1: **Relational Self-Centering.** The total shift Q is defined by the orthogonal projections β and κ . Interaction is causal only when the center of the Object lies within the Observer's horizon ($Q < 1$), ensuring mutual coverage.

Self-centering reciprocity. Every observer performs self-centering:

$$(\beta, \kappa) = (0, 0).$$

When I observe another system, I assign to it (β, κ) and therefore a total shift

$$Q^2 = \beta^2 + \kappa^2.$$

When that system observes me, it again self-centres and obtains the same form. It assigns to me some (β, κ) and again computes $Q^2 = \beta^2 + \kappa^2$.

Reciprocity is therefore not a vector symmetry in a shared space. It is a symmetry of the *self-centering operation*: each observer applies the same rule and only the norm of shift is invariant.

Summary

Relational reciprocity = invariance of the norm Q under self-centering.

There is no common background arena. There are only mutual total shift magnitudes Q computed in each observer's own relational origin.

12 Clear Relational Symmetry Between Kinematic and Potential Projections

Now we can clearly see the underlying symmetry between relativistic and gravitational factors that can be expressed in unified algebraic and trigonometric forms, as shown in Table 1.

Summary


The familiar SR and GR factors emerge here as projections of the same conserved resource. Relativistic (β) and gravitational (κ) modes are not separate "effects" but dual aspects of one energy-transformation constraint revealing their unified origin.

$\theta_1 = \arccos(\beta), \quad \theta_2 = \arcsin(\kappa), \quad \kappa^2 = 2\beta^2$	
Algebraic Form	Trigonometric Form
$\beta = v/c$	$\beta = \cos(\theta_1)$
$\kappa = \sqrt{R_s/r}$	$\kappa = \sin(\theta_2)$
$\beta_Y = \sqrt{1 - \beta^2}$	$\beta_Y = \sin(\theta_1) = \sin(\arccos(\beta))$
$\kappa_X = \sqrt{1 - \kappa^2}$	$\kappa_X = \cos(\theta_2) = \cos(\arcsin(\kappa))$
$p = E_0/c \cdot \beta/\beta_Y$	$p = E_0/c \cdot \cot(\theta_1)$
$p_g = E_0/c \cdot \kappa/\kappa_X$	$p_g = E_0/c \cdot \tan(\theta_2)$
$\tau = \beta_Y \kappa_X$	$\tau = \sin(\theta_1) \cos(\theta_2)$
$\tau_Y = \sqrt{1 - \tau^2}$	$\tau_Y = \sqrt{1 - \tau^2}$
$Q = \sqrt{\kappa^2 + \beta^2} = \sqrt{3}\beta$	$Q = \sqrt{3} \cos(\theta_1)$
$Q_Y = \sqrt{1 - Q^2} = \sqrt{1 - \kappa^2 - \beta^2} = \sqrt{1 - 3\beta^2}$	$Q_Y = \sqrt{1 - 3\cos^2(\theta_1)}$

Table 2: Unified representation of relativistic and gravitational effects for closed systems.

Phenomenon	Radius r	β^2	κ^2	Q^2	Comment
ISCO (Static location)	$3R_s$	0	$\frac{1}{3}$	$\frac{1}{3}$	Pure coordinate location, $r = R_s/\kappa^2$
ISCO (Dynamic state)	$3R_s$	$\frac{1}{6}$	$\frac{1}{3}$	$\frac{1}{2}$	Marginal orbital stability, $Q^2 = Q_Y^2 = 1/2$
Photon sphere (Static)	$\frac{3}{2}R_s$	0	$\frac{2}{3}$	$\frac{2}{3}$	Pure coordinate location
Causal Dynamic horizon	$\frac{3}{2}R_s$	$\frac{1}{3}$	$\frac{2}{3}$	1	Relational horizon under closure $\kappa^2 = 2\beta^2$, $Q = 1$
Causal Static horizon	R_s	0	1	1	Purely gravitational closure limit, $\kappa^2 = 1$
Dynamic horizon	R_s	$\frac{1}{2}$	1	$\frac{3}{2}$	Orbital closure at Schwarzschild radius
Extremal Kerr horizon	$\frac{1}{2}R_s$	1	2	3	Maximal rotation, $\beta = 1$, merged horizons

Table 3: Critical radii and their projectional parameters. Static coordinate locations ($\beta = 0$) are strictly distinguished from their corresponding dynamic orbital states that emerge from the closure law $\kappa^2 = 2\beta^2$.



13 Equivalence Principle as Derived Identity

Lemma 13.1 (Unified Relational Scaling). *Within the relational framework of WILL, both kinematic (S^1) and gravitational (S^2) transformations act as independent projections of the same invariant energy E_0 . Each projection rescales the observable quantities by its respective geometric factor:*

$$E = \frac{E_0}{\beta_Y}, \quad E_g = \frac{E_0}{\kappa_X}.$$

Proof. On the kinematic circle S^1 , the invariant vertical projection corresponds to $\beta_Y = \sin \theta_1$. Preserving the same invariant leg E_0 forces the stretch $E/E_0 = 1/\beta_Y$. On the gravitational sphere S^2 , the invariant horizontal projection

is $\kappa_X = \cos \theta_2$, forcing $E_g/E_0 = 1/\kappa_X$. These transformations are independent and commute, each preserving the closure identity of its respective carrier. $\square$

Theorem 13.2 (Equivalence of Inertial and Gravitational Response). *Composing the independent relational stretches of Lemma 13.1 yields the total local energy scale*

$$E_{\text{loc}} = \frac{E_0}{\tau} = \frac{E_0}{\beta_Y \kappa_X} = \frac{E_0}{\sqrt{(1-\beta^2)(1-\kappa^2)}}.$$

The corresponding inertial and gravitational projections share a single operational factor,

$$\tilde{p} = \frac{E_{\text{loc}}}{c} \beta, \quad \tilde{p}_g = \frac{E_{\text{loc}}}{c} \kappa,$$

both governed by the same effective mass

$$m_{\text{eff}} = \frac{E_0}{\beta_Y \kappa_X c^2} = \frac{E_0}{\tau c^2}.$$

Therefore,

$$\boxed{m_g \equiv m_i \equiv m_{\text{eff}}},$$

and the Einstein equivalence principle follows as a necessary structural identity of WILL.

Corollary 13.3 (Mass Invariance under Relational Scaling). *The invariant core E_0 denotes the complete internal equilibrium state ($\beta_Y = \kappa_X = 1$). Relational factors β_Y and κ_X rescale only external manifestations (energy, momentum, and rates), while E_0 remains unchanged. Hence,*

$$\boxed{m_g \equiv m_i \equiv m = E_0/c^2},$$

is not a dynamical statement but the definition of rest invariance itself.

Remark 13.4 (Composition-Independence). *Decomposing the invariant rest energy into internal channels,*

$$E_0 = \sum_a E_0^{(a)},$$

each term couples identically through the same geometric stretch:

$$E_{\text{loc}} = \sum_a \frac{E_0^{(a)}}{\tau}.$$

Since all channels scale by the same factor $1/\tau = 1/(\beta_Y \kappa_X)$, ratios between channels cancel in all observables. Therefore, composition-independence of motion (Eotvos universality) follows identically, without requiring a postulate $m_g = m_i$.

Remark 13.5 (Quantum Interface). *The relational phase increment inherits the same scaling:*

$$\Delta\phi \propto E_{\text{loc}} \Delta\lambda,$$

where $\Delta\lambda$ is the internal ordering parameter. Thus both kinematic and gravitational phase shifts share the same stretch $1/\tau = 1/(\beta_Y \kappa_X)$, yielding composition-independent matter-wave interference patterns.

Summary:

In WILL, the equivalence of inertial and gravitational mass is not assumed but follows necessarily from the compositional closure of relational geometry. What General Relativity posits as a postulate, WILL reveals as a corollary.

14 Unification of Projections: The Geometric Exchange Rate

Having established that directional (kinematic) and omnidirectional (gravitational) relations are carried by the S^1 and S^2 respectively, we now derive the relationship that unifies them.

14.1 Derivation of the Energetic Closure Condition

Remark 14.1 (From slice to whole S^2). *Although we parametrise a single meridional great circle (κ_X, κ) for algebraic convenience, the amplitude κ^2 denotes the total omnidirectional budget of the S^2 carrier. The exchange-rate factor 2 reflects that S^2 has two independent relational degrees of freedom even when calculations are carried out on a representative great-circle section.*

14.2 Uniqueness of the Exchange Rate (No Hidden Weighting)

Lemma 14.2 (DOF-Indifference). *Under maximal symmetry (no privileged directions) and ontological minimalism (no hidden structure), any admissible conserved budget must assign equal quadratic weight to each independent relational degree of freedom.*

Proof. If two independent DOF contributed unequal weights to the conserved quadratic budget, then the theory would contain an implicit weighting structure that distinguishes DOF. This constitutes a privileged feature not derivable from relations, violating maximal symmetry and minimalism. $\square$

Theorem 14.3 (Closure). *Within WILL, the only exchange rate between the kinematic carrier S^1 (1 DOF) and the gravitational carrier S^2 (2 DOF) compatible with (i) closure in quadratic form, (ii) maximal symmetry, and (iii) ontological minimalism is:*

$$\kappa^2 = 2\beta^2.$$

Proof. Let b denote the conserved quadratic budget associated with a single independent DOF. By Lemma 14.2, each DOF must contribute the same amount b .

The carrier S^1 has 1 DOF, so its total quadratic budget is $B_{S^1} = 1 \cdot b = b$. The carrier S^2 has 2 DOF, so its total quadratic budget is $B_{S^2} = 2 \cdot b$.

An exchange rate is precisely the statement that the omnidirectional budget is a fixed multiple of the directional budget:

$$B_{S^2} = \mathcal{R} B_{S^1}.$$

Substituting $B_{S^2} = 2b$ and $B_{S^1} = b$ gives $\mathcal{R} = 2$, hence

$$\kappa^2 = \mathcal{R} \beta^2 = 2\beta^2.$$

Any $\mathcal{R} \neq 2$ would require unequal DOF weighting (hidden structure) or an extra free parameter, violating the methodology. $\square$

Remark 14.4 (Status). *The factor 2 is not empirical calibration and not a coordinate choice. It is the unique consequence of (1 DOF vs 2 DOF) under symmetry and minimalism.*

Definition 14.5 (Closure Factor).

$$\delta \equiv \frac{\kappa^2}{2\beta^2}$$

A subsystem is energetically closed if $\langle \delta \rangle_{\text{cycle}} = 1$. For circular orbits, $\delta \equiv 1$.

Corollary 14.6 (Energetic Closure Criterion). *Closed systems (momentary or periodic) satisfy $\kappa^2 = 2\beta^2$ identically. Open systems display $\delta \neq 1$, the magnitude of which quantifies the energy flow through unaccounted channels. When all channels are included, closure is restored.*

Remark 14.7 (Physical Interpretation). *The exchange rate between the kinematic and gravitational projections corresponds to the ratio of their relational dimensions. This purely geometric constant (2) replaces the empirical proportionalities of classical dynamics. It is the relational origin of the **virial theorem**: the kinetic and potential aspects of WILL maintain closure through the invariant ratio*

$$\kappa^2 = 2\beta^2.$$

Illustrative Examples.

- **Circular Orbit (Closed).** A body at any orbital phase exactly satisfies $\kappa^2 = 2\beta^2$. The entire conserved resource is partitioned between kinetic and gravitational projections; no internal "breathing" and no external channel exists.
- **Elliptical Orbit (Closed).** A body satisfies $\langle \kappa^2 \rangle = 2 \langle \beta^2 \rangle$ exactly as an average per orbital cycle due to internal "breathing" of elliptical systems. Though this internal "breathing" is restricted by the Energy-Symmetry Law (15) so the difference $W = \frac{1}{2}(\kappa^2 - \beta^2) = \text{constant}$ at any orbital phase. No external channel exists.

- **Radiating Binary (Open).** An elliptical compact binary violates $\langle \kappa^2 \rangle = 2 \langle \beta^2 \rangle$ when only orbital degrees of freedom are counted, the closure defect δ quantifying energy lost to gravitational radiation. Including all channels restores closure.

Summary:

1. WILL is a closed relational structure, $\text{SPACETIME} \equiv \text{ENERGY}$.
2. The simplest maximally symmetric carriers of these relations are S^1 and S^2 .
3. The parameters $\beta = \cos \theta_1$ and $\kappa = \sin \theta_2$ are thus constrained to these carriers.
4. The geometric exchange rate between these modes equals the ratio of their relational dimensionalities: 2.

Remark 14.8 (Geometric Origin of Physical Law). *The relation between kinetic and potential energy is not an empirical coincidence but a geometric necessity of relational closure. Classical mechanics merely approximates this deeper invariant. Explicitly,*

$$\text{Geometric Distribution } (\kappa^2) \equiv 2 \times \text{Kinetic Distribution } (\beta^2).$$

15 Energy-Symmetry Law

In RG, every transformation is bidirectional: each change observed by A corresponds to an equal and opposite change observed by B . This reciprocity is the algebraic form of causal continuity, and its geometric expression is the Energy-Symmetry Law.

15.1 Causal Continuity and Energy Symmetry

Theorem 15.1 (Energy Symmetry). *The specific energy differences (per unit of rest energy) perceived by two observers for a transition between their states balance according to the Energy-Symmetry Law:*

$$\Delta E_{A \rightarrow B} + \Delta E_{B \rightarrow A} = 0. \quad (8)$$

Proof. Consider two observers:

- Observer A at rest on the surface at radius r_A (state defined by $\kappa_A, \beta_A = 0$).
- Observer B orbiting at radius $r_B > r_A$ with orbital velocity v_B (state defined by κ_B, β_B).

Each observer perceives energy transfers as the sum of the change in potential and kinetic energy budgets.

From A 's perspective (transition from surface to orbit):

1. An object gains potential energy by moving away from the gravitational center.
2. It gains kinetic energy by accelerating to orbital velocity.

The total specific energy required for this transition is the sum of these two contributions:

$$\Delta E_{A \rightarrow B} = \underbrace{\frac{1}{2}(\kappa_A^2 - \kappa_B^2)}_{\text{Change in Potential}} + \underbrace{\frac{1}{2}(\beta_B^2 - \beta_A^2)}_{\text{Change in Kinetic}} \quad (9)$$

Since observer A is at rest, $\beta_A = 0$, and the expression simplifies to:

$$\Delta E_{A \rightarrow B} = \frac{1}{2}((\kappa_A^2 - \kappa_B^2) + \beta_B^2) \quad (10)$$

From B 's perspective (transition from orbit to surface):

1. The object loses potential energy descending into a stronger gravitational field.
2. It loses kinetic energy by reducing its velocity to rest.

This results in a specific energy difference:

$$\Delta E_{B \rightarrow A} = \frac{1}{2}((\kappa_B^2 - \kappa_A^2) + (\beta_A^2 - \beta_B^2)) = \frac{1}{2}((\kappa_B^2 - \kappa_A^2) - \beta_B^2) \quad (11)$$

Summing these transfers gives:

$$\Delta E_{A \rightarrow B} + \Delta E_{B \rightarrow A} = 0 \quad (12)$$

Thus, no net energy is created or destroyed in a closed cycle of transitions, confirming the Energy-Symmetry Law as a direct consequence of the closed geometry. $\square$

15.2 The Specific Energy Transfer (ΔE):

This is the projectional energy difference between states, equivalent to the Total Relational Shift Q , corresponding to the classical total energy of a transition (per unit rest energy). It is defined as the **sum of the changes** in the potential and kinetic energy budgets:

$$\Delta E_{A \rightarrow B} = \Delta U_{A \rightarrow B} + \Delta K_{A \rightarrow B} = \frac{1}{2}(\kappa_A^2 - \kappa_B^2) + \frac{1}{2}(\beta_B^2 - \beta_A^2) \quad (13)$$

It is this quantity, ΔE , that is conserved and must balance to zero in any closed system.

When the closure condition for stable, periodic orbits ($\kappa^2 - 2\beta^2 = 0$) is applied, the general Energy-Symmetry Law simplifies into remarkably elegant and direct forms. These simplified equations provide the precise energy balance for transitions involving energetically closed systems, such as planets or satellites in stable orbits.

Case 1: Surface-to-Orbit Transfer. For a transfer from a state of rest (A, where $\beta_A = 0$) to a closed orbit (B) where E_{0B} is the objects rest energy, the specific energy balance is given by:

$$\frac{E_{A \rightarrow B}}{E_{0B}} = \frac{1}{2}(\kappa_A^2 - \beta_B^2) \quad (14)$$

This result is derived by applying the closure condition $\kappa_B^2 = 2\beta_B^2$ to the general energy transfer formula, elegantly linking the initial potential projection to the final kinetic projection.

Case 2: Orbit-to-Orbit Transfer. For a transfer between two different closed orbits (A and B), the simplification is even more profound. The specific energy balance reduces to:

$$\frac{E_{A \rightarrow B}}{E_{0B}} = \frac{1}{2}(\beta_A^2 - \beta_B^2) \quad (15)$$

In this case, applying the closure condition to both the initial and final orbits causes the potential projection terms (κ^2) to cancel out completely. The entire energy balance of the transfer is expressed purely as the difference between the squares of the initial and final kinetic projections. This demonstrates a deep symmetry in the energetic structure of stable orbital systems.

15.3 Physical Meaning of the Factor $\frac{1}{2}$

The factor $\frac{1}{2}$ originate from the quadratic nature of the energy budgets in RG. The energetic significance of a state is proportional to the **square** of its geometric projection. This is the unavoidable consequence of relational carriers closure condition (amplitude² + phase² = 1). By using only amplitudes (β^2 and κ^2) we operating with half's relational budgets of S^1 and S^2 carriers.

The individual energy budgets are:

- **Specific Potential Energy Budget:** $U/E_0 \propto -\frac{1}{2}\kappa^2$
- **Specific Kinetic Energy Budget:** $K/E_0 = \frac{1}{2}\beta^2$

The factor $\frac{1}{2}$ arises naturally when representing a conserved quantity (energy) through a quadratic measure (the square of a projection). The Energy-Symmetry Law deals with the sum of the *changes* in these individual budgets.

15.4 Universal Speed Limit as a Consequence of Energy Symmetry

Theorem 15.2 (Universal Speed Limit). *The universal speed limit ($v \leq c$) emerges naturally from the requirement of energetic symmetry.*

Proof. Assume an object could exceed the speed of light, implying $\beta > 1$. In this scenario, its specific kinetic energy budget, $\frac{1}{2}\beta^2$, would become arbitrarily large.

The energy transfer required to reach this state, $\Delta E_{A \rightarrow B}$, would also become arbitrarily large. Consequently, no finite physical process could provide a balancing reverse transfer, $\Delta E_{B \rightarrow A}$, that would sum to zero. The fundamental symmetry would be broken:

$$\Delta E_{A \rightarrow B} + \Delta E_{B \rightarrow A} \neq 0 \quad (16)$$

Therefore, the condition $\beta \leq 1$ (which implies $v \leq c$) is an intrinsic requirement for maintaining the causal and energetic consistency of the relational universe. $\square$

15.5 Single-Axis Energy Transfer and the Nature of Light

Theorem 15.3 (Single-Axis Transformation). *For light, the kinematic projection amplitude reaches its full extent:*

$$\boxed{\beta = 1 \implies \beta_Y = 0.}$$

*This means that all transformation of the relational resource occurs along a **single X axis** $\implies \beta = 1$. The orthogonal Y axis is absent $\implies \beta_Y = 0$, and the total resource of transformation is entirely expressed on one geometric component.*

Proof. For massive bodies, the Energy-Symmetry Law (Section 15) partitions the transformation resource equally between orthogonal axes. The specific binding energy invariant for orbital motion is therefore:

$$W_{\text{mass}} = \frac{1}{2}(\kappa^2 - \beta^2), \quad (17)$$

where the factor $\frac{1}{2}$ arises from this dual-axis distribution (Theorem 5.4). This invariant is conserved for closed systems and reduces to the classical Keplerian energy under the closure condition $\kappa^2 = 2\beta^2$ (Corollary 14.6).

Now consider light. By the Single-Axis Transformation Principle (Theorem 15.3), its kinematic projection saturates the carrier:

$$\beta = 1 \implies \beta_Y = 0.$$

Geometrically, this collapses the relational architecture:

- The Y-axis (Phase component) vanishes entirely, as $\beta_Y = 0$ indicates no internal state evolution.
- No rest frame exists for self-centering (Section 11), eliminating the dual-framing that justifies the $\frac{1}{2}$ partitioning.
- The entire transformation resource concentrates on the single X-axis (Amplitude component).

Consequently, the energy invariant for photon interactions with a massive body (projection κ) must exclude the partitioning factor:

$$W_\gamma = \kappa^2 - \beta^2 = \kappa^2 - 1. \quad (18)$$

This is verified with no gravity state ($\kappa = 0$):

$$\begin{aligned} W_{\text{mass}} &= \frac{1}{2}(0^2 - 0^2) = 0, \\ W_\gamma &= 0^2 - 1^2 = -1. \end{aligned}$$

The value $W_\gamma = -1$ corresponds to the full rest energy cost of transitioning to a photon state - whereas the $\frac{1}{2}$ factor would erroneously yield $-\frac{1}{2}$, violating energy symmetry.

For a concrete test case, consider a photon interacting with a massive body where $\kappa_M = 0.4$:

$$\begin{aligned} W_{\text{mass}} &= \frac{1}{2}(0.4^2 - 1^2) = \frac{1}{2}(-0.84) = -0.42, \\ W_\gamma &= 0.4^2 - 1^2 = -0.84 = 2 \times W_{\text{mass}}. \end{aligned}$$

Thus, the gravitational effect on light is twice that on massive particles - matching the observed factor in light deflection.

The general energy potential follows directly: for light, the full geometric resource expresses unpartitioned along the single axis:

$$\Phi_\gamma = \kappa^2 c^2,$$

while massive bodies retain the partitioned form:

$$\Phi_{\text{mass}} = \frac{1}{2}\kappa^2 c^2.$$

This factor-of-2 is the geometric signature of axis count in relational space. No auxiliary approximations or background structures are introduced; the result emerges from the topological constraint $\beta_Y = 0$ applied to the closed carrier S^1 . $\square$

Summary

Energy Symmetry Law $\Delta E_{A \rightarrow B} + \Delta E_{B \rightarrow A} = 0$ - universal relational energy bookkeeping.

The Speed of Light is the boundary beyond which the energy symmetry law breaks down.

Causality is built-in feature of Relational Geometry.

Light has no rest frame. The disappearance of the Phase component (Y-axis $\beta_Y = 0$) concentrates the entire transformation resource on a single geometric component. This eliminates the $\frac{1}{2}$ partitioning factor, yielding $\boxed{\Phi_\gamma = \kappa^2 c^2}$ and explaining why light experiences exactly twice the geometric effect of massive bodies.

This explains the experimentally verified factor of 2 in gravitational lensing and Shapiro delay, traditionally requiring full General Relativity to derive.

16 Classical Keplerian Energy and Minkowski Interval as Single-Point Limits of the Relational Energy-Symmetry Law

A striking consequence of the Energy-Symmetry Law (Section 15) emerges when analysing the total specific orbital energy. Since energy in RG is defined *relationally, as the measure of difference between two states*, we naturally select these two states (e.g., the surface of the central body 'A' and the orbit 'B') as the reference points for the potential and kinetic energy budgets. Under this relational approach, the total specific orbital energy (potential + kinetic, per unit rest mass) naturally appears in a form **structurally identical to the Minkowski interval**.

16.1 Classical Result with Surface Reference

For a test body of mass m on a circular orbit of radius a about a central mass $M_{\oplus}$ (Earth in our example), classical Newtonian mechanics gives:

$$\Delta U = -\frac{GM_{\oplus}m}{a} + \frac{GM_{\oplus}m}{R_{\oplus}}, \quad (19)$$

$$K = \frac{1}{2}m\frac{GM_{\oplus}}{a}. \quad (20)$$

Adding these and dividing by the rest-energy $E_0 = mc^2$ yields the dimensionless total:

$$\frac{E_{\text{tot}}}{E_0} = \frac{GM_{\oplus}}{R_{\oplus}c^2} - \frac{1}{2}\frac{GM_{\oplus}}{ac^2}. \quad (21)$$

16.2 Projection Parameters and Minkowski-like Form

Remark 16.1. Here we explicitly define our projections using classical language to ease understanding for readers not yet familiar with WILL RG. For more explicit comparison and clearer derivations, we adopt standard Newtonian notation and define the projection parameters κ^2 and β^2 through their legacy equivalents involving G , M , and c . This is purely a pedagogical choice—the projections themselves remain dimensionless geometric quantities independent of these conventional constants.

Define the WILL projection parameters for the surface and the orbit:

$$\kappa_{\oplus}^2 \equiv \frac{2GM_{\oplus}}{R_{\oplus}c^2}, \quad (22)$$

$$\beta_{\text{orbit}}^2 \equiv \frac{GM_{\oplus}}{ac^2}. \quad (23)$$

Substituting into (16.1) gives the exact identity:

$$\frac{E_{\text{tot}}}{E_0} = \frac{1}{2}(\kappa_{\oplus}^2 - \beta_{\text{orbit}}^2). \quad (24)$$

This is already in the form of a *hyperbolic difference of squares*: if we set $x \equiv \kappa_{\oplus}$ and $y \equiv \beta_{\text{orbit}}$, then

$$\frac{E_{\text{tot}}}{E_0} = \frac{1}{2}(x^2 - y^2), \quad (25)$$

which is structurally identical to a Minkowski interval in $(1+1)$ dimensions, up to the constant factor $\frac{1}{2}$.

Sign convention. We use $U/E_0 = -\frac{1}{2}\kappa^2$ and $K/E_0 = \frac{1}{2}\beta^2$ as *budgets*. The minus sign attaches to the potential budget by convention of reference (surface vs infinity); the budgets themselves are positive quadratic measures, while transfer ΔE is the signed sum of budget *changes*.

16.3 Physical Interpretation

In classical derivations, (16.1) is just the sum $\Delta U + K$ with a particular choice of potential zero. In the RG, (16.2) emerges directly from the energy-symmetry relation:

$$\Delta E_{A \rightarrow B} = \frac{1}{2}((\kappa_A^2 - \kappa_B^2) + \beta_B^2),$$

with $(A, B) = (\text{surface}, \text{orbit})$, and is *invariantly* expressible as a difference of squared projections.

This shows that the Keplerian total energy is not an isolated Newtonian artifact but a special case of a deeper geometric structure. While RG refuse to postulate any spacetime metric in the traditional sense, the emergence of this Minkowski-like structure from purely energetic principles is a powerful indicator of the deep identity between the geometry of spacetime and the geometry of energy transformation.

Why This Matters

- In classical form, the total orbital energy per unit mass depends only on GM and a , and is independent of the test-mass m .
- In WILL form, the same fact is embedded in Energy-Symmetry difference of squared projections, with no need for separate “gravitational” and “kinetic” constructs.
- This re framing answers *why* the Keplerian combination appears: it is enforced by the underlying geometry of energy transformation.

17 Lagrangian and Hamiltonian as Single-Point Limits of the Relational Energy-Symmetry Law

The following section present philosophical and algebraic demonstration: the standard L and H arise as degenerate limits of the relational Energy-Symmetry law.

We now demonstrate that the familiar Lagrangian and Hamiltonian formalisms arise as limiting cases of the two-point relational Energy-Symmetry Law. Specifically, they emerge when the relational structure between two distinct observers A and B is collapsed into a single-point local description. This collapse preserves computational utility but reduces the ontological transparency of the underlying relational structure.

17.1 Definitions of Parameters

We consider a central mass M and a test mass m . The state of the test mass is described in polar coordinates (r, ϕ) relative to the central mass.

- r_A — reference radius associated with observer A (e.g., planetary surface).
- r_B — orbital radius of the test mass m (position of observer B).
- $v_B^2 = \dot{r}_B^2 + r_B^2 \dot{\phi}^2$ — total squared orbital speed at B.
- $\beta_B^2 = v_B^2/c^2$ — dimensionless kinematic projection at B.
- $\kappa_A^2 = 2GM/(r_A c^2)$ — dimensionless potential projection defined at A.

17.2 The Relational Lagrangian

Instead of a relational energy, we define the *clean relational Lagrangian* L_{rel} , which represents the kinetic budget at point B relative to the potential budget at point A:

$$L_{\text{rel}} = T(B) + U(A) = \frac{1}{2}m(\dot{r}_B^2 + r_B^2 \dot{\phi}^2) + \frac{GMm}{r_A}. \quad (26)$$

In dimensionless form, using the rest energy $E_0 = mc^2$, this is:

$$\frac{L_{\text{rel}}}{E_0} = \frac{1}{2}(\beta_B^2 + \kappa_A^2). \quad (27)$$

This two-point, relational form is the clean geometric statement.

17.3 First Ontological Collapse: The Newtonian Lagrangian

If one commits the first ontological violation by identifying the two distinct points, $r_A = r_B = r$, the relational structure degenerates into a local, single-point function:

$$L(r, \dot{r}, \dot{\phi}) = \frac{1}{2}m(\dot{r}^2 + r^2 \dot{\phi}^2) + \frac{GMm}{r}. \quad (28)$$

This is precisely the standard Newtonian Lagrangian. Its origin is not fundamental but arises from the collapse of the two-point relational Energy Symmetry law into a one-point formalism.

17.4 Second Ontological Collapse: The Hamiltonian

Introducing canonical momenta,

$$p_r = \frac{\partial L}{\partial \dot{r}} = m\dot{r}, \quad (29)$$

$$p_\phi = \frac{\partial L}{\partial \dot{\phi}} = mr^2\dot{\phi}, \quad (30)$$

one defines the Hamiltonian via the Legendre transformation $H = p_r\dot{r} + p_\phi\dot{\phi} - L$. This evaluates to the total energy of the collapsed system:

$$H = T + U = \frac{1}{2}m(\dot{r}^2 + r^2\dot{\phi}^2) + \frac{GMm}{r}. \quad (31)$$

17.4.1 Interpretation

In terms of the collapsed WILL projections ($\beta^2 = v^2/c^2$ and $\kappa^2 = 2GM/(rc^2)$, both strictly positive), the match to standard mechanics becomes explicit:

$$L = \frac{1}{2}mv^2 + \frac{GMm}{r} \longleftrightarrow \frac{1}{2}mc^2(\beta^2 + \kappa^2), \quad (32)$$

$$H = \frac{1}{2}mv^2 - \frac{GMm}{r} \longleftrightarrow \frac{1}{2}mc^2(\beta^2 - \kappa^2). \quad (33)$$

Here the “+” or “−” signs do not come from κ^2 itself, which is always positive, but from the ontological collapse of the two-point relational energy law into a single-point formalism. In WILL, both projections are clean and positive; in standard mechanics, the apparent sign difference arises only after this collapse.

Both the Lagrangian and Hamiltonian thus emerge from the same relational Energy-Symmetry Law under the identification $r_A = r_B = r$. The apparent sign difference between L and H is not a fundamental feature but an artifact of this single-point collapse: in the full two-point relational law, both projections β^2 and κ^2 are strictly positive.

Remark 17.1 (Mathematical Status: Groupoid vs. Group). *From a category-theoretic perspective, the relational transitions $\Delta E_{A \rightarrow B}$ form a **Groupoid** structure, where operations are defined only between specific connected states. Standard field theories (Hamiltonian/Lagrangian) rely on the collapse of this structure into a **Group** acting on a global manifold (Quotienting). Thus, the relational Energy-Symmetry Law operates at the Groupoid level, prior to the reduction into global field theories.*

Key Message

The Lagrangian and Hamiltonian arise as single-point limiting cases of the two-point relational Energy-Symmetry Law $\Delta E_{A \rightarrow B} + \Delta E_{B \rightarrow A} = 0$. They remain computationally valid within their domain of applicability. RG provides the more general two-point relational structure from which these formalisms can be systematically derived.

Legacy Dictionary (for conventional formalisms).

Within RG, all physical content is expressed purely in terms of relational projections β and κ on S^1 and S^2 . For readers accustomed to standard frameworks, the following translation rules may help:

1. *General Relativity (metric form):*

$$\kappa_X \triangleq \sqrt{-g_{tt}} \quad (\text{static spacetimes}), \quad \beta \triangleq \frac{\|u_{\text{spatial}}^\mu\|}{u^t c}.$$

2. *Canonical mechanics (Lagrangian/Hamiltonian):* Quantities such as $p_i = \partial L / \partial \dot{q}^i$ do not belong to the ontology of RG. They arise only after collapsing the two-point relational law into a one-point formalism. They are secondary derived quantities arising after the two-point relational structure is reduced to a local single-point formalism.

Here the symbol $\triangleq$ denotes not an ontological identity, but a pragmatic dictionary entry for translation into legacy notation.

17.5 Third Ontological Collapse: Derivation of Newton's Third Law

We now demonstrate that Newton's Third Law, like the Lagrangian and Hamiltonian, emerges as a limiting case of RG. It arises as a necessary mathematical consequence of the same ontological collapse — forcing a two-point relational law into a single-point, instantaneous formalism.

Theorem 17.2 (Newton's Third Law as a Single-Point Limit). *The Energy-Symmetry Law ($\Delta E_{A \rightarrow B} + \Delta E_{B \rightarrow A} = 0$) mathematically necessitates Newton's Third Law ($\vec{F}_{AB} = -\vec{F}_{BA}$) in the classical, non-relativistic limit where the two-point relational energy budget is collapsed into a single-point potential function $U(\vec{r})$.*

Proof. We begin with the foundational Energy-Symmetry Law (Section 15), the principle of causal balance for state transitions:

$$\Delta E_{A \rightarrow B} + \Delta E_{B \rightarrow A} = 0.$$

In the classical, non-relativistic limit, this two-point relational law is "ontologically collapsed" into a single-point potential energy function, U . This function is assumed to depend only on the relative positions of the two interacting entities, A and B :

$$U = U(\vec{r}) \quad \text{where} \quad \vec{r} = \vec{r}_B - \vec{r}_A.$$

This $U(\vec{r})$ is the classical approximation of the system's relational energy budget. In this collapsed formalism, the force $\vec{F}$ is defined as the negative gradient of this potential.

(1) Force on B by A ($\vec{F}_{AB}$): This force is found by taking the gradient with respect to B 's coordinates:

$$\vec{F}_{AB} = -\nabla_B U(\vec{r}_B - \vec{r}_A) \tag{34}$$

$$= -\left(\frac{dU}{d\vec{r}}\right) \cdot \left(\frac{\partial \vec{r}}{\partial \vec{r}_B}\right) \tag{35}$$

$$= -\nabla U(\vec{r}) \cdot (\mathbf{I}) \tag{36}$$

$$= -\nabla U(\vec{r}) \tag{37}$$

(2) Force on A by B ($\vec{F}_{BA}$): This force is found by taking the gradient with respect to A 's coordinates:

$$\vec{F}_{BA} = -\nabla_A U(\vec{r}_B - \vec{r}_A) \tag{38}$$

$$= -\left(\frac{dU}{d\vec{r}}\right) \cdot \left(\frac{\partial \vec{r}}{\partial \vec{r}_A}\right) \tag{39}$$

$$= -\nabla U(\vec{r}) \cdot (-\mathbf{I}) \tag{40}$$

$$= +\nabla U(\vec{r}) \tag{41}$$

(3) Conclusion: By direct comparison of the results, we find:

$$\vec{F}_{AB} = -\nabla U(\vec{r}) \quad \text{and} \quad \vec{F}_{BA} = +\nabla U(\vec{r}).$$

Therefore, within the single-point collapsed formalism:

$$\boxed{\vec{F}_{AB} = -\vec{F}_{BA}}$$

This completes the proof. Newton's Third Law is not an independent physical axiom but a necessary mathematical consequence of representing the two-point Energy-Symmetry Law through a single potential function $U(\vec{r})$. The principle of equal and opposite forces thus emerges as the single-point limit of the more general relational balance condition $\Delta E_{A \rightarrow B} + \Delta E_{B \rightarrow A} = 0$. $\square$


Philosophical Consequences

[Substantialism vs. Relationalism](#)

18 Relational Orbital Mechanics (R.O.M.)

In this section we will prove independence from mass M and gravitational constant G , and will show how all observational phenomenon of orbital systems are generated as the unavoidable consequences of Relational Geometry.

But first lets remind our selves some key definitions established in prior sections:



$\theta_1 = \arccos(\beta), \quad \theta_2 = \arcsin(\kappa), \quad \kappa^2 = 2\beta^2$	
Algebraic Form	Trigonometric Form
$\beta = v/c$	$\beta = \cos(\theta_1)$
$\kappa = \sqrt{R_s/r}$	$\kappa = \sin(\theta_2)$
$\beta_Y = \sqrt{1 - \beta^2}$	$\beta_Y = \sin(\theta_1) = \sin(\arccos(\beta))$
$\kappa_X = \sqrt{1 - \kappa^2}$	$\kappa_X = \cos(\theta_2) = \cos(\arcsin(\kappa))$
$p = E_0/c \cdot \beta/\beta_Y$	$p = E_0/c \cdot \cot(\theta_1)$
$p_g = E_0/c \cdot \kappa/\kappa_X$	$p_g = E_0/c \cdot \tan(\theta_2)$
$\tau = \beta_Y \kappa_X$	$\tau = \sin(\theta_1) \cos(\theta_2)$
$Q = \sqrt{\kappa^2 + \beta^2} = \sqrt{3}\beta$	$Q = \sqrt{3} \cos(\theta_1)$

Table 4: Unified representation of relativistic and gravitational effects for closed systems.

19 Operational Independence and the Role of Constants

In this and following sections we demonstrate that RG's relational projections (κ, β) are ontologically and operationally independent from mass M and gravitational constant G . We will show that R_s as a system scale derived directly from the phase interactions of light.

19.1 Derivation of the Spectroscopic Shifts

Theorem 19.1 (Spectroscopic Phase Shift). *The geometric projection κ^2 of a closed system is strictly determined by the operationally measurable gravitational redshift z_κ via the identity:*

$$\kappa^2 = 1 - \frac{1}{(1 + z_\kappa)^2}. \quad (42)$$

Proof. By the S^2 closure invariant (Theorem 5.4), the gravitational amplitude κ and internal phase κ_X satisfy:

$$\kappa^2 + \kappa_X^2 = 1 \implies \kappa^2 = 1 - \kappa_X^2. \quad (43)$$

By the Principle of Relational Origin (1.3), an observer must self-center at the relational origin $(\beta, \kappa) = (0, 0)$. Consequently, the observer's local phase is maximal: $\kappa_{X(obs)} \equiv 1$.

The observable spectroscopic shift z_κ is the operational measure of the discrepancy between the observer's internal phase and the source's internal phase. The optical scaling ratio is defined as:

$$1 + z_\kappa = \frac{\kappa_{X(obs)}}{\kappa_{X(source)}}. \quad (44)$$

Substituting the observer's maximal phase ($\kappa_{X(obs)} = 1$), the exact phase of the source is isolated:

$$\kappa_{X(source)} = \frac{1}{1 + z_\kappa}. \quad (45)$$

Substituting this internal phase back into the initial closure invariant yields the exact algebraic identity:

$$\kappa^2 = 1 - \left(\frac{1}{1 + z_\kappa} \right)^2. \quad (46)$$

□

Theorem 19.2 (Kinematic Phase Shift (Transverse Doppler)). *The kinematic geometric projection β^2 of a closed system is strictly determined by the operationally measurable transverse Doppler shift z_β via the symmetric identity:*

$$\beta^2 = 1 - \frac{1}{(1 + z_\beta)^2}. \quad (47)$$

Proof. By the S^1 closure invariant (Theorem 5.4), the kinematic amplitude β and internal kinematic phase β_Y satisfy:

$$\beta^2 + \beta_Y^2 = 1 \implies \beta^2 = 1 - \beta_Y^2. \quad (48)$$

Following the Principle of Relational Origin (1.3), the self-centered observer dictates the rest state, where external kinematic projection is zero ($\beta_{obs} = 0$), yielding maximum internal kinematic phase: $\beta_{Y(obs)} \equiv 1$.

The transverse Doppler shift z_β (measured orthogonally to eliminate classical radial kinematics) is the pure operational measure of the discrepancy between the observer's internal kinematic phase and the source's phase. The optical scaling ratio is:

$$1 + z_\beta = \frac{\beta_{Y(obs)}}{\beta_{Y(source)}}. \quad (49)$$

Substituting $\beta_{Y(obs)} = 1$, we isolate the internal phase of the moving source:

$$\beta_{Y(source)} = \frac{1}{1 + z_\beta}. \quad (50)$$

Substituting this back into the S^1 closure invariant yields the exact algebraic identity:

$$\beta^2 = 1 - \left(\frac{1}{1 + z_\beta} \right)^2. \quad (51)$$

□

19.2 Operational Measurability of Relational Projections

Theorem 19.3 (Operational Measurability). *The relational projections are encoded directly in the combined phase interactions of light (spectroscopy) and are operationally independent of G and M .*

Proof. Step 1: The Raw Observable (τ). Spectroscopy does not measure "gravitational potential" or "kinematic velocity" as separate isolates; it measures the total accumulated phase difference between the source and the observer. We define the **Relational Spacetime Factor** τ as the inverse of the systems measured redshift product:

$$\boxed{\kappa^2 = 1 - \frac{1}{(1 + z_\kappa)^2}} \quad (z_\kappa = \text{gravitational redshift})$$

$$\boxed{\beta^2 = 1 - \frac{1}{(1 + z_\beta)^2}} \quad (z_\beta = \text{transverse Doppler shift})$$

$$\tau \equiv \frac{1}{Z_{sys}} = 1/[(1 + z_\kappa)(1 + z_\beta)]. \quad (52)$$

In the WILL framework, this single observable represents the product of the internal phase projections of the carriers S^2 and S^1 :

$$\tau = \underbrace{\kappa_X}_{\text{Gravitational Phase}} \cdot \underbrace{\beta_Y}_{\text{Kinematic Phase}} = \sqrt{1 - \kappa^2} \sqrt{1 - \beta^2}. \quad (53)$$

Step 2: Exact Relational Identity. We reject weak-field approximations. The exact relationship between the signal τ and the structural norm Q is given by the algebraic expansion of the phase product:

$$\tau^2 = (1 - \kappa^2)(1 - \beta^2) = 1 - (\kappa^2 + \beta^2) + \kappa^2\beta^2. \quad (54)$$

Substituting $Q^2 = \kappa^2 + \beta^2$, we obtain the rigorous link between the optical observable and the geometric state:

$$\tau^2 = 1 - Q^2 + \kappa^2\beta^2. \quad (55)$$

This identity demonstrates that the optical signal contains the complete information about the system's structural state, measurable without prior knowledge of mass. □

Orbital dynamics boils down to interplay of frequency shifts. No mass, no G , no manifold, no metric required.

$$\kappa^2 = 1 - \frac{1}{(1 + z_\kappa)^2} \quad z_\kappa = \text{gravitational redshift}$$

$$\beta^2 = 1 - \frac{1}{(1 + z_\beta)^2} \quad z_\beta = \text{transverse Doppler shift}$$

Observational Z Inputs

$Z_{sys} = (1 + z_\kappa)(1 + z_\beta) = \frac{1}{\tau}$ (product of gravitational red shift and transverse Doppler shift)

$\tau = \kappa_X \cdot \beta_Y = \frac{1}{Z_{sys}}$ (product of projectinal phase factors on S^1 and S^2 carriers)

$z_\kappa = \frac{1}{\kappa_X} - 1$ (gravitational redshift)

$z_\beta = \frac{1}{\beta_Y} - 1$ (transverse Doppler shift)

Everything else is Closed Algebraic System of Relational Orbital Mechanics (R.O.M.) Please follow this link to see all [R.O.M. equations and definitions](#)

19.3 Derivation of Eccentricity

Eccentricity is a measure of the projectional deviation from the circular equilibrium state ($\delta = 1$).

Theorem 19.4 (Geometric Eccentricity). *For a closed orbital system governed by the projection invariants of WILL Relational Geometry, the orbital eccentricity e is strictly determined by the closure factor at periapsis, δ_p :*

$$e = \frac{2\beta_p^2}{\kappa_p^2} - 1 = \frac{1}{\delta_p} - 1. \quad (56)$$

Proof. Instead of relying on classical force laws, we derive this relation directly from the conservation of the two fundamental projection invariants of the WILL framework:

1. **Energy Projection Invariant (Binding Energy):** $W = \frac{1}{2}(\kappa^2 - \beta^2) = \text{const.}$
2. **Angular Projection Invariant:** $h = r\beta = \text{const}$ (at turning points).

Consider the two turning points of a closed orbit: periapsis (p) and apoapsis (a). By operational definition of the shape parameter e , the relation between radii is determined by the geometric range:

$$r_a = r_p \left(\frac{1+e}{1-e} \right). \quad (57)$$

Step 1: Relational Mapping. Using the angular invariant h (implying $\beta \propto 1/r$) and the field definition $\kappa^2 \propto 1/r$, we express the apoapsis projections in terms of the periapsis values:

$$\beta_a^2 = [\beta_p \left(\frac{r_p}{r_a} \right)]^2 = \beta_p^2 \left(\frac{1-e}{1+e} \right)^2, \quad (58)$$

$$\kappa_a^2 = \kappa_p^2 \left(\frac{r_p}{r_a} \right) = \kappa_p^2 \left(\frac{1-e}{1+e} \right). \quad (59)$$

Note: Kinematic projection scales quadratically with the radius ratio, while potential projection scales linearly.

Step 2: Energy Balance. Substituting these into the energy invariant conservation condition $W_p = W_a$:

$$\frac{1}{2}(\kappa_p^2 - \beta_p^2) = \frac{1}{2}(\kappa_a^2 - \beta_a^2).$$

Cancelling the factor $\frac{1}{2}$ and substituting the mappings from Step 1:

$$\kappa_p^2 - \beta_p^2 = \kappa_p^2 \left(\frac{1-e}{1+e} \right) - \beta_p^2 \left(\frac{1-e}{1+e} \right)^2.$$

Rearranging to group potential terms (κ) on the left and kinematic terms (β) on the right:

$$\kappa_p^2 \left[1 - \frac{1-e}{1+e} \right] = \beta_p^2 \left[1 - \left(\frac{1-e}{1+e} \right)^2 \right].$$

Step 3: Algebraic Reduction. Expanding the terms in brackets:

$$\begin{aligned} \text{LHS bracket } (\kappa \text{ term}): \quad & 1 - \frac{1-e}{1+e} = \frac{(1+e) - (1-e)}{1+e} = \frac{2e}{1+e}. \\ \text{RHS bracket } (\beta \text{ term}): \quad & 1 - \frac{(1-e)^2}{(1+e)^2} = \frac{(1+e)^2 - (1-e)^2}{(1+e)^2} = \frac{4e}{(1+e)^2}. \end{aligned}$$

Substituting back into the balance equation:

$$\kappa_p^2 \left(\frac{2e}{1+e} \right) = \beta_p^2 \left(\frac{4e}{(1+e)^2} \right).$$

Dividing both sides by $2e$ and multiplying by $(1+e)^2$:

$$\kappa_p^2 (1+e) = 2\beta_p^2.$$

This yields geometric identity for bound orbits:

$$2\beta_p^2 = \kappa_p^2 (1+e). \quad (60)$$

Step 4: Connection to Closure. Recall the definition of the closure factor at periapsis:

$$\delta_p = \frac{\kappa_p^2}{2\beta_p^2}.$$

Substituting Eq.(19.3) into this definition:

$$\delta_p = \frac{\kappa_p^2}{\kappa_p^2 (1+e)} = \frac{1}{1+e}.$$

Solving for e , we obtain the stated result:

$$e = \frac{1}{\delta_p} - 1 = \frac{2\beta_p^2}{\kappa_p^2} - 1 = 1 - \frac{2\beta_a^2}{\kappa_a^2}$$

□

co_speechnotes_pluginfabid : 1339,363

Remark 19.5. This result confirms that eccentricity is strictly a measure of the energetic deviation from the circular equilibrium state ($\delta = 1$), derived entirely from the conservation of relational projections without invoking mass or Newtonian forces.

SUMMARY

$$\frac{2\beta_p^2}{\kappa_p^2} - 1 \equiv e \equiv 1 - \frac{2\beta_a^2}{\kappa_a^2}$$

ECCENTRICITY $\equiv$ CLOSURE DEFECT

SPACETIME $\equiv$ ENERGY

19.4 Orbital Precession as Relational Phase Accumulation

In standard General Relativity, orbital precession is derived via perturbative methods and Taylor expansions of the Schwarzschild metric. In WILL Relational Geometry, precession is derived strictly from the exact algebraic accumulation of the internal phase difference over a closed geometric cycle, without differential approximations.

Theorem 19.6 (Exact Precession Law). The secular angular shift (precession) $\Delta\varphi$ per closed orbit is the strict geometric projection of the system's exact relational state difference τ_Y normalized by the orbital shape factor $(1-e^2)$.

Proof. **Step 1: Exact State Difference.** A system completely at rest with the observer possesses maximal internal phase $\tau = 1$. The true relational phase of an orbiting system is the exact product of the S^1 and S^2 phase projections:

$$\tau^2 = \beta_Y^2 \kappa_X^2 = (1 - \beta^2)(1 - \kappa^2). \quad (61)$$

Expanding this product algebraically yields:

$$\tau^2 = 1 - (\beta^2 + \kappa^2) + \kappa^2\beta^2. \quad (62)$$

Recognizing the total relational shift norm $Q^2 = \beta^2 + \kappa^2$, the exact absolute state divergence from the rest origin ($\tau = 1$) is defined as:

$$\tau_Y^2 \equiv 1 - \tau^2 = Q^2 - \kappa^2\beta^2. \quad (63)$$

The cross-coupling term $\kappa^2\beta^2$ represents the exact, non-linear geometric interaction between the directional and omnidirectional relational carriers.

Step 2: Geometric Accumulation (The Exact Form). The system accumulates this state mismatch over the full cycle (2π). The total angular shift is the base phase scaled by the state divergence, normalized by the elliptical geometry factor ($1 - e^2$).

We do not truncate or simplify this connection. Restricted by the Mathematical Transparency principle (??), the orbital precession is locked in its exact, fully non-linear algebraic form:

$$\Delta\varphi = \frac{2\pi \cdot \tau_Y^2}{1 - e^2} = \frac{2\pi(1 - \tau^2)}{1 - e^2} \quad (64)$$

□

Remark 19.7 (Epistemic Purity). *This derivation replaces the complex integration of perturbed differential equations with an exact algebraic evaluation of the relational phase. The persistent presence of the cross-coupling term $\kappa^2\beta^2$ confirms that RG is structurally complete and relies on exact invariant topologies rather than weak-field truncations.*

Legacy Translation: Recovering the General Relativity Approximation

To demonstrate structural compatibility with historical models, we can intentionally ablate our exact formula to see what happens when the pure geometry is truncated.

If we drop the higher-order geometric cross-coupling term ($\kappa^2\beta^2 \rightarrow 0$), we isolate the primary operational observable Q^2 . This yields the simplified precession law:

$$\Delta\varphi \simeq \frac{2\pi Q^2}{1 - e^2}. \quad (65)$$

Substituting Q^2 , we recover the standard classical form purely algebraically. By selecting the semi-major axis a as the reference scale, the closure condition ($\kappa^2 = 2\beta^2$) defines the specific distribution of the invariant Schwarzschild scale R_s :

$$\kappa^2(a) = \frac{R_s}{a}, \quad \beta^2(a) = \frac{R_s}{2a}.$$

Substituting these into the definition of the relational shift norm $Q^2 = \beta^2 + \kappa^2$ yields $Q^2 = \frac{3R_s}{2a}$. Inserting this back into our ablated equation perfectly recovers the standard GR differential approximation:

$$\Delta\varphi \simeq \frac{2\pi}{1 - e^2} \left(\frac{3R_s}{2a} \right) \equiv \frac{3\pi R_s}{a(1 - e^2)}. \quad (66)$$

Furthermore, mapping this simplified shift strictly to periapsis (p) observables using the geometric identity $(1 + e) = 2\beta_p^2/\kappa_p^2$ yields the direct operational ratio:

$$\Delta\varphi \simeq \frac{3\pi}{2} \frac{\kappa_p^4}{\beta_p^2}. \quad (67)$$

This confirms that the classical tensor calculus results, which cost years of study to master, are merely truncated shadows of the exact relational phase geometry.

19.5 Kinematic Mechanism of the Dynamic Horizon

In classical literature, the event horizon is often heuristically treated as an escape-velocity boundary where spacetime curvature becomes infinitely steep. In RG, the horizon emerges strictly as the limit of geometric precessional invariant, representing the ultimate topological closure of a trajectory.

Using the exact relational phase accumulation law (derived in 19.4), the baseline precession per orbit is given by:

$$\Delta\varphi = \frac{2\pi \cdot \tau_Y^2}{1 - e^2} = \frac{2\pi(1 - \tau^2)}{1 - e^2} \quad (68)$$

For a massive system under total energetic closure ($\kappa^2 = 2\beta^2$), consider the **Dynamic Horizon** state where the potential projection reaches absolute saturation ($\kappa^2 = 1$). By the closure rule, the kinematic projection is $\beta^2 = \frac{1}{2}$.

At this limit, the gravitational phase component collapses ($\kappa_X = \sqrt{1-1} = 0$), causing the entire internal causal structure of the system to vanish:

$$\tau^2 = (1 - \beta^2)(1 - \kappa^2) = 0 \implies \tau_Y = 1 \quad (69)$$

For a perfectly circular orbit ($e = 0$) at this causal limit, the precession evaluates exactly to:

$$\Delta\varphi = \frac{2\pi \cdot (1)}{1 - 0} = 2\pi \quad (70)$$

Geometric Origin of the Horizon

An orbital precession of exactly $\Delta\varphi = 2\pi$ radians per revolution implies that the argument of periapsis shifts by a full circle during a single orbit. Consequently, the trajectory continuously folds onto itself perfectly, creating a closed, self-intersecting locus where forward orbital motion mathematically nullifies itself. This is the true kinematic definition of the Dynamic Horizon: it is not a "point of no return" in an external container, but the unitary topological closure of the relational trajectory itself.

19.6 Algebraic Determination of Absolute System Scale

We now derive algebraic formulas for the system scale R_s using three distinct observational methods. These derivations rely strictly on the **Conservation of the Energy Invariant W** , replacing the need for Newtonian force laws.

Closed Algebraical System of R.O.M. Equations

R.O.M. equations and definitions

19.6.1 Method A: Differential (Two-Point Method)

This method is suitable when the orbital period is unknown, but the trajectory can be traced geometrically.

Theorem 19.8 (Two-Point Schwarzschild Scale). Given measurements of geometric position (r) and kinematic intensity (β) at two arbitrary points along a trajectory:

- Radii: r_1, r_2 (from astrometry)
- Intensities: β_1, β_2 (from de-projected spectral line widths or proper motion)

the Schwarzschild radius is:

$$R_s = \frac{r_1 r_2}{r_2 - r_1} (\beta_1^2 - \beta_2^2). \quad (71)$$

Proof. We invoke the **Conservation of the Relational Invariant $W = \frac{1}{2}(\kappa^2 - \beta^2)$** . This law states that for any closed system, the specific energy difference between the potential and kinematic projections is constant throughout the orbit.

Therefore, at any two points 1 and 2:

$$\frac{1}{2}(\kappa_1^2 - \beta_1^2) = \frac{1}{2}(\kappa_2^2 - \beta_2^2). \quad (72)$$

Rearranging to group the projections:

$$\kappa_1^2 - \kappa_2^2 = \beta_1^2 - \beta_2^2. \quad (73)$$

Substituting the field identity $\kappa^2 = R_s/r$:

$$\frac{R_s}{r_1} - \frac{R_s}{r_2} = \beta_1^2 - \beta_2^2. \quad (74)$$

Factoring out the scale parameter R_s :

$$R_s \left(\frac{r_2 - r_1}{r_1 r_2} \right) = \beta_1^2 - \beta_2^2. \quad (75)$$

Solving for R_s :

$$R_s = \frac{\beta_1^2 - \beta_2^2}{\frac{r_2 - r_1}{r_1 r_2}} = \frac{r_1 r_2}{r_2 - r_1} (\beta_1^2 - \beta_2^2). \quad (76)$$

This formula extracts the "mass" scale purely from geometric gradients. $\square$

19.6.2 Method B: Geometric Resonance (Balance Point Method)

At the specific orbital phase $O_o = \arccos(-e)$, the system passes through its geometric balance point where the instantaneous radius r equals the semi-major axis a . At this unique phase, the closure condition $\kappa^2 = 2\beta^2$ is satisfied instantaneously.

Theorem 19.9 (Balance Point Formula). *Given the geometric scale a and the total light signal τ measured at the balance point ($r = a$):*

$$R_s = \frac{a}{2}(3 - \sqrt{1 + 8\tau^2(O_o)}). \quad (77)$$

Proof. Step 1: Spacetime Factor Expansion. At the balance point ($r = a$), the closure condition implies $\kappa^2 = R_s/a$ and $\beta^2 = R_s/2a$. The observable τ is:

$$\tau^2 = (1 - \kappa^2)(1 - \beta^2) = (1 - \frac{R_s}{a})(1 - \frac{R_s}{2a}). \quad (78)$$

Step 2: Exact Geometric Solution. Expanding the product:

$$\tau^2 = 1 - \frac{R_s}{2a} - \frac{R_s}{a} + \frac{R_s^2}{2a^2} = 1 - \frac{3R_s}{2a} + \frac{R_s^2}{2a^2}. \quad (79)$$

Multiplying by $2a^2$ to clear denominators and rearranging into standard quadratic form ($Ax^2 + Bx + C = 0$):

$$R_s^2 - 3aR_s + 2a^2(1 - \tau^2) = 0. \quad (80)$$

Solving for R_s using the quadratic formula ($x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a_{cof}}$):

$$R_s = \frac{3a \pm \sqrt{(3a)^2 - 4(1)(2a^2(1 - \tau^2))}}{2}. \quad (81)$$

Simplifying the term under the radical:

$$9a^2 - 8a^2(1 - \tau^2) = 9a^2 - 8a^2 + 8a^2\tau^2 = a^2(1 + 8\tau^2). \quad (82)$$

Thus:

$$R_s = \frac{3a \pm a\sqrt{1 + 8\tau^2}}{2} = \frac{a}{2}(3 \pm \sqrt{1 + 8\tau^2}). \quad (83)$$

For stable orbits, we must select the negative root. □

Remark 19.10 (Light Separation). *This unit-less factor in brackets is:*

- **Potential projection at semi-major axis:** $\kappa^2 = \frac{1}{2}(3 - \sqrt{1 + 8\tau^2(O_o)})$

A unique way to separate gravitational part from the light signal.

19.6.3 Method C: Instantaneous (Arbitrary Phase Method)

Most generally, if the orbital geometry (a) is known, the scale R_s can be derived from a **single epoch** observation at any arbitrary radius r_o .

Theorem 19.11 (Arbitrary Phase Formula). *Given the orbital geometry (a, r_o) and the single light observable τ_{W_o} :*

$$R_s = \frac{r_o}{2(2a - r_o)}(4a - r_o - \sqrt{(4a - r_o)^2 - 8a(2a - r_o)(1 - \tau_{W_o}^2)}). \quad (84)$$

Proof. Step 1: Relational Invariant Form. We start with the energy invariant relation $W = \frac{1}{2}(\kappa^2 - \beta^2)$. For a bound orbit, $W = R_s/4a$. We express the kinematic intensity β^2 at arbitrary radius r in terms of the field intensity κ^2 :

$$\frac{1}{2}(\kappa^2 - \beta^2) = \frac{R_s}{4a} \implies \beta^2 = \kappa^2 - \frac{R_s}{2a}. \quad (85)$$

Substituting $\kappa^2 = R_s/r$:

$$\beta^2 = R_s(\frac{1}{r} - \frac{1}{2a}). \quad (86)$$

Step 2: Constraint via Observables. We substitute the expressions for κ^2 and β^2 into the exact observable constraint $\tau_{Wo}^2 = (1 - \kappa^2)(1 - \beta^2)$:

$$\tau_{Wo}^2 = (1 - \frac{R_s}{r})(1 - R_s[\frac{1}{r} - \frac{1}{2a}]). \quad (87)$$

Let $A = \frac{1}{r}$ and $B = \frac{1}{r} - \frac{1}{2a}$. The equation becomes:

$$\tau_{Wo}^2 = (1 - AR_s)(1 - BR_s) = 1 - (A + B)R_s + ABR_s^2. \quad (88)$$

Rearranging into quadratic form:

$$(AB)R_s^2 - (A + B)R_s + (1 - \tau_{Wo}^2) = 0. \quad (89)$$

Step 3: Coefficient Expansion. We compute the coefficients explicitly:

$$AB = \frac{1}{r} \left(\frac{2a - r}{2ar} \right) = \frac{2a - r}{2ar^2} \quad (90)$$

$$A + B = \frac{1}{r} + \frac{2a - r}{2ar} = \frac{2a + 2a - r}{2ar} = \frac{4a - r}{2ar} \quad (91)$$

Multiplying the entire quadratic equation by $2ar^2$ to clear denominators:

$$(2a - r)R_s^2 - r(4a - r)R_s + 2ar^2(1 - \tau_{Wo}^2) = 0. \quad (92)$$

Step 4: Solution. Solving for R_s :

$$R_s = \frac{r(4a - r) \pm \sqrt{r^2(4a - r)^2 - 4(2a - r)(2ar^2)(1 - \tau_{Wo}^2)}}{2(2a - r)}. \quad (93)$$

Factoring out r^2 from the radical term $\sqrt{r^2(\dots)} = r\sqrt{(\dots)}$:

$$R_s = \frac{r}{2(2a - r)}((4a - r) \pm \sqrt{(4a - r)^2 - 8a(2a - r)(1 - \tau_{Wo}^2)}). \quad (94)$$

For stable orbits, we select the negative root, yielding the exact algebraic link between the observed light phase and the system's geometric scale. $\square$

19.7 The Role of G as Translation Constant

The presence of R_s in these formulas does not imply a dependency on mass M or the constant G . The objection that $R_s = 2GM/c^2$ makes G fundamental rests on a categorical error: it mistakes a unit conversion factor for a physical source.

Theorem 19.12 (Constants as Converters). *In WILL RG, G and M are derived calibration tools used to translate geometric scales into legacy units.*

Proof. The operational procedure is strictly geometric:

1. **Measure:** Light phase τ (dimensionless) via spectroscopy.
2. **Measure:** Geometric scale r (meters/AU) via astrometry.
3. **Calculate:** System Scale $R_s = f(r, \tau)$ via Theorems above.

The physical calculation ends here. The system is fully defined. If, and only if, one wishes to interface with legacy catalogues, one employs the **unit converter**:

$$M \equiv \frac{R_s c^2}{2G}. \quad (95)$$

The constant G describes the units we use (kilograms vs. meters), not the physics of the system. $\square$

Remark 19.13 (Historical Artifact). *The kilogram is a human convention. In WILL RG, the fundamental quantity is the dimensionless ratio $\kappa^2 = R_s/r$, which encodes the energy density ratio $\rho/\rho_{\max}$. The "mass" M is a secondary bookkeeping device.*

19.8 Summary

We have derived three algebraically distinct formulas for R_s , each optimized for different observational scenarios:

1. **Two-point** Requires two velocity measurements
2. **Balance point** Simplified form at special phase
3. **Arbitrary phase** Works for any single epoch

All formulas are:

- **Operationally independent** of G and M
- **Non-circular** (inputs are direct observables)
- **Algebraically exact** (no approximations beyond Keplerian closure)

The operational input is always $\tau = 1/[(1 + z_\kappa)(1 + z_\beta)]$, a dimensionless quantity directly measurable from spectroscopy. This demonstrates that WILL RG formulas are empirically grounded and do not rely on hidden assumptions about gravitational constants.

20 Relational Parameterization and the Fundamental Primitives

Theorem 20.1 (Relational Parameterization). *The complete structural and dynamical parameterization of any closed orbital system - specifically its geometric shape (e = eccentricity), secular phase shift ($\Delta\varphi$ = orbital precession), critical scale (R_s = Schwarzschild radius), global scale (a = semi-major axis), strictly determined by the algebraic relations of its dimensionless spectroscopic and chronometric observables. The derivation requires no independent mass parameter (M), gravitational constant (G), or an a priori spatial metric.*

Proof. The proof is constructive and algebraically closed. It generalises to any bound orbital system. We demonstrate this using the Mercury-Sun system as an operational test case. All governing equations are derived exclusively from the internal topology of the WILL relational carriers (S^1, S^2) and have zero theoretical dependence on Newtonian mechanics, Special Relativity, or General Relativity. The derivation is executed using only four dimensionless, epistemologically direct observables: ($e, \theta_\odot, T_M/T_\oplus, z_{sun}$):

INPUTS (Cross-Cultural Invariants)

The following inputs are strictly dimensionless ratios. They represent direct physical observables that remain identical for any observer in the Universe, regardless of their local system of units (meters/seconds) or theoretical paradigms.

Parameter	Value	Direct Observational Method
$\theta_\odot$	0.0046524 rad	Visual Size: The angular radius of the Sun in the sky (how many angular units it occupies from the observer's viewpoint).
$T_M/T_\oplus$	0.2408	Cycle Ratio: The observed system's full orbital period divided by our own local orbital period.
e_{obs}	0.2056	Visual Kinematics: Derived directly from the ratio of the planet's fastest (ω_{max}) and slowest (ω_{min}) apparent angular speeds across the sky: $e = \frac{\sqrt{\omega_{max}/\omega_{min}-1}}{\sqrt{\omega_{max}/\omega_{min}+1}}$.
z_{sun}	2.1224×10^{-6}	Spectroscopic Tension: The raw fractional frequency shift of light ($1 - \nu_{obs}/\nu_{emit}$). Methodologically, it is defined strictly as an optical geometric observable, explicitly bypassing any a priori theoretical dependence on mass (M) or Newtonian constants (G).

Algebraic Derivation

1. Relational Scale Factor

The ratio of the Sun's radius to the planet's perihelion distance, derived entirely from visual size and cycle ratios, without metric distances:

$$R_{ratio} = \frac{\theta_{\odot}}{(T_M/T_{\oplus})^{2/3} \cdot (1 - e_{obs})} \approx 0.0151290$$

2. Potential Projection (at perihelion)

Translating the spectral shift at the Sun's surface (z_{sun}) to the potential projection at Mercury's perihelion (κ_p^2) using the derived scale factor (R_{ratio}):

$$\kappa_p^2 = \left(1 - \frac{1}{(1 + z_{sun})^2}\right) \cdot R_{ratio} \approx 6.4219 \times 10^{-8}$$

3. Kinematic Projection (at perihelion)

Deriving the orbital kinematics strictly from the potential and eccentricity, requiring no Doppler or radar data:

$$\beta_p^2 = \frac{\kappa_p^2}{2}(1 + e_{obs}) \approx 3.8712 \times 10^{-8}$$

4. Global Orbital Invariants

Deriving the system's global projections (at the semi-major axis balance point) via the invariant binding energy $W = \frac{1}{2}(\kappa_p^2 - \beta_p^2)$:

$$\beta^2 = 2W \approx 2.5508 \times 10^{-8} \quad \text{and} \quad \kappa^2 = 4W \approx 5.1016 \times 10^{-8}$$

5. Exact Orbital Precession

The secular phase shift emerges from accumulation of the relational state divergence ($\tau_Y^2 = 1 - (1 - \kappa^2)(1 - \beta^2)$) over the full cycle:

$$\Delta\varphi = \frac{2\pi \cdot \tau_Y^2}{1 - e_{obs}^2} \approx 5.0203 \times 10^{-7} \text{ rad/orbit}$$

Converting this pure geometric phase shift to standard observational units (arcseconds per century) yields exactly **42.99 arcsec/century**, matching the observed precession of Mercury within observational uncertainties.

6. Absolute System Scale (Metric Translation)

To translate this into human units (meters), we express Mercury's orbital period in seconds ($T_M = 7,600,521.6$ s) and the speed of light ($c = 299,792,458$ m/s) to the derived global kinematic projection ($\beta = \sqrt{2W}$):

$$R_{sRG} = T_M \cdot c \cdot \frac{\beta^3}{\pi} \approx 2954.8 \text{ m}$$

and the standard value is:

$$R_{sGR} = \frac{2GM_{sun}}{c^2} \approx 2953.33 \text{ m} \quad \text{where} \quad M_{sun} = 1.98847 \times 10^{30} \text{ kg}$$

Comparing the values:

$$\frac{R_{sGR} - R_{sRG}}{R_{sGR}} \times 100 = |0.0476|\%$$

This difference is well within the observational uncertainties of the input parameters.

7. Semi-major axis (Standard value $\approx 5.79 \times 10^{10}$ m):

$$a = \frac{R_{sRG}}{\kappa^2} = 5.792288 \times 10^{10} \text{ m}$$

8. Orbital radius at perihelion (Standard value $\approx 4.60 \times 10^{10}$ m):

$$r_p = \frac{R_{sRG}}{\kappa_p^2} = 4.601394 \times 10^{10} \text{ m}$$

9. Conclusion: The logical chain is complete and algebraically closed. All major parameters of gravitational system recovered without use of mass, gravitational constant G, metric manifold, differential formalism, postulation of "container-space" and external flow of time. $\square$

No differential formalism. No metric space. Just light frequency shift red vs. blue

Corollary 20.2 (Epistemic Mandate and Ontological Redundancy). *In information theory and formal logic, if a parameter is strictly absent from the complete algebraic generative chain of a system, its reintroduction constitutes an epistemic violation. Because the full structural and dynamical parameterization $(e, \Delta\varphi, R_s)$ is algebraically closed using only directly measurable parameters $(e, \theta_\odot, T_M/T_\oplus, z_{sun})$ and derived relational projections (κ, β) , the variables G and M possess zero independent predictive power. They are not fundamental primitives. Their retention required only for conversion of pure relational geometry into legacy units of kilograms.*

Mercury-Sun Relational Parameterization

DESMOS COLAB

21 Rotational Systems (Kerr Without Metric)

21.1 Contextual Bounds

- **For a gravitationally closed (static) system**, the physical boundary is defined by the condition $\kappa^2 = 1$. The closure principle ($\kappa^2 = 2\beta^2$) is what dictates that this corresponds to a kinetic state of $\beta^2 = 1/2$.
- **For a kinematically closed (maximally rotating) system**, the physical boundary is defined by the condition $\beta^2 = 1$. The same closure principle ($\kappa^2 = 2\beta^2$) then necessitates that the corresponding gravitational state must be $\kappa^2 = 2$.

For rotating black holes, we establish the connection between relational kinetic projection and the Kerr metric by defining:

$$\beta = \frac{ac^2}{Gm_0}, \quad \kappa = \sqrt{2}\beta$$

where:

- β is the relational rotation parameter, with $0 \leq \beta \leq 1$,
- κ is related to the geometry and gravity,
- $R_s = \frac{2Gm_0}{c^2}$ is the Schwarzschild radius,
- $a = \frac{J}{m_0c}$ is the Kerr rotation parameter,
- J is the angular momentum of the black hole,
- m_0 is the mass of the black hole.

We also derive a key invariant relationship:

$$a_{\max} = \frac{Gm_0}{c^2} = \frac{R_s}{2} = \beta_{\max}^2 r$$

This relationship holds when $r = \frac{R_s}{2\beta^2}$, providing an elegant connection between the parameters.

21.2 Event Horizon

Using our approach, the inner and outer event horizons of the Kerr metric are expressed as:

$$r_{\pm} = \frac{R_s}{2} (1 \pm \beta_Y)$$

For the extreme case where $\beta = 1$ (maximal rotation), the horizons merge at:

$$r_+ = r_- = \frac{R_s}{2}$$

This coincides with the minimum radius in our model predicted using maximum value of κ parameter $\kappa_{\max} = \sqrt{2}$:

$$r_{\min} = \frac{1}{\kappa_{\max}^2} R_s = \frac{1}{2} R_s$$

21.3 Ergosphere

The radius of the ergosphere in our model is described as:

$$r_{\text{ergo}} = \frac{R_s}{2} \left(1 + \sqrt{1 - \beta^2 \cos^2 \theta} \right)$$

This formulation correctly reproduces the key features of the ergosphere:

- At the equator ($\theta = \pi/2$), $r_{\text{ergo}} = R_s$ for any rotation parameter,
- At the poles ($\theta = 0$), r_{ergo} coincides with the event horizon radius.

21.4 Ring Singularity

Unlike the Schwarzschild metric with its point singularity, the Kerr metric features a ring singularity located at:

$$r = 0, \quad \theta = \frac{\pi}{2}$$

The "size" of this ring is proportional to $a = \frac{Gm_0}{c^2} \beta$, reaching its maximum for extreme black holes ($\beta = 1$).

21.5 Naked Singularity

For $\beta \leq 1$, a naked singularity does not emerge, aligning with the cosmic censorship Principle. In our model, Energy Symmetry Law enforce constraint by limiting β to the range $[0, 1]$.

21.6 Chiral Asymmetry: The Algebraic Origin of Frame-Dragging

In standard General Relativity, the chiral asymmetry of a rotating black hole (where prograde and retrograde orbits diverge) is modeled via the off-diagonal metric tensor component $g_{t\phi}$. In the Relational Orbital Mechanics (R.O.M.), this asymmetry is not a geometric deformation of a background manifold, but a strict algebraic consequence of linear phase superposition on the directed S^1 carrier.

21.6.1 Composite Kinematic Projection

When a test particle (or photon) with orbital kinematic projection β_{orb} interacts with a central body possessing a spin projection β_{spin} , the total relational kinematic shift must be aggregated linearly on the S^1 carrier before the quadratic energy invariant is calculated.

The composite kinematic projection β_{Σ} is defined as:

$$\beta_{\Sigma} = \beta_{\text{orb}} \pm \beta_{\text{spin}} \quad (96)$$

where the (+) sign denotes co-rotating (prograde) relational motion, and the (−) sign denotes counter-rotating (retrograde) relational motion.

21.6.2 The Algebraic Symmetry Breaker (Cross-Term)

Substituting the composite projection into the fundamental relational energy invariant $W = \frac{1}{2}(\kappa^2 - \beta_{\Sigma}^2)$, we obtain the expanded state balance:

$$W = \frac{1}{2} [\kappa^2 - (\beta_{\text{orb}} \pm \beta_{\text{spin}})^2] \quad (97)$$

$$W = \frac{1}{2} [\kappa^2 - (\beta_{\text{orb}}^2 \pm 2\beta_{\text{orb}}\beta_{\text{spin}} + \beta_{\text{spin}}^2)] \quad (98)$$

The emergent cross-term $\pm 2\beta_{\text{orb}}\beta_{\text{spin}}$ is the exact algebraic equivalent of the frame-dragging effect. It enforces a chiral split in the required gravitational projection κ^2 to maintain the invariant W :

- **Prograde Orbits (+):** The cross-term increases the total subtracted kinetic budget, requiring a deeper gravitational projection (larger κ^2 , hence smaller r) to maintain closure.
- **Retrograde Orbits (−):** The cross-term decreases the total kinetic budget, requiring a shallower gravitational projection (smaller κ^2 , hence larger r).

21.6.3 Orbital and Optical Splitting

Because all dynamic orbital parameters in R.O.M. (such as precession $\Delta\varphi \propto \tau_Y^2$ and deflection angles) are strictly proportional to the ratio of projections, replacing the static β_{orb} with the chiral β_Σ automatically generates the asymmetric splitting of trajectories.

For a photon ($\beta_{orb} = 1$) grazing a rotating mass, the relative kinematic budget becomes $\beta_\Sigma = 1 \pm \beta_{spin}$. This directly splits the photon sphere radii algebraically, eliminating the need to integrate geodesics over a Kerr metric. The spatial "dragging" is entirely replaced by the interference of relational phase projections.

21.7 The Relationship Between $\kappa > 1$ and Rotation

For extreme rotation ($\beta = 1$), we find $\kappa = \sqrt{2} > 1$, which reflects the displacement of the event horizon and the geometric properties of rotating black holes. This suggests that values of $\kappa > 1$ are inherently connected to the physics of relational rotation.

This connection suggests that the rotation of a black hole can be understood through geometric parameters analogous to orbital mechanics. Physically, it indicates that the rotational properties of the black hole, encapsulated in a_* , mirror the orbital velocity parameter β , providing a unified description of spacetime dynamics.

Physical Interpretation

- **No need for pre-existing spacetime** - geometry emerges from energy distributions.
- **All parameters** are dimensionless and directly derived from the speed of light as finite resource.
- **Scale invariance:** The same structure applies from Planck-scale objects to galactic black holes.

22 Time and Phase Evolution

In strictly empirical terms, RG refrains from ontological speculation regarding an absolute, continuous background flow of time. Instead, time is strictly operationalized through measurement: it is the physically recorded accumulation of cyclic oscillations from a local reference clock relative to the geometric phase evolution of the observed system.

22.1 Continuous Model and Closed-Form Evaluation

The time duration of a given phase interval is defined via the integral:

$$\Delta_{to}(o) = \int_0^o \frac{1}{\omega_\theta(\theta)} d\theta = \frac{a}{\beta c} \tau_o = \frac{T}{2\pi} \tau_o \quad (99)$$

where $\omega_o(o)$ is the instantaneous angular frequency at phase o :

$$\omega_o(o) = \frac{\beta \cdot c}{a} \cdot \frac{(1 + e \cdot \cos(o))^2}{(1 - e^2)^{\frac{3}{2}}} \quad (100)$$

and τ_o is the temporal phase interval:

$$\tau_o = (1 - e^2)^{\frac{3}{2}} \int_0^o (1 + e \cos \theta)^{-2} d\theta \quad (101)$$

This integral admits an exact closed-form evaluation expressed entirely in R.O.M. quantities. Via the Weierstrass half-angle substitution $t = \tan(\theta/2)$ and subsequent reduction through the shape factor $e_X = (1 + e)/(1 - e)$ and the complementary eccentricity $e_Y = \sqrt{1 - e^2}$, the temporal phase interval evaluates to:

$$\tau_o(o) = 2 \arctan\left(\frac{\sin(o/2)}{\sqrt{e_X} \cos(o/2)}\right) - \frac{e e_Y \sin(o)}{1 + e \cos(o)} \quad (102)$$

Every symbol in this expression is native to R.O.M.:

- o — orbital phase (true anomaly),
- $e = 1/\delta_p - 1$ — eccentricity (closure defect),
- $e_X = (1 + e)/(1 - e) = r_a/r_p = \beta_p/\beta_a = \kappa_p^2/\kappa_a^2$ — shape factor,
- $e_Y = \sqrt{1 - e^2}$ — complementary eccentricity.

The forward direction (phase $\rightarrow$ time) is therefore algebraically closed:

$$\Delta_{to}(o) = \frac{T}{2\pi} \tau_o(o) \quad (103)$$

22.2 Irreducible Transcendence of the Inverse Problem

The inverse problem — determining the orbital phase o from a given elapsed time Δt — requires solving:

$$\tau_o(o) = \frac{2\pi}{T} \Delta t \quad (104)$$

Since the unknown o appears simultaneously inside the arctan and the trigonometric terms, this equation is irreducibly transcendental: no finite algebraic rearrangement to the form $o = f(\Delta t, e)$ exists. This is a mathematical property of elliptical geometry, not a limitation of any formalism.

Graphically, the equation is solved by plotting $\tau_o(x)$ against the constant right-hand side and reading the intersection directly.

22.3 Operational Discrete Alternative

By strictly adhering to the operational definition of time as discrete, quantized measurements (ticks of a reference clock), the orbital trajectory is generated via a discrete iterative map that replaces the transcendental inversion with an algebraic recurrence at the cost of first-order discretization error.

For a given discrete reference clock interval Δt_{tick} (e.g., one standard second), the first-order discrete phase shift Δo_n executed by the system is dictated directly by the local angular frequency:

$$\Delta o_n \approx \omega_o(o_n) \cdot \Delta t_{\text{tick}} \quad (105)$$

The complete orbital phase evolution over any measured macroscopic time interval Δt is computationally recovered by iterating the discrete map:

$$o_{n+1} = o_n + \left(\frac{\beta \cdot c}{a} \cdot \frac{(1 + e \cdot \cos(o_n))^2}{(1 - e^2)^{\frac{3}{2}}} \right) \Delta t_{\text{tick}} \quad (106)$$

Remark 22.1 (Operational Discretization). This iterative state machine represents the operational reality of the physical measurement apparatus. The accumulated discretization error within this first-order map physically corresponds to the intra-tick geometric evolution that remains fundamentally unresolvable to an observer bounded by the finite frequency of their local clock. The continuous, transcendental integral is recovered identically only in the unobservable limit $\Delta t_{\text{tick}} \rightarrow 0$.

23 Strong Field Test (S2 Star)

For distant stars like S2 (orbiting Sgr A*), we **reconstruct** the potential depth κ_p purely from the visible orbital shape (e) and velocity (β_p), using the relation derived from Eq.(19.3):

$$\kappa_p = \beta_p \sqrt{\frac{2}{1 + e}}.$$

Data (GRAVITY Collaboration):

$$e \simeq 0.8846, \quad \beta_p \simeq 0.0255 \text{ (7650 km/s)}.$$

Prediction: 1. Reconstruct Potential: $\kappa_p \approx 0.02627$. 2. Calculate Precession Ratio using Eq. (??):

$$\frac{\kappa_p^4}{\beta_p^2} = \frac{(0.02627)^4}{(0.0255)^2} \approx 7.32 \times 10^{-4}.$$

3. Result:

$$\Delta\varphi = \frac{3\pi}{2} (7.32 \times 10^{-4}) \approx 3.45 \times 10^{-3} \text{ rad} \approx \mathbf{11.85'}.$$

Comparison:

- **WILL prediction:** $\approx 11.89'$.
- **Observed shift:** $12' \pm 1.5'$.

The result lies well within observational uncertainty.

Red vs. Blue

This demonstrates that orbital geometry is encoded in the ratio of gravitational redshift to Doppler shift, revealing that mass is not a fundamental input but a derived bookkeeping quantity

24 Blind Prediction for S4716 (In Silico Experiment)

Blind Prediction Protocol

*This section documents a specific numerical prediction made in November 2025, prior to the observational confirmation of the relativistic precession for the star S4716. By de-projecting the raw line-of-sight velocity from 2009 SINFONI data, we reconstruct the relativistic state vector and predict a precession of **14.80 arcmin/orbit**. The fact that this result, usually requiring the full machinery of General Relativity, can be derived via algebraic closure suggests that strong-field gravity may be fully described by the algebra of optical observables.*

24.1 Operational Derivation from Observation

To derive the orbital precession of S4716 without relying on the mass of Sgr A* or the Schwarzschild metric, we utilize a scale-invariant reconstruction based on the geometric shape of the orbit (e) and the kinetic intensity (β) derived from optical observables.

While the Decryption Invariant (Section 25.6) allows autonomous extraction of the inclination i given full phase coverage at the extrema, historical data often provides only sparse or single-epoch measurements. In such cases, the full spectroscopic curve is unavailable.

However, the R.O.M. framework is algebraically modular. By adopting the astrometric inclination to resolve the single-epoch projection, the internal closure algebra autonomously generates the relativistic precession without any reliance on mass models or the Schwarzschild metric.

24.2 Geometric De-projection of Single-Epoch Velocity

The observational input is the Line-of-Sight (LOS) velocity measured by SINFONI in 2009 ($v_{LOS} \approx 1690$ km/s) [?]. Because the dataset lacks the extrema required to deploy Equation 124, we input the independently measured inclination ($i \approx 161.13^\circ$) to de-project the total velocity.

The geometric projection factor $\mathcal{P}$ at true anomaly o is:

$$\mathcal{P}(o) = \left| \sin(i) \frac{\cos(\omega + o) + e \cos(\omega)}{\sqrt{1 + e^2 + 2e \cos(o)}} \right|. \quad (107)$$

Using the orbital parameters for S4716 ($e = 0.756$, $i = 161.13^\circ$, $\omega = 2.25^\circ$) and the calculated phase for 2009 ($o \approx 1.122$ rad), we find $\mathcal{P} \approx 0.25$. The total velocity is:

$$v_{total} = \frac{v_{LOS}}{\mathcal{P}} \approx \frac{1690}{0.25} \approx 6760 \text{ km/s}. \quad (108)$$

This yields the scale-invariant intensity parameter at phase o :

$$\beta_o = \frac{v_{total}}{c} \approx 0.02255. \quad (109)$$

24.3 Propagation to Periapsis

We propagate this state to periapsis using the geometric invariants of WILL.

1. Closure Factor at Observation:

$$\delta_o = \frac{1 + e \cos(o)}{1 + e^2 + 2e \cos(o)} \approx 0.38317314. \quad (110)$$

2. Potential Projection: From the closure definition, the potential depth κ_o at that phase is:

$$\kappa_o = \beta_o \cdot \sqrt{\delta_o \cdot 2} \approx 0.02462222. \quad (111)$$

3. The Energy Invariant (W): The orbital energy invariant W is constant throughout the trajectory:

$$W = \frac{1}{2}(\kappa_o^2 - \beta_o^2) \approx 0.00004889967. \quad (112)$$

4. **Solution at Periapsis:** At periapsis ($o = 0$), the closure factor simplifies to $\delta_p = (1 + e)^{-1}$. Solving for the periapsis velocity β_p via invariance of W :

$$\beta_p = \sqrt{\frac{W}{\delta_p - 0.5}} \approx 0.02652. \quad (113)$$

This yields $\beta_p \approx 0.0265$ ($v_p \approx 7956$ km/s), derived independent of mass models.

24.4 Prediction

With the derived periapsis intensity β_p and the corresponding potential $\kappa_p = \beta_p \sqrt{\frac{2}{1+e}} \approx 0.02831$ we apply the Precession Law (Eq. ??):

$$\Delta\varphi = \frac{3\pi}{2} \frac{\kappa_p^4}{\beta_p^2} \approx \frac{3\pi}{2} (9.13 \times 10^{-4}). \quad (114)$$

$$\Delta\varphi \approx 14.7909706972 \approx 14.80 \text{ arcmin/orbit.} \quad (115)$$

Summary

Orbital phenomena require no mass, G , metrics, manifolds, no deferential field equations or any additional assumptions. All observable phenomena arise entirely from algebraic relations among observable frequency projections.

25 Sidestepping the Classical $M \sin i$ Degeneracy

25.1 The 130-Year Limitation

For more than a century, radial-velocity and astrometric orbital analyses have been fundamentally limited by the $M \sin i$ (or equivalently $\beta \sin i$) degeneracy. The observed velocity semi-amplitude K only constrains the product

$$K = \frac{\beta_{\text{int}} \sin i}{\sqrt{1-e^2}},$$

leaving an infinite family of physically distinct (β, i) pairs consistent with the data. This degeneracy has been the principal obstacle to model-independent mass and inclination determination in exoplanet, binary, and Galactic-center studies.

By deriving Relational Orbital Mechanics (R.O.M. 18) RG sidesteps this degeneracy at the algebraic level by introducing two independent higher-order kinematic invariants that depend on β (or β_{int}) alone, without reference to the observer's inclination i .

25.2 The Interior Kinematic Projection

We define the **interior kinematic projection** (the projection that appears in all relativistic invariants) as

$$\beta_{\text{int}} \equiv \frac{\beta}{\sqrt{1-e^2}}.$$

This is the natural variable once the elliptical geometry is accounted for. All higher-order effects now depend strictly on powers of β_{int} .

25.3 Non-Linear Relational Invariants

R.O.M. supplies two independent constraints that are quadratic (or higher) in β_{int} :

Theorem 25.1 (Decoupling of β and i). *The classical linear amplitude $K \propto \beta_{\text{int}} \sin i$ is supplemented by two non-linear invariants:*

1. **Precessional Phase Shift** (from total relational shift accumulation):

$$\Delta\phi = 6\pi\beta_{\text{int}}^2.$$

2. **Systemic Transverse Baseline** $Z_{\text{sys}}(\phi)$ (phase-dependent modulation arising from the transverse components of the kinematic and potential projections).

Both invariants depend only on β_{int}^2 (and orbital phase), independent of the observer's line-of-sight inclination i .

Proof. The precession law follows directly from the accumulation of the total relational shift $Q^2 = 3\beta^2$ over one closed orbit. Substituting $\beta = \beta_{\text{int}} \sqrt{1-e^2}$ immediately yields $\Delta\phi = 6\pi\beta_{\text{int}}^2$.

The Z_{sys} term arises from the exact product of the phase projections on the carriers S^1 and S^2 and is likewise quadratic in the interior kinematic amplitude. Neither invariant carries a factor of $\sin i$. Therefore, once these two

higher-order signals are included, β_{int} is fixed independently of the linear amplitude K . The inclination i is then uniquely recovered from

$$\sin i = \frac{K\sqrt{1-e^2}}{\beta_{\text{int}}}.$$

□

Operational Summary

The degeneracy is not fundamental; it is an artifact of truncating the kinematic description to its first-order linear term. R.O.M. restores the full non-linear structure of the relational carriers and renders β and i separately measurable from a single dataset.

25.4 Empirical Validation: Bayesian MCMC Analysis on S0-2 Data

To strictly test the resolution quantitatively, a full 5-dimensional Bayesian Markov Chain Monte Carlo (MCMC) analysis was performed on the 82 radial-velocity measurements of the S0-2 star. The parameter space was mapped across a 5D posterior: $p(\beta, i, v_{z0}, e, \omega_0 | \text{data})$.

An affine-invariant ensemble sampler (emcee) was utilized with 50 walkers to compare two models:

- **Full R.O.M.:** Includes both Z_{sys} and precession (β_{int}^2 invariants).
- **Ablated model:** Classical Keplerian RV only (no Z_{sys} , no precession).

25.4.1 Ablation Study Results

Table 5 demonstrates that the classical (ablated) RV model fundamentally fails to capture the true physical state of the system, while the Full R.O.M. successfully encompasses the independently verified GRAVITY Collaboration parameters within its 95% confidence interval (CI).

Parameter	Full R.O.M. (Median $\pm 1\sigma$)	Ablated (Median)	GRAVITY Ref.	In 95% CI?
β (Kinematic)	$0.00597^{+0.00108}_{-0.00078}$	0.00477	0.00645	YES
i (Inclination)	$130.8^{+9.6}_{-11.7}^\circ$	105.8°	134.0°	YES
v_{z0} (Drift)	$-18.2^{+4.5}_{-6.5}$ km/s	-2.2 km/s	-20.0 km/s	YES
e (Eccentricity)	0.88501 ± 0.00038	0.88496	0.88466	YES
ω_0 (Periaapsis)	$65.99^{+0.63}_{-0.39}^\circ$	64.43°	66.13°	YES

Table 5: MCMC marginal posteriors for S0-2. The Ablated model fails to capture the true parameters for β , i , and v_{z0} , confirming that classical RV analysis is strictly degenerate without geometric invariants.

Detailed Analysis

https://willrg.com/msini_test

25.4.2 Topology of the Posterior and the Degeneracy Ridge

While R.O.M. successfully localizes the physical truth, the MCMC mapping reveals that the $\beta - i$ degeneracy is not reduced to a singular point constraint, but rather narrowed to a well-defined topological ridge.

Model	$\text{corr}(\beta, i)$	β width	i width
Full R.O.M.	+0.971	31%	21.3°
Ablated (classical)	+0.811	12%	29.3°
Full R.O.M. + v_{z0} prior	+0.978	20%	12.8°

Table 6: Quantitative degeneracy metrics. The Full R.O.M. correctly positions the degeneracy ridge, while the ablated model displaces it entirely into non-physical parameter space. Column widths refer to 68% credible intervals.

Colab Notebook

[Beta i comparison.ipynb](#)

Because the invariant constraints (precession and Z_{sys}) both scale strictly as $\sim \beta^2$, they manifest as phase-differentiated signals of the same geometric order. This overdetermines the system, effectively moving the probability ridge to intersect the physically true values, but preserves a narrow 1σ distribution width due to the instrumental noise floor.




Figure 2: Direct comparison of the β - i marginal posteriors. Left: Full R.O.M. (blue) — the degeneracy ridge is correctly centred on the true values. Right: Ablated model (orange) — the ridge is displaced and the true values lie outside the 95% contour.

25.4.3 Independent Validation via Background Drift (v_{z0})

The behaviour of the background drift v_{z0} serves as an independent diagnostic. In the Full model, v_{z0} accurately converges to -18.2 ± 5.5 km/s, exhibiting a strong physical anti-correlation with the kinematic projection ($r(\beta, v_{z0}) = -0.851$). Higher theoretical values of β generate a stronger Z_{sys} baseline shift, forcing the optimizer to correctly adjust the background velocity.

In the Ablated model, this anti-correlation vanishes ($r = +0.031$), and v_{z0} collapses to an absurd -2.2 km/s. This proves that Z_{sys} is an active, physically requisite invariant in the orbital data.




Figure 3: Full 5D Posterior mapping for S0-2 demonstrating the correctly positioned probability ridge.

25.5 Independent Validation via Synthetic 1PN Data

To rigorously eliminate observational bias and prove that the algebraic decoupling of β and i is not an artefact of the specific S0-2 dataset or instrumental noise profile, the R.O.M. framework was subjected to a strict blind in silico experiment.

25.5.1 Data Generation and Observational Strategy

A synthetic radial velocity dataset was generated using a standard 1st Post-Newtonian (1PN) Keplerian engine. The true parameters were randomly seeded, simulating a highly eccentric test mass ($e > 0.9$) orbiting a supermassive black hole.

Crucially, the generative model utilized the classical additive approximations for relativistic effects:

$$v_{\text{rel_shift}} = c \left(\frac{GM}{rc^2} + \frac{v_{\text{orb}}^2}{2c^2} \right), \quad (116)$$

and injected a realistic Gaussian noise profile of $\sigma = 3.0$ km/s. To simulate a realistic observational campaign (e.g., the GRAVITY Collaboration), the epoch sampling included sparse background monitoring and dense sampling around two consecutive periapsis passages to capture the accumulated precessional shift.

25.5.2 Three-Stage Algebraic Extraction

The recovery algorithm operated blindly, utilizing only the exact geometric constraints of R.O.M. without any reference to M or G . The optimization was driven entirely by the unified phase observable:

$$Z_{\text{sys}} = \frac{1}{\sqrt{1 - \beta_o^2} \sqrt{1 - \kappa_o^2}}, \quad (117)$$

coupled with the continuous geometric precession rate $3\beta_{\text{int}}^2$ per radian of true anomaly.

To avoid topological traps, a physics-first, three-stage extraction architecture was employed:

1. **Global Keplerian Scout:** An unconstrained differential evolution scan isolated the geometric timing (Period P , eccentricity e , and epoch T_{peri}).
2. **R.O.M. Global Sniper:** The algorithm isolated the macroscopic amplitude $K \propto \beta \sin i$, breaking the i vs $180^\circ - i$ projection symmetry by folding the domain to $[0, \pi/2]$.
3. **MCMC Bayesian Mapping:** A 32-walker ensemble mapped the residual probability ridge to extract the final parameters.

25.5.3 Blind Test Results

The R.O.M. framework successfully isolated the physical truth, breaking the $\beta \sin i$ degeneracy purely through the structural imposition of the S^1 and S^2 relational carrier constraints.

Colab Notebook

[ROM Validation via Synthetic 1PN Data.ipynb](#)

Parameter	True (Hidden 1PN Data)	R.O.M. MCMC Median	Deviation
Period (yrs)	12.237	12.237	0.00%
Eccentricity (e)	0.91024	0.91023	< 0.01%
Inclination (i folded)	49.85°	49.92°	0.07°
Background Drift (v_{z0})	−1.15 km/s	−1.00 km/s	0.15 km/s

Table 7: Strict blind test recovery results ($\sigma = 3.0$ km/s). The R.O.M. algorithm cleanly separates the inclination i from the kinematic projection β without prior mass constraints.

Methodological Confirmation

The extraction of the true inclination angle i from 1D radial synthetic data confirms that the $\beta - i$ degeneracy is strictly broken by the non-linear mathematical structure of the R.O.M. invariants. The framework explicitly maps 1D observables back to their complete 3D generative geometry.

25.6 Analytical Extraction of the Inclination Angle: The Decryption Invariant

While Bayesian MCMC provides a robust statistical mapping of the probability ridge (Section 25.4), the Relational Orbital Mechanics (R.O.M.) allows for a strict, closed-form algebraic resolution of the $M \sin i$ degeneracy. By evaluating the raw observational shifts at the orbital extrema, the inclination angle i can be completely isolated and extracted.

R.O.M. equations and definitions

Closed Algebraic System of Relational Orbital Mechanics

25.6.1 The Observer Equations at Extrema

The raw spectroscopic shift observed by a telescope, $Z_{\text{raw}}(o)$, is the product of the line-of-sight Doppler projection and the systemic transverse baseline $Z_{\text{sys}}(o)$. The extrema of the radial velocity occur at phases $o_{i1} = -\omega_i$ (maximum redshift) and $o_{i2} = \pi - \omega_i$ (minimum redshift/maximum blueshift).

Utilizing the relational energy invariant $\kappa_o(o)^2 - \beta_o(o)^2 = \beta^2$, we eliminate the gravitational projection κ_o from Z_{sys} entirely. The systemic baselines at the extrema, denoted D_{max} and D_{min} , become strict functions of the global kinematic projection β , eccentricity e , and the argument of periaapsis ω_i :

$$D_{\text{max}}(\beta, e, \omega_i) = \sqrt{1 - 2\beta^2 \frac{1 + e \cos \omega_i}{1 - e^2}} \sqrt{1 - \beta^2 \frac{1 + e^2 + 2e \cos \omega_i}{1 - e^2}} \quad (118)$$

$$D_{\text{min}}(\beta, e, \omega_i) = \sqrt{1 - 2\beta^2 \frac{1 - e \cos \omega_i}{1 - e^2}} \sqrt{1 - \beta^2 \frac{1 + e^2 - 2e \cos \omega_i}{1 - e^2}} \quad (119)$$

The observed raw extrema can then be written linearly with respect to the classical semi-amplitude invariant $K_i = \frac{\beta \sin i}{\sqrt{1 - e^2}}$:

$$Z_{\text{rawmax}} \cdot D_{\text{max}} = 1 + K_i(1 + e \cos \omega_i) \quad (120)$$

$$Z_{\text{rawmin}} \cdot D_{\text{min}} = 1 - K_i(1 - e \cos \omega_i) \quad (121)$$

25.6.2 Algebraic Decoupling of β and i

To break the degeneracy, we isolate the inclination-dependent term. Subtracting Equation 121 from Equation 120 yields exactly twice the semi-amplitude:

$$2K_i = Z_{\text{rawmax}}D_{\text{max}} - Z_{\text{rawmin}}D_{\text{min}} \quad (122)$$

Adding the two equations yields:

$$Z_{\text{rawmax}}D_{\text{max}} + Z_{\text{rawmin}}D_{\text{min}} = 2 + 2K_i e \cos \omega_i \quad (123)$$

Substituting $2K_i$ from Equation 122 into this sum and grouping the Z terms isolates the fundamental **Decryption Invariant**:

The Decryption Invariant

$$Z_{\text{rawmax}}D_{\text{max}}(1 - e \cos \omega_i) + Z_{\text{rawmin}}D_{\text{min}}(1 + e \cos \omega_i) = 2 \quad (124)$$

This equation contains the raw observational signals ($Z_{\text{rawmax}}, Z_{\text{rawmin}}$), the known geometry (e), and the internal R.O.M. variables (β, ω_i) locked inside the D terms. Crucially, **the inclination angle i has been strictly eliminated**. This proves analytically that the true kinematic projection β is determined entirely by the asymmetry of the transverse baselines at the extrema, independent of the observer's viewing angle.

25.6.3 Analytical Extraction of $\sin i$

Once β and ω_i are constrained via Equation 124 (and verified via the balance phase O_o), the true orbital inclination is trivially extracted by substituting the definition of K_i back into Equation 122:

$$\sin i = \frac{\sqrt{1 - e^2}}{2\beta} [Z_{\text{rawmax}}D_{\text{max}}(\beta, e, \omega_i) - Z_{\text{rawmin}}D_{\text{min}}(\beta, e, \omega_i)] \quad (125)$$

25.6.4 Numerical Verification: S0-2 Extrema Synthesis

To verify this architecture, a forward-synthesis and reverse-extraction test was performed using the empirically established parameters of the S0-2 star (GRAVITY Collaboration: $e = 0.88466$, $\omega_i = 66.13^\circ$, $\beta = 0.00645$, $i = 134.0^\circ$).

Forward synthesis generates the exact transverse baselines $D_{\max} = 0.999501$ and $D_{\min} = 0.999775$, yielding the raw spectroscopic extrema:

- $Z_{\text{rawmax}} = 1.014019$
- $Z_{\text{rawmin}} = 0.993834$

Evaluating the Decryption Invariant (Equation 124) with these values yields exactly 2.000000, confirming absolute algebraic closure. Applying the extraction equation directly outputs $\sin i = 0.71931$, which immediately returns the true inclination $i = 133.99^\circ$.

Remark 25.2 (Epistemological Proof vs. Empirical Extraction). While the Decryption Invariant (Equation 124) provides the rigorous epistemological proof that the classical $M \sin i$ degeneracy is not a fundamental law of physics, its direct algebraic application is inherently bounded by instrumental limitations.

Because the baseline asymmetry ($D_{\max} - D_{\min}$) scales strictly as $\mathcal{O}(\beta^2)$ (e.g., $\approx 2.7 \times 10^{-4}$ for the S0-2 system), the algebraic inversion is hypersensitive to spectrographic noise. Any instrumental fluctuation or finite-sampling interpolation error at the exact extrema (Z_{rawmax} and Z_{rawmin}) will be mathematically amplified by the inversion matrix.

Therefore, the algebraic and statistical methods must operate in strict symbiosis. The algebraic system (Section 25.6) provides the exact topological initial locus, instantly dispatching the optimizer to the correct degeneracy ridge. The Bayesian MCMC framework (Section 25.4) then takes over, integrating the continuous phase curve to suppress instrumental noise and establish robust empirical confidence intervals.

This confirms that the classical $M \sin i$ degeneracy is algebraically penetrable without external metric priors or assumptions regarding the mass.

25.7 Conclusion

The classical $M \sin i$ degeneracy is not a fundamental limitation of orbital mechanics. It is an artifact of an incomplete, linearized description of the kinematic projection.

By rejecting speculative ontological baggage and restoring the full non-linear structure of the relational carriers S^1 and S^2 , R.O.M. provides two independent quadratic invariants ($\Delta\phi \propto \beta_{\text{int}}^2$ and $Z_{\text{sys}} \propto \beta^2$) that operationally decouple β from i . The inclination is then uniquely recovered from the linear amplitude.

This result removes a 130-year observational barrier and demonstrates that what appear as fundamental degeneracies in descriptive physics are often resolved as soon as the generative relational geometry is fully expressed. The Bayesian analysis provides rigorous statistical confirmation that the degeneracy ridge is correctly positioned to contain the independently verified physical truth.

Philosophical Implication

Degeneracies are not features of Nature; they are symptoms of incomplete substantival description. When the full relational geometry is restored, the apparent ambiguity dissolves into algebraic necessity.

26 Derivation of Gravitational Deflection and Lensing

In General Relativity, the deflection of light is obtained by integrating the null geodesic equations over a curved spacetime manifold, often relying on weak-field approximations and Taylor expansions. Within WILL Relational Geometry (RG), we reject both the background manifold and the use of mathematical approximations as non-operational ontological artifacts.

The system consists exclusively of its participants: the Source, the Lens (at periapsis p), and the Receiver. The total deflection angle must be derived as a strict, exact algebraic difference between their measurable relational phase states, without resorting to series expansions.

26.1 Gravitational Deflection and Interaction Gradient

Energy projections distribution among axis of relational carriers has to obey the conservation law (distribution between axis does not create nor destroy energy). There must exist a strict, algebraically closed gradient connecting all states, governed entirely by the kinematic projection β .

Theorem 26.1 (Unified Interaction Gradient). *The gravitational interaction capacity of any entity is determined by its available phase buffer β_Y . The geometric scaling factor Γ that dictates the distribution of the potential projection κ^2 onto the spatial trajectory is strictly defined by the arithmetic mean of the saturated carrier S^1 :*

$$\Gamma(\beta) = \frac{1 + \beta^2}{2} = 1 - \frac{\beta_Y^2}{2} \quad (126)$$

Proof. By the S^1 closure invariant ($\beta^2 + \beta_Y^2 = 1$), an object at rest ($\beta = 0$) possesses maximum internal phase ($\beta_Y = 1$). This phase acts as a geometric buffer, absorbing half of the relational gradient, yielding the classical partitioning $\Gamma = \frac{1}{2}$. As the spatial projection β increases, the internal clock slows, and the phase buffer β_Y depletes. At the topological limit $\beta \rightarrow 1$ (light), the internal phase collapses ($\beta_Y \rightarrow 0$). The buffer is exhausted, forcing the entity to absorb the full, unpartitioned gravitational gradient, yielding $\Gamma = 1$. $\square$

Using this gradient, we define the **Unified Closure Defect** for any trajectory—from a slow asteroid to a photon—as:

$$\delta_\varphi = \frac{\kappa_p^2}{\beta_p^2} \Gamma(\beta_p) = \frac{\kappa_p^2(1 + \beta_p^2)}{2\beta_p^2} \quad (127)$$

The geometric shape parameter (eccentricity) of the trajectory is derived directly from this defect:

$$e_\varphi = \frac{1}{\delta_\varphi} - 1 = \frac{2\beta_p^2}{\kappa_p^2(1 + \beta_p^2)} - 1 \quad (128)$$

Applying the exact algebraic transit equation for a distant observer ($\kappa_o \rightarrow 0$), we have $\cos o_\infty = -1/e_\varphi$. Using the same trigonometric extraction as established for light ($\sin(\frac{\Delta\varphi}{2}) = \frac{1}{e_\varphi}$), we arrive at the absolute, unified equation for gravitational deflection:

$$\Delta\varphi = 2 \arcsin\left(\frac{\kappa_p^2(1 + \beta_p^2)}{2\beta_p^2 - \kappa_p^2(1 + \beta_p^2)}\right) \quad (129)$$

Solving for $\beta_p = 1$ the total deflection angle $\Delta\varphi$ gives non-linear equation for light deflection in RG:

$$\Delta\gamma = 2 \arcsin\left(\frac{\kappa_p^2}{\kappa_{Xp}^2}\right) \quad (130)$$

Desmos and Colab

Desmos: [Algebraic Light Deflection \(one input derivation\)](#) Colab: [Gravitational Deflection.ipynb](#)

Verification of Topological Limits:

- **Newtonian Limit** ($\beta_p \ll 1$): The phase buffer is full ($\Gamma \rightarrow 0.5$). The eccentricity is dominated by $2\beta_p^2/\kappa_p^2$. The deflection reduces to the classical Rutherford/Newton scattering: $\Delta\varphi \approx \frac{\kappa_p^2}{\beta_p^2}$.
- **Relativistic Limit** ($\beta_p \rightarrow 0.99$): The phase buffer is nearly depleted ($\Gamma \rightarrow 0.99$). The trajectory stiffens, and the deflection angle approaches the photonic maximum, smoothly capturing the post-Newtonian factor without Taylor expansions.
- **Photonic Limit** ($\beta_p = 1$): The phase buffer is completely exhausted ($\Gamma = 1$). Substituting $\beta_p = 1$ yields $e = \frac{2}{\kappa_p^2(2)} - 1 = \frac{1}{\kappa_p^2} - 1 = \frac{\kappa_{Xp}^2}{\kappa_p^2}$. The equation resolves perfectly into the exact light deflection identity: $\Delta\varphi = 2 \arcsin\left(\frac{\kappa_p^2}{\kappa_{Xp}^2}\right)$.

Elimination of the "Massless" Myth

This formulation suggests that light is not a distinct ontological category governed by separate physical laws. There are no "massless particles" in RG. Light is simply the rightward topological limit of matter, where the internal phase β_Y reaches zero. The historical factor of 2 in gravitational lensing is not an anomaly of curved spacetime, but the inevitable geometric consequence of an exhausted phase buffer.

26.2 Gravitational Lensing and the Einstein Ring

In Relational Geometry, the description of gravitational lensing (deflection, convergence, shear, and magnification) does not require a background metric or the postulation of a curved manifold. It arises entirely as the geometric projection of the deflection angle onto the observer's plane.

By employing the standard geometric baseline equation for the observer-lens-source system, the true angular position θ_S of a source is related to its observed image position θ_I via:

$$\theta_S = \theta_I - \frac{D_{LS}}{D_S} \Delta\varphi_{\text{unified}}(\beta_p, \kappa_p(\theta_I)) \quad (131)$$

where D_{LS} and D_S are the relational baseline distances.

The potential projection at periapsis κ_p^2 is strictly a function of the observed angle θ_I , defined by the physical impact parameter $r_p = D_L \theta_I$:

$$\kappa_p^2(\theta_I) = \frac{R_s}{D_L \theta_I} \quad (132)$$

Theorem 26.2 (Exact Algebraic Einstein Ring). *For a perfectly aligned system ($\theta_S = 0$), the image forms a symmetric ring. The exact angular radius of this Einstein ring, θ_E , valid for all velocities and field strengths, is given by the implicit algebraic equation:*

$$\theta_E = 2 \frac{D_{LS}}{D_S} \arcsin \left(\frac{\kappa_p^2(\theta_E)(1 + \beta_p^2)}{2\beta_p^2 - \kappa_p^2(\theta_E)(1 + \beta_p^2)} \right) \quad (133)$$

Recovery of Topological Limits: To reveal the underlying gradient without altering the exact equation, we examine the weak-field astrophysical limit ($\kappa_p^2 \ll \beta_p^2$), where $\arcsin(x) \approx x$. The unified deflection simplifies algebraically to:

$$\theta_E \approx \frac{D_{LS}}{D_S} \frac{R_s}{D_L \theta_E} \left(\frac{1 + \beta_p^2}{\beta_p^2} \right) \implies \theta_E^2 \approx \frac{R_s D_{LS}}{D_L D_S} \left(\frac{1 + \beta_p^2}{\beta_p^2} \right) \quad (134)$$

Colab Notebook

[Gravitational Lensing.ipynb](#)

- **The Photonic Limit** ($\beta_p \rightarrow 1$): The kinematic factor evaluates exactly to $\frac{1+1}{1} = 2$. The equation resolves to $\theta_{E,\gamma} = \sqrt{\frac{2R_s D_{LS}}{D_L D_S}}$, perfectly matching the full General Relativistic prediction for light (equivalent to $4GM/c^2$).
- **The Relativistic Massive Limit** ($0 < \beta_p < 1$): The continuous phase-buffer depletion natively supplies the correct post-Newtonian scaling without external ad-hoc gradients.

Epistemological Consequence

The properties of the lens—convergence (κ_{lens}), shear (γ), and magnification (μ)—are simply the components of the 2D Jacobian matrix of the deflection field $\vec{\alpha}(\vec{\theta})$. Because the physics of the phase-buffer β_Y is fundamentally woven into the exact definition of $\Delta\varphi_{\text{unified}}$, no manual "weighting" of shear or convergence is required. The optical mapping emerges naturally from the pure algebraic closure of the S^1 and S^2 relational carriers.

27 Geometric Mechanism of Orbital Decay and Radiation

In standard formulations, the orbital decay of binary systems (e.g., the Hulse-Taylor pulsar) is attributed to the emission of gravitational waves carrying away energy and angular momentum. In WILL Relational Geometry, orbital contraction is not a mechanical loss of substance, but a strict topological necessity governed by the conservation of the Unitary State Vector.

Theorem 27.1 (Topological Orbital Contraction). *The emission of radiation from a closed relational system mandates a compensatory geometric contraction of its orbital radius to strictly preserve the unitary state invariant.*

Proof. By the structure of the Unitary State Manifold, the system's exact internal causal phase (τ) and its external relational projection (τ_Y) are strictly bound by the geometric invariant:

$$\tau_Y^2 + \tau^2 = 1. \quad (135)$$

Radiation - whether electromagnetic or gravitational—is operationally defined in RG as the transfer of internal causal order to the external relational network. Geometrically, this translates to the continuous depletion of the system's internal phase: $\tau \rightarrow \tau - \delta\tau$.

To maintain the unitary invariant ($= 1$), the exact external projection must concurrently grow to compensate for the lost internal phase:

$$\tau \downarrow \implies \tau_Y^2 \uparrow. \quad (136)$$

Recalling that the state difference is defined by the projection intensities $\tau_Y^2 = Q^2 - \kappa^2\beta^2$, and that $Q^2 \propto 1/r$, an absolute increase in τ_Y^2 mathematically forces an increase in both κ^2 and β^2 .

Consequently, the operational coordinate radius r between the interacting bodies must decrease. The orbit shrinks not due to an external mechanical drag, but because the system is geometrically forced to descend into higher state projections to mathematically balance the loss of its internal causal budget. $\square$

Remark 27.2 (The Radiation Limit). *This geometric decay naturally terminates at the Dynamic Horizon ($\kappa^2 = 1$). At this limit, the internal phase is completely exhausted ($\tau = 0$) and the state projection is maximal ($\tau_Y^2 = 1$). The system can no longer radiate, strictly forbidding the extraction of further relational phase from a merged causal horizon.*

28 Derivation of Density, Mass, and Pressure

28.1 Derivation of Density

Translating RG (2D) to Conventional Density (3D). *In RG κ^2 is the 2D parameter defined in the relational carrier S^2 . In conventional physics, the source term is volumetric density ρ , a 3D concept defined by the "cultural artifact" (a Newtonian "cannonball" model) of mass-per-volume.*

To bridge our 2D theory with 3D empirical data, we must create a "translation interface". We do this by explicitly adopting the conventional (Newtonian) definition of density, $\rho \propto M/r^3$, as our "translation target".

From the projective analysis established in the previous sections:

$$\kappa^2 = \frac{R_s}{r},$$

where κ emerges from the energy projection on the area of unit sphere S^2 , and $R_s = 2GM/c^2$ links to the mass scale factor $M = E_0/c^2$.

This leads to mass definition:

$$M = \frac{\kappa^2 c^2 r}{2G}$$

To translate this into a volumetric density, we first adopt the conventional 3D (volumetric) proxy, r^3 . This is not a postulate of RG, but the first step in applying the legacy (3D) definition of density:

$$\frac{M}{r^3} = \frac{\kappa^2 c^2}{2Gr^2}$$

This expression, however, is incomplete. Our κ^2 "lives" on the 2D surface S^2 (which corresponds to 4π), while the r^3 proxy implicitly assumes a 3D volume. To correctly normalize the 2D parameter κ^2 against the 3D volume, we must apply the geometric normalization factor of the S^2 carrier by deviding on to area of the sphere, which is $1/4\pi$.

This normalization is the necessary geometric step to interface the 2D relational carrier (S^2) with the 3D legacy definition of density:

$$\begin{aligned} \rho &= \frac{1}{4\pi} \left(\frac{\kappa^2 c^2}{2Gr^2} \right) \\ \rho &= \frac{\kappa^2 c^2}{8\pi Gr^2} \end{aligned}$$

Local Density $\equiv$ Relational Projection

Maximal Density. *At $\kappa^2 = 1$ (the horizon condition (for non rotating systems), $r = R_s$), this density reaches its natural bound, $\rho_{\max}$, which is derived purely from geometry:*

$$\rho_{\max} = \frac{c^2}{8\pi Gr^2}$$

Normalized Relation. *Thus, our "translation" reveals an identity: the geometric projection κ^2 is simply the ratio of density to the maximal density:*

$$\boxed{\kappa^2 = \frac{\rho}{\rho_{\max}} \quad \text{arrow} \quad \kappa^2 \equiv \Omega}$$

28.2 Self-Consistency Requirement

The mass scale factor can be expressed in two equivalent ways.

From the geometric definition:

$$M = \frac{\kappa^2 c^2 r}{2G}.$$

From the energy density:

$$M = \alpha r^n \rho.$$

Substituting $\rho = \frac{\kappa^2 c^2}{8\pi G r^2}$ into $M = \alpha r^n \rho$ gives

$$M = \frac{\alpha \kappa^2 c^2 r^{n-2}}{8\pi G}.$$

Equating the two forms:

$$\frac{\alpha r^{n-2}}{8\pi} = \frac{r}{2}.$$

For the mass M to remain a constant independent of the measurement scale r , the exponent must be $n = 3$, yielding $\alpha = 4\pi$. Hence,

$$M = 4\pi r^3 \rho,$$

which closes the consistency loop between the geometric and density-based formulations.

Desmos Project

[Derivation of Density and Pressure](#)

28.3 Pressure as Surface Curvature Gradient

In the RG framework pressure is not a thermodynamic assumption but the direct consequence of curvature gradients. The radial balance relation gives

$$P(r) = \frac{c^4}{8\pi G} \frac{1}{r} \frac{d\kappa^2}{dr}.$$

Using $\kappa^2 = R_s/r$, one finds $d\kappa^2/dr = -\kappa^2/r$, hence

$$P(r) = -\frac{\kappa^2 c^4}{8\pi G r^2}.$$

Since the local energy density is

$$\rho(r) = \frac{\kappa^2 c^2}{8\pi G r^2},$$

this yields the invariant equation of state

$$P(r) = -\rho(r) c^2.$$

Interpretation. P is a surface-like negative pressure (isotropic tension), not a bulk volume pressure. It expresses the resistance of energy-geometry itself to changes in projection.

Consistency. If one formally freezes the projection parameter ($d\kappa^2/dr = 0$), then $P = 0$. But in this case the angular curvature terms remain uncompensated, and the field equation is no longer satisfied. Any nontrivial radial dependence of κ inevitably generates the negative tension

$$P = -\rho c^2,$$

which precisely cancels the residual curvature. Thus the negative pressure is not optional but a necessary ingredient for full self-consistency.

Desmos Project

[Derivation of Density and Pressure](#)

Maximum pressure. At the geometric bound $\kappa^2 = 1$ (horizon condition), the density saturates at

$$\rho_{\max} = \frac{c^2}{8\pi G r^2},$$

and the corresponding pressure is

$$P_{\max} = -\rho_{\max} c^2 = -\frac{c^4}{8\pi G r^2}.$$

This negative surface pressure represents the ultimate tension limit of spacetime fabric at a given scale r .

Pressure in WILL is the intrinsic surface tension of energy-geometry, saturating at $P_{\max} = -c^4/(8\pi G r^2)$.

Physical meaning

The negative pressure is not an exotic substance but the geometric tension required to maintain self-consistency when κ^2 varies radially. It's the relational analogue of surface tension in a soap bubble.

29 Unified Geometric Field Equation

From the energy-geometry equivalence, the complete description of gravitational phenomena reduces to a single algebraic relation linking the geometric scale to the energy density ratio:

$$\kappa^2 = \frac{R_s}{r} = \frac{\rho_{\text{field}}}{\rho_{\max}}$$

This identity defines the **local energy state of the geometry itself**. Here $\rho_{\max} = c^2/(8\pi G r^2)$ is the saturation density limit, and ρ_{field} is the effective energy density of the relational curvature.

29.1 Field Equation and Matter Sources

For a static, spherically symmetric configuration containing matter with density $\rho_{\text{matter}}(r)$, the relationship is governed by the differential accumulation of the potential:

$$\frac{d}{dr}(r \kappa^2) = \frac{8\pi G}{c^2} r^2 \rho_{\text{matter}}(r) \quad (137)$$

This expression reproduces the tt -component of the Einstein field equations.

The Vacuum Solution ($\rho_{\text{matter}} = 0$). In the vacuum region outside a central mass, the source density vanishes ($\rho_{\text{matter}} = 0$). The field equation implies conservation of the projection budget:

$$\frac{d}{dr}(r \kappa^2) = 0 \implies r \kappa^2 = \text{const} = R_s.$$

Thus, we recover the potential law of WILL RG:

$$\kappa^2 = \frac{R_s}{r}.$$

Resolution of Roles

1. **The Identity** $\kappa^2 = \rho/\rho_{\max}$ describes the state of the field geometry.
 2. **The Equation** $(r \kappa^2)' \sim \rho_{\text{matter}}$ describes how matter generates that geometry.
- In vacuum, the generator is zero, but the field persists as the algebraic structure $\kappa^2 = R_s/r$.

30 No Singularities, No Hidden Regions

This framework introduces no interior singularities, no coordinate patches hidden behind horizons, and no ambiguous initial conditions. The geometric field equation:

$$\frac{R_s}{r} = \frac{8\pi G}{c^2} r^2 \rho = \kappa^2$$

ensures that curvature and energy density evolve smoothly and remain bounded across all observable scales.

WILL Relational Geometry resolves the singularity problem not by regularizing divergent terms, nor by introducing quantum effects, but by geometrically constraining the domain of valid projections. Curvature is always finite, and energy remains bounded by construction. Black holes become energetically saturated but non-singular regions, described entirely by finite, dimensionless parameters.

This projectional approach provides a clean, intrinsic termination to gravitational collapse, replacing singular end-points with structured, maximally curved boundaries.

- **Surface-scaled closure (vs. volume filling).** Mass follows the algebraic closure $M = 4\pi r^3 \rho$ with $\rho = \kappa^2 c^2 / (8\pi G r^2)$; the 4π is the spherical projection measure, not a Newtonian volume average.
- **Natural bounds.** The constraint for non rotating systems $\kappa^2 \leq 1$ enforces $\rho \leq \rho_{\max}$ and $|P| \leq |P_{\max}| = c^4 / (8\pi G r^2)$, avoiding singularities without extra hypotheses.

31 Theoretical Ouroboros

Closure

Ontological principle is proven as the inevitable consequence of geometric consistency. Field Equation $\Longleftrightarrow$ Ontological Principle

We have shown that this single Ontological Principle (3.3), through pure geometric reasoning, necessarily leads to an equation which mathematically expresses the very same equivalence we began with. We started with $SPACETIME \equiv ENERGY$, from which geometry and physical laws are logically derived, and these derived laws then loop back to intrinsically define and limit the very nature of energy and spacetime, proving the self-consistency of the initial idea.

$$SPACETIME \equiv ENERGY.$$

The ratio of geometric scales equals the ratio of energy densities.

$$\kappa^2 = \frac{R_s}{r} = \frac{\rho}{\rho_{\max}}$$

$$\begin{array}{c} SPACETIME \text{ GEOMETRY} \\ \equiv \\ ENERGY \text{ DISTRIBUTION} \end{array}$$

Summary

All physical structure emerges from the single relational equivalence:

$$SPACETIME \equiv ENERGY$$

From this, by enforcing geometric self-consistency, one necessarily arrives at the Unified Geometric Field Equation:

$$\kappa^2 = \frac{R_s}{r} = \frac{\rho}{\rho_{\max}}.$$

This is not an external law but an intrinsic closure relation: geometry and energy are two mutually defining projections of a single entity. It represents the completion of the theoretical Ouroboros — where the principle generates its own mathematical expression and the expression in turn validates the principle.

From a philosophical and epistemological point of view, this can be considered the crown achievement of any theoretical framework - the "Theoretical Ouroboros". But regardless of the aesthetic beauty of this result, let us remain sceptical.

32 W_{ILL} : Unity of Relational Structure

The ontological principle

$$SPACETIME \equiv ENERGY$$

states that there is only one closed relational resource - WILL. What we call space, time and energy are different projections of the same relational structure. For any energy-closed system observed from a relational origin, this resource appears through four operational projections:

Desmos project

WILL = 1 Derivation of the W_{ILL} invariant.

These quantities are defined as correlated projections of the same underlying WILL structure. In the dimensionful form we write:

$$M = \frac{\beta^2}{\beta_Y} \frac{c^2 a}{G}, \quad E = \frac{\kappa^2}{\kappa_X} \frac{c^4 a}{2G}, \quad T = \kappa_X \left(\frac{2Gm_0}{\kappa^2 c^3} \right)^2, \quad L = \beta_Y \left(\frac{Gm_0}{\beta^2 c^2} \right)^2,$$

where (β, β_Y) and (κ, κ_X) are the kinematic and gravitational projections on the carriers S^1 and S^2 , m_0 is the rest mass, $E_0 = m_0 c^2$ is the rest energy, and a is the relational scale of the system as semi-major axis (average length per cycle within this system).

The closure conditions of the carriers,

$$\beta^2 + \beta_Y^2 = 1, \quad \kappa_X^2 + \kappa^2 = 1,$$

together with the energetic exchange condition

$$\kappa^2 = 2\beta^2,$$

fix a unique dimensionless combination of these four projections. Combining E , T , M , and L we obtain

$$W_{ILL} \equiv \frac{ET}{ML} = \frac{\frac{E_0}{\kappa_X} \kappa_X t^2}{\frac{m_0}{\beta_Y} \beta_Y a^2} = \frac{E_0 t^2}{m_0 a^2}.$$

By the relations that tie temporal and $t = a/c$ and spacial $a = R_s/\kappa^2$ scales this ratio is identically equal to unity for any closed system:

$$W_{ILL} = \frac{ET}{ML} = 1.$$

All dimensionful constants cancel automatically; the value is fixed by the geometry of the carriers, not by a choice of units.

Desmos project

WILL EARTH GPS calculates GPS satellite time shift and testing $W_{ILL} = 1$ on the Earth-GPS system

The same invariant can be written in a phase-normalized form, using local projections

$$E_o = \frac{E_0}{\kappa_{X_o}}, \quad M_o = \frac{m_0}{\beta_{Y_o}}, \quad T_o = \kappa_{X_o} t_o^2, \quad L_o = \beta_{Y_o} r_o^2,$$

Equivalently, for any state (β_o, κ_o) , any scale r_o and any phase along the orbit. Then

$$W_{ILL} = \frac{E_o T_o}{M_o L_o} = 1 \quad \text{for ALL ENERGY'S, ALL SCALES and ALL PHASES.}$$

$$\frac{E_o}{M_o} = \frac{L_o}{T_o},$$

so the energy sector and the spacetime sector are not independent. Every change of the relational state rescales (E_o, M_o) and (T_o, L_o) coherently so that this equality is always preserved. The familiar practice of treating energy-mass and space-time as separate blocks is therefore an ontological approximation: in WILL they are locked by a single relational constraint.

$$\text{Geometry} \equiv \text{Energy} \equiv \text{Causality} \equiv \text{WILL},$$

$$W_{ILL} = 1.$$

in the precise sense that one and the same conserved relational resource appears as mass, energy, time and length, but always in a way that keeps their ratio fixed.

32.1 Interpretive Note: The Name "WILL"

The term **WILL** stands for **SPACE-TIME-ENERGY**. It is both a formal shorthand and a philosophical statement: the universe is not a stage where energy acts through time upon space, but a single self-balancing structure whose internal distinctions generate all phenomena. The name also serves as a gentle irony toward anthropic thinking: the Cosmos does not possess "will" - yet through WILL, it manifests All that Is.

Summary

$$\mathbf{WILL} \equiv \frac{ET}{ML} = 1 \quad \Longleftrightarrow \quad \text{Geometry} = \text{Energy} = \text{Causality.}$$

WILL is not the unit of something - but the Unity of Everything.

33 Ontological Shift: From Descriptive to Generative Physics

In conventional physics the methodology follows a descriptive paradigm:

1. Observable phenomena are identified.
2. Empirical regularities are codified as "laws of nature."
3. Mathematical formalisms are constructed to describe these regularities.

Thus, physical laws are always introduced as external assumptions that model what is seen. Even in General Relativity, where geometry plays the central role, the equivalence principle and the metric postulate are still external inputs.

The RG framework inverts this paradigm. Laws are not added on top of observations; they are generated as inevitable consequences of relational geometry:

- There are no independent axioms such as "inertial mass equals gravitational mass."
- Such relations appear automatically as algebraic identities enforced by the geometry.
- What classical physics calls "laws of nature" are secondary shadows of the single relational principle:

$$\text{SPACETIME} \equiv \text{ENERGY.}$$

Summary

Standard Physics: Laws describe what we observe.

Relational Geometry: Laws are generated as necessary products of closure and self-consistency.

In this sense, the ontological role of physical law is transformed. Physics ceases to be a catalog of empirical descriptions, and becomes the logical unfolding of a single relational structure. WILL identifies the necessary conditions under which all observed phenomena arise.

Descriptive Physics (Standard)	Generative Physics (WILL)
Phenomena are <i>observed</i> first, then summarized into empirical laws.	Laws emerge as <i>inevitable consequences</i> of relational geometry.
Physical laws are <i>assumptions</i> introduced to model reality.	Physical laws are <i>identities</i> , enforced by geometric self-consistency.
Time and space are treated as external backgrounds.	Time and space are <i>projections of energy relations</i> .
Dynamics = evolution of states <i>in time</i> .	Dynamics = ordered succession of balanced configurations; <i>time is emergent</i> .
Goal: <i>describe</i> what is observed.	Goal: <i>show why nothing else is possible</i> .

Table 8: Ontological contrast between standard descriptive physics and the generative paradigm of WILL Relational Geometry.

Phenomenon	Standard (GR) Result	Relational Geometry (RG)
GPS time shift / gravitational redshift	Frequency shift = combination of kinetic (SR) and gravitational (GR) effects.	Single symmetric law: $\tau = \beta_Y \cdot \kappa_X$, $E_{\text{loc}} = \frac{E_0}{\sqrt{(1-\beta^2)(1-\kappa^2)}} = \frac{E_{\text{loc}}}{\tau} 38.52 \mu\text{s/day}$ verified directly with GPS satellites.
Photon sphere, ISCO, horizons	Derived by solving geodesic equations in Schwarzschild metric.	Critical radii emerge from simple symmetry's (Photon sphere: $\theta_1 = \theta_2 = 54.73^\circ$ ("magic angle") $Q^2 = \kappa^2 + \beta^2 = 1$. ISCO: $Q = Q_Y$,
Mercury's perihelion precession	Complex expansion of Einstein field equations.	Same number obtained from RG with $\Delta\varphi = \frac{2\pi\tau_Y^2}{1-e^2} = 43''/\text{century}$. Using simple algebra.
Binary pulsar orbital decay	Explained via quadrupole radiation formula; requires asymptotic Bondi mass.	Emerges from balance of projection invariants without asymptotic constructs. $\Delta P \approx -2.42 \times 10^{-12} \text{ s/s}$ (predicted)
Cosmological redshift	Photon "loses energy" as universe expands.	Energy conserved; redshift = redistribution of projection parameters. (Details in WILL PART II)
Cosmological absolute scale (Supernovae fit)	Hubble-like expansion, ΛCDM fits	$H_0 \equiv \sqrt{8\pi G\rho_\gamma/(3\alpha^2)} = 68.15$ Derived from CMB temperature and α connecting micro and macro scales (Details in WILL PART II-III).
Cosmological constant Λ	Added by hand to fit data ("dark energy").	Arises naturally as $\Lambda = 2/3r^2$. No extra entities required. (More details in WILL PART I and II)
Singularities	Predicted in black holes and big bang ($\rho \rightarrow \infty$).	Forbidden: density bounded by $\rho_{\text{max}} = c^2/(8\pi G r^2)$.
Local gravitational energy	"Cannot be localized" (only ADM/Bondi at infinity).	Directly measurable via κ , e.g. from light deflection angle or red shift.
Unification with QM	No natural unification in GR framework.	Same projectional law applies from microscopic $\alpha = \beta_1$ (QM) to cosmic $\kappa^2 = \Omega_\Lambda$ (GR, COSMO) scales. (Details in WILL PART II and III)

Table 9: Classical GR results vs. WILL RG outcomes. Known effects are recovered by simpler symmetric laws, while new predictions eliminate singularities and explain cosmology without dark entities.

34 Conclusion

WILL Relational Geometry fully reproduces the predictive content and central equations of both Special and General Relativity, while simultaneously addressing their foundational inconsistencies:

- (1) the lack of an operational definition of local gravitational energy density in GR,
- (2) the artificial separation of kinetic and gravitational energy in SR and GR, and
- (3) the emergence of singularities as pathological artifacts of coordinate-based models. By treating energy and its transformations as the true basis of geometry, RG unifies and extends these frameworks into a fully relational and operationally grounded description of spacetime and energy.

By focusing on the projectional nature of energy, we have shown that spacetime itself is merely the manifestation of energy.

From a single Ontological Principle-that spacetime is equivalent to energy - we derived all the mathematical apparatus needed to describe gravitational and relativistic phenomena. This unification, showing that energy, time and space are merely different projections of the same underlying structure.

Special and General Relativity emerge from the same geometric principles.

This approach offers distinct advantages:

- *Conceptual clarity* - understanding physics through pure geometry
- *Computational efficiency* - significantly reducing complexity
- *Epistemological hygiene* - deriving results from minimal assumptions
- *Philosophical depth* - redefining our understanding of time, mass, and causality

WILL Relational Geometry inverts our fundamental understanding:

Spacetime and energy are mutually defining aspects of a single relational structure.

Final Summary

SPACETIME $\equiv$ ENERGY.

35 References:

References

- [1] Schwarzschild, K. (1916). *Sitzungsberichte der Königlich Preussischen Akademie der Wissenschaften*.
- [2] Tolman, R.C. (1939). *Phys. Rev.* **55**, 364; Oppenheimer, J.R., Volkoff, G.M. (1939). *Phys. Rev.* **55**, 374.
- [3] Dyson, F.W., Eddington, A.S., Davidson, C. (1920). *Phil. Trans. R. Soc. A* **220**, 291-333.
- [4] Lebach, D.E. et al. (1995). *Phys. Rev. Lett.* **75**, 1439.
- [5] Ashby, N. (2003). *Living Rev. Relativity* **6**:1.
- [6] Will, C.M. (2014). *Living Rev. Relativity* **17**:4.
- [7] Williams, J.G., Turyshev, S.G., Boggs, D.H. (2006). *Adv. Space Res.* **37**, 67-71.
- [8] Kramer, M. et al. (2021). *Phys. Rev. X* **11**, 041050.
- [9] Misner, Thorne, Wheeler (1973). *Gravitation*.
- [10] et al 2022 *ApJ* 933 49 DOI 10.3847/1538-4357/ac752f
- [11] Espenak, F. (2014). *Seven Century Catalog of Mercury Transits: 2001 to 2700*. NASA Eclipse Web Site.
- [12] Williams, M. (2016). *How Far is Mercury From the Sun? Universe Today* (citing NASA data).
- [13] Mamajek, E.E. et al. (2015). *IAU 2015 Resolution B3 on Recommended Nominal Conversion Constants*. *arXiv:1510.07674*.