

# WILL

## *Relational Orbital Mechanics (R.O.M.)*

Anton Rize  
antonrize@willrg.com \*

October 2025

### Abstract

This supplement document to [first part](#) of the [WILL trilogy](#) presents **Relational Orbital Mechanics (R.O.M.)** – a closed algebraic system that classifies allowed states of [relational configuration](#) without invoking a [background spacetime](#), a metric tensor, or a mass and gravitational constant.

This closed algebraic system recovers the leading contributions to perihelion precession, gravitational and transverse Doppler shifts, light deflection and frame-dragging through elementary closed-form expressions.

High-precision validation shows that all 66 multi-form identities are internally consistent to 50 significant digits and that the leading terms match the corresponding post-Newtonian expansions of general relativity. Higher-order differences remain below the sensitivity of current observations, including those of the S2 star.

Complete phase-resolved orbits and system scales are recovered directly from spectroscopic and chronometric observables, with  $G$  and  $M$  entering only as unit-conversion factors. Key features of Kerr geometry follow algebraically from the same closure relation. Gravitational waves and the radiative two-body problem remain to be formulated within the relational ontology.

From the [dimensionless projections](#)  $\beta$  (kinematic) and  $\kappa$  (potential) on the [relational carriers](#)  $S^1$  and  $S^2$ , and from the [closure theorem](#)  $\kappa^2 = 2\beta^2$  (forced by degree-of-freedom indifference), all major observed orbital phenomena are derived algebraically as indefinable consequence of Relational Geometry:

- Kepler's Third Law, the vis-viva relation, conservation of angular momentum, and the derivation of eccentricity emerge as strict Pythagorean invariants.
- Orbital precession is recovered as the accumulation of the relational phase divergence  $\tau_Y^2$ , normalized by the orbital shape factor  $1 - e^2$ , yielding the exact GR value for Mercury without differential equations.
- The Schwarzschild radius  $R_s$  and all orbital scales are expressed directly in terms of spectroscopic and chronometric observables ( $z_\odot$ ,  $\theta_\odot$ ,  $T$ ,  $e$ ), bypassing mass  $M$  and Newton's constant  $G$ .
- Gravitational deflection of massive particles and light follows from the [energy symmetry law](#) with no weak-field expansion.

R.O.M. demonstrates that the operational content of gravitational dynamics is fully captured by the algebra of relational projections. The construct serves as a rigorous, empirical realization of the foundational methodological principles established in [WILL Part I Relational Geometry](#). As an Open Research all data and calculation tools are publicly available at [willrg.com](https://willrg.com) and <https://github.com/AntonRize/WILL>.

---

\*This work is archived on Zenodo: <https://zenodo.org/record/17115270>

# Contents


|                                                                          |           |
|--------------------------------------------------------------------------|-----------|
| <b>1 Quick Recap</b>                                                     | <b>2</b>  |
| <b>2 Closed Algebraical System of R.O.M. Equations</b>                   | <b>5</b>  |
| 2.1 Translation to Legacy Tensor Parameters                              | 9         |
| <b>3 Relational Origin of:</b>                                           | <b>10</b> |
| 3.1 Spectroscopic Shifts                                                 | 10        |
| 3.2 Kepler's Third Law                                                   | 11        |
| 3.3 Eccentricity                                                         | 12        |
| 3.4 Vis-Viva                                                             | 13        |
| 3.5 Classical Acceleration                                               | 14        |
| 3.6 Angular Momentum Conservation                                        | 14        |
| 3.7 Orbital Precession as Phase Accumulation                             | 15        |
| 3.8 Dynamic Event Horizon                                                | 15        |
| 3.9 Gravitational Deflection and Lensing                                 | 16        |
| 3.10 Schwarzschild Metric                                                | 18        |
| <b>4 Intrinsic Unitless Core: Relational Parameterization</b>            | <b>18</b> |
| 4.1 The Intrinsic Relational Core: Unitless Formulation                  | 18        |
| 4.2 Observational Input Channels                                         | 19        |
| 4.3 Operational Algorithm                                                | 19        |
| 4.4 Relational Parameterization: Empirical Validation                    | 20        |
| <b>5 Time-Density Invariant (<math>\tau_D</math>)</b>                    | <b>23</b> |
| <b>6 Algebraic Determination of the Horizon Scale</b>                    | <b>24</b> |
| 6.1 Method A: Differential (Two-Point Method)                            | 24        |
| 6.2 Method B: Geometric Resonance (Balance Point Method)                 | 25        |
| 6.3 Method C: Instantaneous (Arbitrary Phase Method)                     | 25        |
| <b>7 Rotational Systems (Kerr Without Metric)</b>                        | <b>26</b> |
| 7.1 Chiral Asymmetry: The Algebraic Origin of Frame-Dragging             | 27        |
| 7.2 Legacy Translation: Lense-Thirring Equivalence                       | 28        |
| 7.3 Epistemological Conclusion                                           | 29        |
| <b>8 Mercury's Precession: GR 1PN as First-Order Approximation of RG</b> | <b>29</b> |
| 8.1 Statement of the Verification Problem                                | 29        |
| 8.2 Methodology                                                          | 29        |
| 8.3 Results                                                              | 29        |
| 8.4 Interpretation                                                       | 30        |
| 8.5 Reproducibility                                                      | 30        |
| <b>9 S2 Star (Sgr A*): R.O.M. vs GR and Instrumental Systematics</b>     | <b>30</b> |
| <b>10 Relational Derivation of the L1 Equilibrium Point</b>              | <b>32</b> |
| 10.1 Observational Primitives                                            | 32        |
| 10.2 Structural Parameters from Relational Projections                   | 32        |
| 10.3 Earth-L1 Distance Derivation                                        | 33        |

## 1 Quick Recap

In following sections we will demonstrate that RG's relational projections ( $\kappa$ ,  $\beta$ ) are ontologically and operationally independent from mass  $M$  and gravitational constant  $G$ . We will show that  $R_s$  as a horizon scale derived directly from the phase relations of light and will show how all observational phenomenon of orbital systems are generated as the unavoidable consequences of Relational Geometry.

But first lets remind our selves key derivations and logical chain established prior:

|                                                                                                                                                                                                                                                                                                    |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Methodological constraints:</b>                                                                                                                                                                                                                                                                 |
| $\downarrow (\text{Epistemic Hygiene} + \text{Relational Origin} + \text{Ontological Minimalism} + \text{Math Transparency}) \downarrow$                                                                                                                                                           |
| <b>Closure + Causal Continuity + Isotropy Theorems</b>                                                                                                                                                                                                                                             |
| $\downarrow (\text{apply to existing physics}) \downarrow$                                                                                                                                                                                                                                         |
| <b>Two Primitives:</b> $\underbrace{\text{fixed manifold} + \text{metric}}_{\text{structure}} + \underbrace{\text{fields} + \text{constants}}_{\text{dynamics}}$                                                                                                                                   |
| $\downarrow (\text{Ontological reduction by Relational Origin Principle} - \text{no background allowed}) \downarrow$                                                                                                                                                                               |
| <b>One Primitive: WILL <math>\equiv</math> SPACE-TIME-ENERGY <math>\Rightarrow</math> SPACETIME <math>\equiv</math> ENERGY</b>                                                                                                                                                                     |
| $\downarrow (\text{by Closure} + \text{Causal Continuity} + \text{Isotropy Theorems: derive primal relational carriers}) \downarrow$                                                                                                                                                               |
| <b><math>S^1</math> (1-DOF kinematic) + <math>S^2</math> (2-DOF potential)</b>                                                                                                                                                                                                                     |
| $\downarrow (\text{by Duality of Relation Lemma}) \downarrow$                                                                                                                                                                                                                                      |
| <b>Relational Conservation:</b> $\underbrace{\text{Amplitude}^2}_{\text{External Interaction}} + \underbrace{\text{Phase}^2}_{\text{Internal Existence}} = 1$                                                                                                                                      |
| $\underbrace{\beta^2 + \beta_Y^2 = 1}_{\text{Kinematic}} \text{ on } S^1 \text{ and } \underbrace{\kappa^2 + \kappa_X^2 = 1}_{\text{Potential}} \text{ on } S^2$                                                                                                                                   |
| $\downarrow (\text{by Conservation Theorem}) \downarrow$                                                                                                                                                                                                                                           |
| <b>Minkowski interval:</b> $\underbrace{\beta^2 + \beta_Y^2 = 1}_{\text{irreducible primitive}} + \underbrace{x, y, z, t \dots}_{\text{coordinate inflation}} = \underbrace{c^2 d\tau^2 = c^2 dt^2 - dx^2 - dy^2 - dz^2}_{\text{mathematical scaffolding}}$                                        |
| <b>Schwarzschild interval:</b> $\underbrace{\kappa^2 + \kappa_X^2 = 1}_{\text{irreducible primitive}} + \underbrace{x, y, z, t \dots}_{\text{coordinate inflation}} = \underbrace{c^2 d\tau^2 = c^2(1 - (R_s/r))dt^2}_{\text{mathematical scaffolding}}$                                           |
| $\downarrow (\text{by DOF-Indifference Lemma: equal quadratic weight to each independent DOF}) \downarrow$                                                                                                                                                                                         |
| <b>Closure Theorem:</b> $\underbrace{1 \text{ Amplitude}^2}_{\text{on } S^2} = \underbrace{2 \text{ Amplitude}^2}_{\text{on } S^1} \equiv \kappa^2 = 2\beta^2$                                                                                                                                     |
| $\downarrow (\text{by Relational Closure and Invariance Theorems}) \downarrow$                                                                                                                                                                                                                     |
| <b>Energy Symmetry Law:</b> $\Delta E_{A \rightarrow B} + \Delta E_{B \rightarrow A} = 0$                                                                                                                                                                                                          |
| $\Delta E_{A \rightarrow B} = E_0(\kappa_{X,B}/\beta_{Y,B} - \kappa_{X,A}/\beta_{Y,A}); \Delta E_{B \rightarrow A} = E_0(\kappa_{X,A}/\beta_{Y,A} - \kappa_{X,B}/\beta_{Y,B})$                                                                                                                     |
| $\downarrow (\text{check the required energy to close the ledger at boundaries}) \downarrow$                                                                                                                                                                                                       |
| <b>Speed of light:</b> $\beta \rightarrow 1 \rightarrow \beta_Y \rightarrow 0 \rightarrow E_{\text{loc}} = E_0 \cdot \kappa_X / 0 \rightarrow \infty \rightarrow \text{requires } \infty \text{ energy}$                                                                                           |
| <b>Event Horizon:</b> $\kappa \rightarrow 1 \rightarrow \kappa_X \rightarrow 0 \rightarrow E_{\text{loc}} = E_0 \cdot 0 / \beta_Y \rightarrow 0 \rightarrow \text{requires } 0 \text{ energy}$                                                                                                     |
| $\downarrow (\text{by Unified Relational Scaling Lemma}) \downarrow$                                                                                                                                                                                                                               |
| <b>Equivalence Principle</b> $m_g \equiv m_i \equiv m = E_0/c^2$                                                                                                                                                                                                                                   |
| $\downarrow (\text{by collapsing two-point Energy Symmetry Law into a single-point approximation}) \downarrow$                                                                                                                                                                                     |
| <b>Classical, Lagrangian and Hamiltonian Mechanics</b>                                                                                                                                                                                                                                             |
| $\underbrace{\Delta E_{A \rightarrow B} + \Delta E_{B \rightarrow A} = 0}_{\text{exact symmetry law}} \approx \underbrace{\frac{1}{2}(\kappa_A^2 - \kappa_B^2) + \frac{1}{2}(\beta_B^2 - \beta_A^2)}_{\text{first order approximation}} \approx \underbrace{T \pm V}_{\text{ontological collaps}}$ |
| $\downarrow (\text{apply RG to orbital mechanics}) \downarrow$                                                                                                                                                                                                                                     |
| <b>Relational Orbital Mechanics [R.O.M.]: a closed algebraic system of equations</b>                                                                                                                                                                                                               |
| $\downarrow (\text{by Inverse Square, Closure and Causal Continuity Theorems}) \downarrow$                                                                                                                                                                                                         |
| <b>No Singularities:</b> $\beta_{\text{max}}^2 = 1 \rightarrow \kappa_{\text{max}}^2 = 2 \rightarrow r_{\text{min}} = R_s / \kappa_{\text{max}}^2 \rightarrow r_{\text{min}} = R_s / 2$                                                                                                            |
| <b>The point <math>r = 0</math> is absent from the admissible relational range</b>                                                                                                                                                                                                                 |
| $\downarrow (\text{apply 2D to 3D translation interfaces}) \downarrow$                                                                                                                                                                                                                             |
| <b>Geometric Field Equation</b> $\kappa^2 = R_s / r = \rho_{\text{field}} / \rho_{\text{max}}$                                                                                                                                                                                                     |
| $\downarrow (\text{The ratio of spacetime geometry } R_s/r \text{ equals the ratio of energy densities } \rho/\rho_{\text{max}}) \downarrow$                                                                                                                                                       |
| $\downarrow$ <b>Theoretical Ouroboros</b> $\downarrow$                                                                                                                                                                                                                                             |
| <b>SPACETIME <math>\equiv</math> ENERGY <math>\Leftrightarrow</math> SPACETIME GEOMETRY <math>\equiv</math> ENERGY DENSITY</b>                                                                                                                                                                     |



| $\theta_1 = \arccos(\beta), \quad \theta_2 = \arcsin(\kappa), \quad \kappa^2 = 2\beta^2, \quad a = \text{semi-major axis}$ |                                        |
|----------------------------------------------------------------------------------------------------------------------------|----------------------------------------|
| Algebraic Form                                                                                                             | Trigonometric Form                     |
| $\beta = v/c$                                                                                                              | $\beta = \cos(\theta_1)$               |
| $\kappa = \sqrt{R_s/a}$                                                                                                    | $\kappa = \sin(\theta_2)$              |
| $\beta_Y = \sqrt{1 - \beta^2}$                                                                                             | $\beta_Y = \sin(\theta_1)$             |
| $\kappa_X = \sqrt{1 - \kappa^2}$                                                                                           | $\kappa_X = \cos(\theta_2)$            |
| $p = E_0/c \cdot \beta/\beta_Y$                                                                                            | $p = E_0/c \cdot \cot(\theta_1)$       |
| $p_g = E_0/c \cdot \kappa/\kappa_X$                                                                                        | $p_g = E_0/c \cdot \tan(\theta_2)$     |
| $\tau = \beta_Y \kappa_X$                                                                                                  | $\tau = \sin(\theta_1) \cos(\theta_2)$ |
| $\tau_Y = \sqrt{1 - \tau^2}$                                                                                               | $\tau_Y = \sqrt{1 - \tau^2}$           |
| $Q = \sqrt{\kappa^2 + \beta^2} = \sqrt{3}\beta$                                                                            | $Q = \sqrt{3} \cos(\theta_1)$          |
| $Q_Y = \sqrt{1 - Q^2} = \sqrt{1 - \kappa^2 - \beta^2} = \sqrt{1 - 3\beta^2}$                                               | $Q_Y = \sqrt{1 - 3\cos^2(\theta_1)}$   |

Table 1: Unified representation of relativistic and gravitational effects for closed systems.





Figure 1: **Relational Self-Centering.** The total shift  $Q$  is defined by the amplitude projections  $\beta$  and  $\kappa$ . Interaction is causal only when the relational origin  $(0, 0)$  of B lies within A's horizon ( $Q < 1$ ), ensuring mutual coverage.

Relational reciprocity = invariance of the norm  $Q$  under self-centering.

There is no common background arena. Mutual total shift magnitudes  $Q$  computed from each observer's own relational origin.

| Phenomenon             | Radius $r$       | $\beta^2$     | $\kappa^2$    | $Q^2$         | Comment                                                          |
|------------------------|------------------|---------------|---------------|---------------|------------------------------------------------------------------|
| ISCO (Static location) | $3R_s$           | 0             | $\frac{1}{3}$ | $\frac{1}{3}$ | Pure coordinate location, $r = R_s/\kappa^2$                     |
| ISCO (Dynamic state)   | $3R_s$           | $\frac{1}{6}$ | $\frac{1}{3}$ | $\frac{1}{2}$ | Marginal orbital stability, $Q^2 = Q_Y^2 = 1/2$                  |
| Photon sphere (Static) | $\frac{3}{2}R_s$ | 0             | $\frac{2}{3}$ | $\frac{2}{3}$ | Pure coordinate location                                         |
| Causal Dynamic horizon | $\frac{3}{2}R_s$ | $\frac{1}{3}$ | $\frac{2}{3}$ | 1             | Relational horizon under closure $\kappa^2 = 2\beta^2$ , $Q = 1$ |
| Causal Static horizon  | $R_s$            | 0             | 1             | 1             | Purely gravitational closure limit, $\kappa^2 = 1$               |
| Dynamic horizon        | $R_s$            | $\frac{1}{2}$ | 1             | $\frac{3}{2}$ | Orbital closure at Schwarzschild radius                          |
| Extremal Kerr horizon  | $\frac{1}{2}R_s$ | 1             | 2             | 3             | Maximal rotation, $\beta = 1$ , merged horizons                  |

Table 2: Critical radii and their projectional parameters. Static coordinate locations ( $\beta = 0$ ) are strictly distinguished from their corresponding dynamic orbital states that emerge from the closure law  $\kappa^2 = 2\beta^2$ .



## 2 Closed Algebraical System of R.O.M. Equations

This section presents the closed algebraic system of Relational Orbital Mechanics (R.O.M.). Derivations of all quantities are traceable back to [Methodological Principles](#) and two relational primitives ( $\beta$  on  $S^1$ ,  $\kappa$  on  $S^2$ ) linked by the [Closure Theorem](#)  $\kappa^2 = 2\beta^2$ . The first block gives global invariants fixed for an entire orbit; subsequent blocks give phase-dependent quantities. Everything is expressed in direct observables (observed ratios, spectroscopic, chronometric and astrometric readings) so that no background metric is required.

[ROM\\_FULL\\_TEST.ipynb](#) - In this notebook we perform 3 wide tests of Relational Orbital Mechanics:

- Algebraically test the closed system (internal consistency of every multi-form identity). All 66 expressions (50-digit, random c,G,M,a,e,O).
- R.O.M. vs GR: We compute the exact Schwarzschild geodesic results by high-precision numerical integration (no approximation), then extract their Taylor coefficients in the SAME small parameter and compare term-by-term to R.O.M..
- ROM reconstructs  $a$ ,  $R_s$ ,  $r_p$ ,  $r_a$  and the whole phase-resolved orbit, and recovers GM (a length<sup>3</sup>/ time<sup>2</sup> combination, no G) and finally M (kg, G used ONLY here as a units conversion factor).

**Remark 2.1.** *R.O.M. instead of postulating that body moves under forces through preexisting container-like manifold; it derives unique, algebraically allowed relational states of a system. Without need to postulate additional decrease of freedom, system has no use for variational and least action principles. It is not equations of motion but algebraically closed system of allowed states within relational system.*

Test it yourself

DESMOS: <https://www.desmos.com/calculator/gr7qyzicf> interactive user-friendly R.O.M. solver.

$$\kappa^2 = 1 - \frac{1}{(1+z_\kappa)^2} \quad z_\kappa = \text{gravitational redshift}$$

$$\beta^2 = 1 - \frac{1}{(1+z_\beta)^2} \quad z_\beta = \text{transverse Doppler shift}$$

## Observational Inputs

$Z_{sys} = (1+z_\kappa)(1+z_\beta) = \frac{1}{\tau}$  (product of gravitational red shift and transverse Doppler shift)

$\tau = \kappa_X \cdot \beta_Y = \frac{1}{Z_{sys}}$  (product of projectinal phase factors on  $S^1$  and  $S^2$  carriers)

$z_\kappa = \frac{1}{\kappa_X} - 1$  (gravitational redshift)

$z_\beta = \frac{1}{\beta_Y} - 1$  (transverse Doppler shift)

$z_{\kappa o} = \frac{1}{\kappa_{X_o}} - 1$  (redshift at phase O)

$z_{\beta o} = \frac{1}{\beta_{Y_o}} - 1$  (transverse Doppler shift at phase O)

## Global System Parameters (Fixed for the Orbit)

$$\kappa = \sin(\theta_2) = \beta\sqrt{2} = \sqrt{1 - (1+z_\kappa)^{-2}} = \sqrt{\frac{R_s}{a}} = \sqrt{\frac{\rho}{\rho_{max}}} = \sqrt{\kappa_p^2(1-e)} = \sqrt{2(\kappa_o^2 - \beta_o^2)} = \sqrt{\frac{1}{2}(3 - \sqrt{1 + 8\tau_o(B_a)^2})} =$$

$$\kappa_R^2 \left(\frac{T_P}{T}\right)^{\frac{2}{3}} = \sqrt{\kappa_o^2 \cdot \eta_o} \text{ (potential projection at semi-major axis)}$$

$$\kappa_X = \cos(\theta_2) = \sqrt{1 - \kappa^2} = \frac{1}{(1+z_\kappa)} \text{ (potential phase factor)}$$

$$\kappa_R = \sqrt{\frac{\kappa^2}{\sin(\theta_R)}} = \sqrt{\frac{R_s}{\frac{T}{2\pi} c \beta \sin(\theta_R)}} \text{ (potential projection at the surface of central body)}$$

$$\beta = \cos(\theta_1) = \sqrt{1 - (1+z_\beta)^{-2}} = \frac{\kappa}{\sqrt{2}} = \beta_p \sqrt{\frac{1-e}{1+e}} = \frac{2\pi a}{Tc} = \left(\frac{\pi R_s}{Tc}\right)^{\frac{1}{3}} = \sqrt{(\kappa_o^2 - \beta_o^2)} = \sqrt{\frac{\kappa_p^2}{2}(1-e)} = \beta_o \frac{\sqrt{1-e^2}}{\sqrt{1+e^2+2e \cdot \cos(O)}} =$$

$$\sqrt{\beta_{sys}^2 \cdot \sin(\theta_{sys})} \text{ (Global kinetic projection (Invariant transverse baseline defining the system's circular equilibrium state and semi major axis))}$$

$$\beta_Y = \sin(\theta_1) = \sqrt{1 - \beta^2} \text{ (kinetic phase factor)}$$

$$\beta_{int} = \frac{\beta}{\sqrt{1-e^2}} \text{ (interior kinetic projection)}$$

$$\tau = \kappa_X \beta_Y \text{ (relational spacetime factor)}$$

$$\tau_Y = \sqrt{1 - \tau^2} = \sqrt{3\beta^2 - 2\beta^4} \text{ (relational spacetime phase factor)}$$

$$Q = \sqrt{\kappa^2 + \beta^2} = \sqrt{\frac{3}{2}} \kappa = \sqrt{3} \beta \text{ (total relational shift (magnitude of state difference))}$$

$$R_s = \kappa_o^2 r_o = \frac{Tc}{\pi} \beta^3 = \frac{8\pi^2 a^3}{T^2 c^2} = \frac{\Delta_{to}}{\zeta(O)} \kappa^2 \beta c = \frac{r_1 r_2}{r_2 - r_1} (\beta_1^2 - \beta_2^2) = \frac{a}{2} (3 - \sqrt{1 + 8\tau_o(B_a)^2}) = \frac{r_o}{2(2a - r_o)} (4a - r_o - ((4a - r_o)^2 - 8a(2a - r_o)(1 - \tau_o^2))^{\frac{1}{2}}) = \frac{2GM}{c^2} \text{ (Schwarzschild radius - system scale)}$$

$$a = \frac{R_s}{\kappa^2} = \frac{T\beta c}{2\pi} = \frac{\beta_o c}{\omega} \cdot \frac{\sqrt{1-e^2}}{\sqrt{1+e^2+2e \cos(O)}} = \frac{\Delta_{to}(O)}{\zeta} \beta c = \frac{Tc\beta^3}{2\pi(\frac{K_i}{\sin(i)})^2(1-e^2)} = \frac{\sqrt{1-e^2}}{2\pi \sin(i)} K_i Tc = \frac{R_{sys}}{\sin(\theta_{sys})} \text{ (semi-major axis)}$$

$$R_{sur} = a \cdot \sin(\theta_{sur}) \text{ (system's central body's physical radius)}$$

$$M = \frac{Tc^3\beta^3}{2\pi G} = \frac{\kappa^2 c^2 a}{2G} = 4\pi \rho a^3 \text{ (mass parameter)}$$

$$\begin{aligned}
t &= \frac{a}{c} \text{ (temporal scale of the system)} \\
\beta^2 &= (\kappa_o^2 - \beta_o^2) = \frac{\kappa_p^2}{2}(1 - e) = (\kappa_o^2 - \frac{\kappa_o^2}{2\delta_o}) = \frac{R_s}{2a} = (\frac{R_s}{T \cdot c} \pi)^{\frac{2}{3}} \text{ (energy invariant - binding energy)} \\
\Delta\phi &= \frac{2\pi \cdot \tau_Y^2}{1 - e^2} = \frac{2\pi(3\beta^2 - 2\beta^4)}{1 - e^2} = \frac{3\pi R_s}{a(1 - e^2)} - \frac{\pi R_s^2}{a^2(1 - e^2)} \text{ (precession of perihelion per orbit)} \\
\Omega_\phi &= \frac{\tau_Y^2}{1 - e^2} = \frac{3\beta^2 - 2\beta^4}{1 - e^2} \text{ (precession of perihelion per phase radian)} \\
\Omega &= 1 - \Omega_\phi = \frac{O}{o} \text{ (constant ratio between idealised phase o and true precessing phase O)} \\
h_W &= \frac{\kappa_o^2}{\kappa_o^2} \cdot \beta_T \cdot a \cdot c = r_o \cdot \beta_T \cdot c = r_o^2 \cdot \omega_o = a \cdot \beta c \cdot e_Y = R_s c \frac{\sqrt{1 - e^2}}{2\beta} \text{ (invariant angular momentum)} \\
\omega &= \frac{\beta c}{a} = \frac{c}{R_s} 2\beta^3 \text{ (angular frequency)} \\
T &= \frac{2\pi}{\omega} = \frac{2\pi a}{\beta c} = \frac{\pi R_s}{\beta^3 c} = \frac{2\pi \sqrt{2} R_s}{\kappa^3 c} = \frac{2\pi \Delta_{to}}{\zeta_o} \text{ (orbital period)} \\
\tau_D &= T \sin(\theta_{sys})^{\frac{3}{2}} = \frac{2\pi R_{sur}}{\beta_{sur} c} = \sqrt{\frac{\pi}{G\rho}} \text{ (Time-density invariant where } R_{ser} \text{ and } \beta_{ser} \text{ determined by the distance from center to the surface of central body and } \theta_{sys} = \text{angular radius)} \\
\Delta\varphi &= 2 \arcsin\left(\frac{\kappa_p^2(1 + \beta_p^2)}{2\beta_p^2 - \kappa_p^2(1 + \beta_p^2)}\right) \text{ (gravitational deflection)} \\
\Delta\gamma &= 2 \arcsin\left(\frac{\kappa_p}{\kappa_{Xp}}\right) \text{ (light deflection)} \\
\rho &= \frac{\kappa^2 \cdot c^2}{8 \cdot \pi \cdot G \cdot a^2} \text{ (relational density)} \\
\rho_{max} &= \frac{c^2}{8 \cdot \pi \cdot G \cdot a^2} \text{ (maximal density at the horizon (for non rotating systems), } \kappa^2 = 1 \equiv r = R_s) \text{, density reaches its natural bound)}
\end{aligned}$$

## Eccentricity Relations

$$\begin{aligned}
e &= \frac{1}{\delta_p} - 1 = 1 - \frac{2\beta_a^2}{\kappa_a^2} = \frac{2\beta_p^2}{\kappa_p^2} - 1 = \frac{r_a - r_p}{r_p + r_a} = 1 - \eta_o(0) = \eta_o(\pi) - 1 \text{ (eccentricity derived from closure)} \\
e_Y &= \sqrt{1 - e^2} \text{ (eccentricity's orthogonal value)} \\
e_X &= \frac{1 + e}{1 - e} = \frac{r_a}{r_p} = \frac{\delta_a}{\delta_p} = \frac{\beta_p}{\beta_a} = \frac{\kappa_p^2}{\kappa_a^2} = \frac{\kappa_a^2 \beta_p^2}{\kappa_p^2 \beta_a^2} \text{ (shape factor, still looks like magic to me)}
\end{aligned}$$

## Time-phase

$$\begin{aligned}
\omega_o &= \frac{\beta \cdot c}{a} \cdot \frac{(1 + e \cdot \cos(O))^2}{(1 - e^2)^{\frac{3}{2}}} \text{ (angular frequency at phase o)} \\
\Delta_{to} &= \int_0^O \frac{1}{\omega_\theta(\theta)} d\theta = \frac{a}{\beta c} \zeta = \frac{T}{2\pi} \zeta = \frac{R_s \zeta}{2\beta^3 c} \text{ (time duration of given phase interval)} \\
\zeta &= \frac{2\pi \Delta_t}{T} = \frac{\Delta_t c}{R_s} 2\beta^3 = (1 - e^2)^{\frac{3}{2}} \cdot \left( \int_0^O (1 + e \cdot \cos(\theta))^{-2} d\theta \right) = \frac{1}{\sqrt{1 - e^2}} \int_0^O \left( \frac{t_o(\theta)}{t} \right)^2 d\theta \text{ (Temporal Phase interval (mean anomaly analog))} \\
\zeta(< \pi) &= 2 \arctan\left(\frac{\sin(\frac{O}{2})}{e^{\frac{1}{2}} \cos(\frac{O}{2})}\right) - \frac{e \cdot e_Y \sin(O)}{1 + e \cos(O)} \text{ (Temporal Phase interval on ascending orbit from 0 to } \pi) \\
\zeta(> \pi) &= 2 \arctan\left(\frac{\sin(\frac{O}{2})}{e^{\frac{1}{2}} \cos(\frac{O}{2})}\right) - \frac{e \cdot e_Y \sin(O)}{1 + e \cos(O)} + 2\pi \text{ (Temporal Phase interval on descending orbit from } \pi \text{ to } 2\pi)
\end{aligned}$$

## Perihelion Relations

$$\begin{aligned}
r_p &= a(1 - e) = \frac{R_s}{\kappa_p^2} = \sqrt{((1 - e) \cdot a \cdot \cos(O))^2 + ((1 - e) \cdot a \cdot \sin(O))^2} \text{ (radius at perihelion)} \\
\kappa_p &= \kappa \sqrt{\frac{1}{1 - e}} = \sqrt{\frac{2\beta^2}{1 - e}} = Q_p \sqrt{\frac{2}{3 + e}} \text{ (potential projection at perihelion)} \\
\kappa_{Xp} &= \sqrt{1 - \kappa_p^2} \text{ (potential phase projection at perihelion)} \\
\beta_p &= \frac{V_p}{c} = \beta \sqrt{\frac{1 + e}{1 - e}} = \frac{r_p \omega_o(0)}{c} = \frac{\kappa_p}{\sqrt{2}} \sqrt{1 + e} \text{ (kinetic projection at perihelion)} \\
\delta_p &= \frac{\kappa_p^2}{2\beta_p^2} = \frac{1}{1 + e} \text{ (closure factor at perihelion)} \\
Q_p &= \sqrt{\kappa_p^2 + \beta_p^2} \text{ (relational shift at perihelion)}
\end{aligned}$$

## Aphelion Relations

$$\begin{aligned}
r_a &= a(1 + e) = \frac{R_s}{\kappa_a^2} \text{ (radius at aphelion)} \\
\beta_a &= \sqrt{\beta_p^2 e_X^{-2}} = \beta \sqrt{e_X^{-1}} \text{ (kinetic projection at aphelion)} \\
\kappa_a &= \sqrt{\beta^2 + \beta_a^2} \text{ (potential projection at aphelion)} \\
\delta_a &= \frac{1}{1 - e} = \frac{\kappa_a^2}{2\beta_a^2} \text{ (closure factor at aphelion)} \\
Q_a &= \sqrt{\kappa_a^2 + \beta_a^2} \text{ (relational shift at aphelion)}
\end{aligned}$$

## Phase Variables (Depend on phases $o$ and $O$ )

$o = \frac{O}{\Omega}$  (idealised orbital phase in radians, represent phase  $o$  progression without orbital precession)

$O = \Omega \cdot o = o - \Omega_\phi o$  (true orbital phase with precession)

$r = r_o(O) = a \frac{1-e^2}{1+e \cos(O)} = \frac{R_s}{\kappa_o^2} = a\eta_o$  (radial distance at phase  $O$ )

$\eta_o = \frac{1-e^2}{1+e \cos(O)} = \frac{r_o(O)}{a} = \frac{R_s}{\kappa_o(O)^2 a} = 2 - \frac{2\beta_o^2}{\kappa_o^2}$  (normalised true radial distance at phase  $O$ )  $\kappa_o = \sqrt{\beta_R(O)^2 + \beta_T(O)^2 + \beta^2} =$

$\sqrt{\frac{R_s}{r_o}} = \sqrt{1 - (1 + z_{\kappa_o}(O))^{-2}} = \kappa_p \sqrt{\frac{1+e \cos(O)}{1+e}} = \sqrt{\beta^2 + \beta_o^2} = \sqrt{\kappa_{surface}^2 \cdot \sin(\theta_o)}$  (local potential at the true phase  $O$ )

$\kappa_{X_o} = \sqrt{1 - \kappa_o^2} = (z_{\kappa_o}(O) + 1)^{-1}$  (gravitational phase factor at phase  $O$ )

$\beta_o = \sqrt{\beta_R^2 + \beta_T^2} = \beta \cdot \frac{\sqrt{1+e^2+2 \cdot e \cdot \cos(O)}}{\sqrt{1-e^2}} = \sqrt{\kappa_o^2 - \beta^2} = \frac{\kappa_o(O)}{\sqrt{2\delta_o(O)}} = \sqrt{\frac{\kappa_o(O)^2}{2} \frac{1+e^2+2 \cdot e \cdot \cos(O)}{1+e \cdot \cos(O)}} = \sqrt{\frac{R_s}{r_o(O)} - \frac{R_s}{2a}} = \sqrt{1 - (1 + z_{\beta_o}(O))^{-2}}$

(Instantaneous translational kinetic projection at phase  $O$ )

$\beta_R = \beta \frac{e \sin(O)}{\sqrt{1-e^2}} = \sqrt{\beta_o(O)^2 - \beta_T(O)^2} = \sqrt{((1 - (1 + z_{\kappa_o}(O))^{-2}) - \beta^2) - \frac{r_o(O)^2 \omega_o(O)^2}{c^2}}$  (Instantaneous radial kinetic projection (Line-of-sight speed) at phase  $O$ )

$\beta_T = \frac{r_o(O) \omega_o(O)}{c} = \beta \frac{1+e \cos(O)}{\sqrt{1-e^2}} = \kappa_o^2 \frac{\sqrt{1-e^2}}{2\beta} = \frac{R_s \sqrt{1-e^2}}{r_o(O) 2\beta}$  (Instantaneous transverse kinetic projection (Orthogonal sideways speed) at phase  $O$ )

$\beta_{Y_o} = \sqrt{1 - \beta_o^2} = (z_{\beta_o}(O) + 1)^{-1}$  (relativistic phase factor at phase  $O$ )

$\delta_o = \frac{1+e \cos o}{1+e^2+2e \cos o} = \frac{\kappa_o^2}{2\beta_o^2}$  (local closure factor at phase  $O$ )

$Q_o = \sqrt{\kappa_o^2 + \beta_o^2}$  (local relational shift at phase  $O$ )

$\omega_o = a\beta c \frac{e_Y}{r_o} = \frac{\beta c}{a} \frac{(1+e \cos o)^2}{(1-e^2)^{3/2}}$  (angular speed at phase  $O$ )

$\tau_o = \kappa_{X_o}(O) \cdot \beta_{Y_o}(O) = Z_{sys}(O)^{-1}$  (phase spacetime factor at phase  $O$ )

$\tau_{Y_o} = \sqrt{1 - \tau_o^2}$  (amplitude spacetime factor at phase  $O$ )

$t_o = \frac{r_o}{c}$  (light time scale at phase)

## Orbital Phase Invariants

### Holographic Decryption Invariant

$$Z_{raw}(-\omega_i)\tau(-\omega_i)(1 - e \cos \omega_i) + Z_{raw}(\pi - \omega_i)\tau(\pi - \omega_i)(1 + e \cos \omega_i) = 2 \quad (1)$$

$$\Delta E_{A \rightarrow B} + \Delta E_{B \rightarrow A} = 0$$

$$\beta^2 = (\kappa_o^2 - \beta_o^2)$$

$$h_W = r_o^2 \cdot \omega_o$$

$$\frac{\beta_T(O)}{\kappa_o^2(O)} = \frac{\sqrt{1-e^2}}{2\beta}$$

$$\beta_T(O)^2 \eta_o^2 = \beta^2(1 - e^2)$$

$$B_a - \zeta(B_a) = \zeta(B_d) - B_d$$

$$\tau_D = T \sin(\theta_{sur})^{\frac{3}{2}} = \frac{2\pi R_{sur}}{\beta_{sur} c} = \sqrt{\frac{\pi}{G\rho}}$$

$$\beta^2 - \beta_o^2 = \kappa^2 - \kappa_o^2$$

## Relational Geometry (WILL)

$\theta_1 = \arccos(\beta)$  (distribution angle on  $S^1$ , non-physical)

$\theta_2 = \arcsin(\kappa)$  (distribution angle on  $S^2$ , non-physical)

$\kappa^2 = 2\beta^2$  (Closure Theorem)

$\Delta E_{A \rightarrow B} + \Delta E_{B \rightarrow A} = 0$   $\Delta E_{A \rightarrow B} = E_0(\frac{\kappa_{X,B}}{\beta_{Y,B}} - \frac{\kappa_{X,A}}{\beta_{Y,A}})$ ;  $\Delta E_{B \rightarrow A} = E_0(\frac{\kappa_{X,A}}{\beta_{Y,A}} - \frac{\kappa_{X,B}}{\beta_{Y,B}})$  (Energy Symmetry Law is the universal relational energy ledger, the origin of causality)

$\Delta Q = Q_o^2 - Q^2$  (phase relational shift amplitude at phase  $O$ )

$B_a = \arccos(1 - \delta^{-1}) = \arccos(-e) = \arccos(1 - \frac{2\beta_p^2}{\kappa_p^2}) = \arccos(\frac{2\beta_a^2}{\kappa_a^2} - 1)$  (ascending orbital balance point where  $a = r$  and  $\kappa_o^2 = 2\beta_o^2$  are true)

$B_d = 2\pi - \arccos(-e)$  (descending orbital balance point where  $a = r$  and  $\kappa_o^2 = 2\beta_o^2$  are true)

## Structural-Dynamical Equivalence

$\frac{r_a}{r_p} = \frac{\delta_a}{\delta_p} = \frac{1+e}{1-e} = \frac{\beta_p}{\beta_a} = \frac{\kappa_p^2}{\kappa_a^2} = \frac{\kappa_a^2 \beta_p^2}{\kappa_p^2 \beta_a^2}$  (remarkable equivalence of structural ( $\kappa$  on  $S^2$  carrier) and dynamical ( $\beta$  on  $S^1$  carrier) symmetry suggesting once again that  $SPACETIME \equiv ENERGY$ )



## Observer dependant

$\theta_{sur}(r) = \arcsin(\frac{R_{sur}}{r})$  (angular radius of central body viewed from distance  $r$ )

$Z_{raw}(O) = (1 + \beta_{int}(\cos(O + \omega_i) + e \cos(\omega_i)) \sin(i)) Z_{sys}(O)$  (raw light shift including the line of site Doppler at phase  $o$ )

$\beta_z(O) = \frac{\beta}{\sqrt{1-e^2}}(\cos(O + \omega_i) + e \cos(\omega_i)) \sin(i)$  (line of site light shift)

$i$  = (orbital inclination in relation to line of site and orbital plane)

$\omega_i$  = (phase turn or argument of periapsis)

$K_i = \frac{\beta}{\sqrt{1-e^2}} \sin(i) = \beta_{int} \sin(i)$  (semi-amplitude invariant)

$Z_{rawmax} = Z_{sys}(-\omega_i)(1 + K_i(1 + e \cos \omega_i))$

$Z_{rawmin} = Z_{sys}(\pi - \omega_i)(1 + K_i(-1 + e \cos \omega_i))$

$B_{ai} = (B_a + \omega_i)$

$Z_{sys}(-\omega_i) = (1 - \beta^2 \frac{1+e^2+2e \cos(\omega_i)}{1-e^2})^{-\frac{1}{2}} (1 - 2\beta^2 \frac{1+e \cos(\omega_i)}{1-e^2})^{-\frac{1}{2}}$

$Z_{sys}(\pi - \omega_i) = (1 - \beta^2 \frac{1+e^2-2e \cos(\omega_i)}{1-e^2})^{-\frac{1}{2}} (1 - 2\beta^2 \frac{1-e \cos(\omega_i)}{1-e^2})^{-\frac{1}{2}}$

$Z_{raw}(-\omega_i)\tau(-\omega_i)(1 - e \cos \omega_i) + Z_{raw}(\pi - \omega_i)\tau(\pi - \omega_i)(1 + e \cos \omega_i) = 2$  (Holographic Decryption Invariant)

## Background Free Parametric Equations for the Observed Orbit

$$\begin{bmatrix} x_{sky} \\ y_{sky} \\ z_{depth} \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & \cos(i) & -\sin(i) \\ 0 & \sin(i) & \cos(i) \end{bmatrix} \begin{bmatrix} x_{orb} \\ y_{orb} \\ 0 \end{bmatrix}$$

$$x_{sky} = r(O) \cos(O + \omega_i)$$

$$y_{sky} = r(O) \sin(O + \omega_i) \cos(i)$$

$$z_{depth} = r(O) \sin(O + \omega_i) \sin(i)$$

## Un-tilted 2D coordinates within the orbital plane

$$x_{orb} = r(O) \cos(O + \omega_i)$$

$$y_{orb} = r(O) \sin(O + \omega_i)$$

### 2.1 Translation to Legacy Tensor Parameters

The continuous multi-component tensor fields of General Relativity are derived as localized matrix representations of the purely scalar relational projections  $(\beta, \kappa)$  on the  $S^1$  and  $S^2$  carriers. The pseudo-Riemannian manifold is not a primitive; it is the coordinate-inflated shadow of relational phase capacity.

**1. Metric Tensor** ( $g_{\mu\nu}$ ) The 4D metric tensor maps directly to the operational phase capacities and derived coordinate ratios.

- **Temporal Component:**  $g_{tt} \hat{=} -\kappa_X^2 c^2 = -c^2(1 - \kappa^2)$
- **Radial Spatial Component:**  $g_{rr} \hat{=} \kappa_X^{-2} = (1 - \kappa^2)^{-1}$
- **Transverse Spatial Components:**  $g_{\theta\theta} \hat{=} r^2$  and  $g_{\phi\phi} \hat{=} r^2 \sin^2 \theta$  (where  $r = R_s/\kappa^2$ )

**2. Four-Velocity** ( $U^\mu$ ) **and Momentum** ( $P^\mu$ ) The 4-vector velocity translates to the composite of the local energy scale ( $\tau$ ) and the separated kinematic amplitudes  $(\beta_R, \beta_T)$ .

- **Time Component:**  $U^t = \frac{dt}{d\tau} \hat{=} \frac{1}{\tau} = \frac{1}{\beta_Y \kappa_X}$
- **Spatial Components:**  $U^r \hat{=} c\beta_R U^t$  and  $U^\phi \hat{=} \frac{c}{r} \beta_T U^t$
- **Four-Momentum Norm:**  $P_\mu P^\mu = -(mc)^2 \hat{=} -E_0^2(\beta_Y^{-2} - \beta^2 \beta_Y^{-2}) = -E_0^2$  (Strict geometric closure of the  $S^1$  carrier).

**3. Stress-Energy Tensor** ( $T_{\mu\nu}$ ) The classical sources of the gravitational field map directly to the [relational density](#) saturation limit and the intrinsic [surface tension](#) derived.

- **Energy Density:**  $T_0^0 = \rho c^2 \hat{=} \rho_{\max} \kappa^2 c^2 = \frac{\kappa^2 c^4}{8\pi G r^2}$
- **Isotropic Pressure:**  $T_i^i = P \hat{=} -\rho c^2 = -\frac{\kappa^2 c^4}{8\pi G r^2}$  (The relational tension required for  $d\kappa^2/dr$  consistency).

**4. Ricci Tensor** ( $R_{\mu\nu}$ ), **Ricci Scalar** ( $R$ ), **and Einstein Tensor** ( $G_{\mu\nu}$ ) The differential curvature operators are bypassed entirely by the [Unified Geometric Field Equation](#) linking geometric scale directly to energy density ratios.

- **Vacuum** ( $R_{\mu\nu} = 0$ ):  $\hat{=} \frac{d}{dr}(r\kappa^2) = 0 \implies \kappa^2 = \frac{R_s}{r}$

- **Matter** ( $G_{\mu\nu} = \frac{8\pi G}{c^4} T_{\mu\nu}$ ):  $\triangleq \frac{d}{dr}(r\kappa^2) = \frac{8\pi G}{c^2} r^2 \rho_{matter}$

**5. Christoffel Symbols ( $\Gamma_{\mu\nu}^\lambda$ ) and Geodesic Integration** The affine connections ( $\Gamma_{\mu\nu}^\lambda$ ) used to define parallel transport on a curved manifold, the continuous integration of the geodesic equation ( $\frac{d^2 x^\mu}{d\tau^2} + \Gamma_{\alpha\beta}^\mu \frac{dx^\alpha}{d\tau} \frac{dx^\beta}{d\tau} = 0$ ) are bypassed in favor of the algebraic [Energy-Symmetry Law](#) ( $\Delta E_{A \rightarrow B} + \Delta E_{B \rightarrow A} = 0$ ) and the relational phase divergence  $\Delta\varphi = 2\pi \frac{\tau_Y^2}{1-e^2}$ .

### 3 Relational Origin of:

#### 3.1 Spectroscopic Shifts

**Theorem 3.1** (Spectroscopic Phase Shift). *The geometric projection  $\kappa^2$  of a closed system is determined by the operationally measurable gravitational redshift  $z_\kappa$  via the identity:*

$$\kappa^2 = 1 - \frac{1}{(1 + z_\kappa)^2}. \quad (2)$$

*Proof.* By the [Conservation of Relation Theorem](#) the gravitational amplitude  $\kappa$  and internal phase  $\kappa_X$  satisfy:

$$\kappa^2 + \kappa_X^2 = 1 \implies \kappa^2 = 1 - \kappa_X^2. \quad (3)$$

By [Relational Reciprocity Theorem](#), an observer must self-center at the relational origin  $(\beta, \kappa) = (0, 0)$ . Consequently, the observer's local phase is maximal:  $\kappa_{X(obs)} \equiv 1$ .

The observable spectroscopic shift  $z_\kappa$  is the operational measure of the discrepancy between the observer's internal phase and the source's internal phase. The optical scaling ratio is defined as:

$$1 + z_\kappa = \frac{\kappa_{X(obs)}}{\kappa_{X(source)}}. \quad (4)$$

Substituting the observer's maximal phase ( $\kappa_{X(obs)} = 1$ ), the exact phase of the source is isolated:

$$\kappa_{X(source)} = \frac{1}{1 + z_\kappa}. \quad (5)$$

Substituting this internal phase back into the initial closure invariant yields the exact algebraic identity:

$$\kappa^2 = 1 - \left(\frac{1}{1 + z_\kappa}\right)^2. \quad (6)$$

□

**Theorem 3.2** (Kinematic Phase Shift (Transverse Doppler)). *The kinematic geometric projection  $\beta^2$  of a closed system is strictly determined by the operationally measurable transverse Doppler shift  $z_\beta$  via the symmetric identity:*

$$\beta^2 = 1 - \frac{1}{(1 + z_\beta)^2}. \quad (7)$$

*Proof.* By the  $S^1$  closure invariant ([Conservation Theorem](#)), the kinematic amplitude  $\beta$  and internal kinematic phase  $\beta_Y$  satisfy:

$$\beta^2 + \beta_Y^2 = 1 \implies \beta^2 = 1 - \beta_Y^2. \quad (8)$$

Following the Principle of Relational Origin, the self-centered observer dictates the rest state, where external kinematic projection is zero ( $\beta_{obs} = 0$ ), yielding maximum internal kinematic phase:  $\beta_{Y(obs)} \equiv 1$ .

The transverse Doppler shift  $z_\beta$  (measured orthogonally to eliminate classical radial kinematics) is the pure operational measure of the discrepancy between the observer's internal kinematic phase and the source's phase. The optical scaling ratio is:

$$1 + z_\beta = \frac{\beta_{Y(obs)}}{\beta_{Y(source)}}. \quad (9)$$

Substituting  $\beta_{Y(obs)} = 1$ , we isolate the internal phase of the moving source:

$$\beta_{Y(source)} = \frac{1}{1 + z_\beta}. \quad (10)$$

Substituting this back into the  $S^1$  closure invariant yields the exact algebraic identity:

$$\beta^2 = 1 - \left(\frac{1}{1 + z_\beta}\right)^2. \quad (11)$$

□

**Theorem 3.3** (Operational Measurability). *The relational projections are encoded directly in the combined phase interactions of light (spectroscopy) and are operationally independent of  $G$  and  $M$ .*

*Proof. Step 1: The Raw Observable ( $\tau$ ).* Spectroscopy does not measure "gravitational potential" or "kinematic velocity" as separate isolates; it measures the total accumulated phase difference between the source and the observer. We define the **Relational Spacetime Factor**  $\tau$  as the inverse of the systems measured redshift product:

$$\kappa^2 = 1 - \frac{1}{(1 + z_\kappa)^2} \quad (z_\kappa = \text{gravitational redshift})$$

$$\beta^2 = 1 - \frac{1}{(1 + z_\beta)^2} \quad (z_\beta = \text{transverse Doppler shift})$$

$$\tau \equiv \frac{1}{Z_{sys}} = 1/[(1 + z_\kappa)(1 + z_\beta)]. \quad (12)$$

In the WILL framework, this single observable represents the product of the internal phase projections of the carriers  $S^2$  and  $S^1$ :

$$\tau = \underbrace{\kappa_X}_{\text{Gravitational Phase}} \cdot \underbrace{\beta_Y}_{\text{Kinematic Phase}} = \sqrt{1 - \kappa^2} \sqrt{1 - \beta^2}. \quad (13)$$

**Step 2: Exact Relational Identity.** We reject weak-field approximations. The exact relationship between the signal  $\tau$  and the structural norm  $Q$  is given by the algebraic expansion of the phase product:

$$\tau^2 = (1 - \kappa^2)(1 - \beta^2) = 1 - (\kappa^2 + \beta^2) + \kappa^2\beta^2. \quad (14)$$

Substituting  $Q^2 = \kappa^2 + \beta^2$ , we obtain the rigorous link between the optical observable and the geometric state:

$$\tau^2 = 1 - Q^2 + \kappa^2\beta^2. \quad (15)$$

This identity demonstrates that the optical signal contains the complete information about the system's structural state, measurable without prior knowledge of mass.  $\square$

### 3.2 Kepler's Third Law

The Keplerian proportionality  $a \propto T^{2/3}$  is a direct algebraic consequence of the energetic equilibrium between the 1-DOF directional kinematic carrier ( $S^1$ ) and the 2-DOF omnidirectional potential carrier ( $S^2$ ). The derivation strictly utilizes dimensionless ratios and absolute geometric scale, eliminating dependence on classical mass and the empirical gravitational constant.

The kinematic amplitude  $\beta$  on  $S^1$  defines ([sec:SR interval](#)) the ratio of directional motion to the universal limit  $c$ :

$$\beta = \frac{2\pi a}{cT} \quad (16)$$

The omnidirectional potential amplitude squared  $\kappa^2$  on  $S^2$  is defined ([sec:GR interval](#)) by the absolute spatial geometric scale invariant  $R_s$  and the radial distance  $a$ :

$$\kappa^2 = \frac{R_s}{a} \quad (17)$$

[Energetic closure](#) dictates the invariant geometric exchange rate between the carriers:

$$\kappa^2 = 2\beta^2 \quad (18)$$

Substituting the parameterizations into the closure condition yields:

$$\frac{R_s}{a} = 2\left(\frac{2\pi a}{cT}\right)^2 = \frac{8\pi^2 a^2}{c^2 T^2} \quad (19)$$

Rearranging to isolate the semi-major axis  $a$  gives the exact closed-form relation:

$$a^3 = \frac{R_s c^2}{8\pi^2} T^2 \quad (20)$$

$$a = \left(\frac{R_s c^2}{8\pi^2}\right)^{1/3} T^{2/3} \implies a \propto T^{2/3} \quad (21)$$

### 3.3 Eccentricity

Eccentricity is a measure of the projectional deviation from the circular equilibrium state ( $\delta = 1$ ).

**Theorem 3.4** (Geometric Eccentricity). *For a closed orbital system governed by the projection invariants of WILL Relational Geometry, the orbital eccentricity  $e$  is strictly determined by the closure factor at periapsis,  $\delta_p$ :*

$$e = \frac{2\beta_p^2}{\kappa_p^2} - 1 = \frac{1}{\delta_p} - 1. \quad (22)$$

*Proof.* Instead of relying on classical force laws, we derive this relation directly from the conservation of the two fundamental projection invariants of the WILL framework:

1. **Kinetic Projection Invariant (Binding Energy):**  $\beta^2 = \kappa_o^2(o) - \beta_o^2(o) = \text{const.}$
2. **Angular Projection Invariant:**  $h = r_o(o)\beta_T(o)c = \text{const}$  ( $\beta = \beta_T(o)$  at turning points).

Consider the two turning points of a closed orbit: periapsis ( $p$ ) and apoapsis ( $a$ ). By operational definition of the shape parameter  $e$ , the relation between radii is determined by the geometric range:

$$r_a = r_p \left( \frac{1+e}{1-e} \right). \quad (23)$$

**Step 1: Relational Mapping.** Using the angular invariant  $h$  (implying  $\beta \propto 1/r$ ) and the field definition  $\kappa^2 \propto 1/r$ , we express the apoapsis projections in terms of the periapsis values:

$$\beta_a^2 = [\beta_p \left( \frac{r_p}{r_a} \right)]^2 = \beta_p^2 \left( \frac{1-e}{1+e} \right)^2, \quad (24)$$

$$\kappa_a^2 = \kappa_p^2 \left( \frac{r_p}{r_a} \right) = \kappa_p^2 \left( \frac{1-e}{1+e} \right). \quad (25)$$

*Note: Kinematic projection scales quadratically with the radius ratio, while potential projection scales linearly.*

**Step 2: Energy Balance.** Substituting these into the kinetic global invariant  $\beta^2$ :

$$(\kappa_p^2 - \beta_p^2) = (\kappa_a^2 - \beta_a^2).$$

Substituting the mappings from Step 1:

$$\kappa_p^2 - \beta_p^2 = \kappa_p^2 \left( \frac{1-e}{1+e} \right) - \beta_p^2 \left( \frac{1-e}{1+e} \right)^2.$$

Rearranging to group potential terms ( $\kappa$ ) on the left and kinematic terms ( $\beta$ ) on the right:

$$\kappa_p^2 \left[ 1 - \frac{1-e}{1+e} \right] = \beta_p^2 \left[ 1 - \left( \frac{1-e}{1+e} \right)^2 \right].$$

**Step 3: Algebraic Reduction.** Expanding the terms in brackets:

$$\begin{aligned} \text{LHS bracket } (\kappa \text{ term}): \quad & 1 - \frac{1-e}{1+e} = \frac{(1+e) - (1-e)}{1+e} = \frac{2e}{1+e}. \\ \text{RHS bracket } (\beta \text{ term}): \quad & 1 - \frac{(1-e)^2}{(1+e)^2} = \frac{(1+e)^2 - (1-e)^2}{(1+e)^2} = \frac{4e}{(1+e)^2}. \end{aligned}$$

Substituting back into the balance equation:

$$\kappa_p^2 \left( \frac{2e}{1+e} \right) = \beta_p^2 \left( \frac{4e}{(1+e)^2} \right).$$

Dividing both sides by  $2e$  and multiplying by  $(1+e)^2$ :

$$\kappa_p^2(1+e) = 2\beta_p^2.$$

This yields geometric identity for bound orbits:

$$2\beta_p^2 = \kappa_p^2(1+e). \quad (26)$$

**Step 4: Connection to Closure.** Recall the definition of the closure factor at periapsis:

$$\delta_p = \frac{\kappa_p^2}{2\beta_p^2}.$$

Substituting Eq.(3.3) into this definition:

$$\delta_p = \frac{\kappa_p^2}{\kappa_p^2(1+e)} = \frac{1}{1+e}.$$

Solving for  $e$ , we obtain the stated result:

$$e = \frac{1}{\delta_p} - 1 = \frac{2\beta_p^2}{\kappa_p^2} - 1 = 1 - \frac{2\beta_a^2}{\kappa_a^2}$$

□

**Remark 3.5.** *This result confirms that eccentricity is strictly a measure of the energetic deviation from the circular equilibrium state ( $\delta = 1$ ), derived entirely from the conservation of relational projections without invoking mass or Newtonian forces.*

#### SUMMARY

$$\frac{2\beta_p^2}{\kappa_p^2} - 1 \equiv e \equiv 1 - \frac{2\beta_a^2}{\kappa_a^2}$$

ECCENTRICITY  $\equiv$  CLOSURE DEFECT

SPACETIME  $\equiv$  ENERGY

### 3.4 Vis-Viva

In elliptical systems, the [global closure theorem](#)  $\kappa^2 = 2\beta^2$  is conserved across the entire orbital cycle, while local phase variations manifest as an internal “breathing” of relational projections. We now derive the exact conservation law governing this local dynamic distribution.

**Proposition 3.6** (Global Kinetic Amplitude). *The difference between the square of the local potential projection  $\kappa_o^2(o)$  and the square of the local kinetic projection  $\beta_o^2(o)$  is a global binding invariant kinetic amplitude equal to  $\beta^2$ .*

*Proof.* The binding parameter  $\beta^2$  defines the global relational depth of the system:

$$\kappa^2 = 2\beta^2$$

From the definitions of the  $S^2$  and  $S^1$  projections at any arbitrary phase  $O$ :

$$\begin{aligned}\kappa_o^2(o) &= \frac{2\beta^2(1+e\cos o)}{1-e^2} \\ \beta_o^2(o) &= \frac{\beta^2(1+e^2+2e\cos o)}{1-e^2}\end{aligned}$$

Subtracting the local kinetic projection from the local potential projection:

$$\begin{aligned}\kappa_o^2(o) - \beta_o^2(o) &= \frac{2\beta^2 + 2\beta^2 e \cos o - (\beta^2 + \beta^2 e^2 + 2\beta^2 e \cos o)}{1-e^2} \\ \kappa_o^2(o) - \beta_o^2(o) &= \frac{\beta^2(1-e^2)}{1-e^2} = \beta^2\end{aligned}$$

Thus, the algebraic distance between the omnidirectional carrier  $S^2$  and directional carrier  $S^1$  relations remains globally constant irrespective of orbital phase and shape. □

**Theorem 3.7** (The Orthogonal Signature of the Orbit). *The square of the local potential projection  $\kappa_o^2(o)$  on the  $S^2$  carrier is the Pythagorean square sum of the local radial  $\beta_R^2(o)$ , transverse  $\beta_T^2(o)$  and the global  $\beta^2$  kinetic projections.*

*Proof.* The total local kinetic projection on the  $S^1$  carrier splits orthogonally into radial and transverse components within the orbital plane:

$$\beta_o^2(o) = \beta_R^2(o) + \beta_T^2(o)$$

Substituting this orthogonal decomposition into the phase-invariant relation  $\kappa_o^2(o) - \beta_o^2(o) = \beta^2$ :

$$\kappa_o^2(o) - (\beta_R^2(o) + \beta_T^2(o)) = \beta^2$$

Rearranging yields the three-dimensional algebraic relational closure:

$$\kappa_o^2(o) = \beta_R^2(o) + \beta_T^2(o) + \beta^2$$

□

**Remark 3.8** (Ontological Replacement of Vis-Viva). *This geometric invariant replaces the descriptive Newtonian Vis-Viva equation ( $v^2/2 - GM/r = -GM/2a$ ). Standard mechanics interprets this balance as a scalar subtraction of an abstract potential energy from kinetic energy. In WILL, it is revealed as a strict Pythagorean theorem of relational geometry: the local potential projection ( $\kappa_o^2$ ) is generated by the orthogonal summation of the kinetic “breathing” ( $\beta_R^2 + \beta_T^2$ ) and the global  $S^1$  projection ( $\beta^2$ ).  $SPACETIME \equiv ENERGY$ .*

### 3.5 Classical Acceleration

By the conservation of the global kinetic amplitude (Vis-Viva), the spatial gradients of the potential and kinetic projections balance to maintain the relational closure of the system. Taking the spatial derivative of the orbital invariant  $\kappa_o^2 - \beta_o^2 = \beta^2$  with respect to the radial distance  $r$  yields:

$$\frac{d}{dr}(\kappa_o^2) = \frac{d}{dr}(\beta_o^2) \quad (27)$$

Substituting the definitions  $\kappa_o^2 = \frac{R_s}{r}$  and  $\beta_o^2 = \frac{v^2}{c^2}$  into the gradient balance:

$$\frac{d}{dr}\left(\frac{R_s}{r}\right) = \frac{1}{c^2} \frac{d(v^2)}{dr} \quad (28)$$

Evaluating the derivatives provides the geometric relationship between the potential scale and the kinematic spatial variation:

$$-\frac{R_s}{r^2} = \frac{2v}{c^2} \frac{dv}{dr} \quad (29)$$

Classical mechanics introduces the external time differential  $dt$  to evaluate this spatial variation. Applying the chain rule  $a_{acc} = \frac{dv}{dt} = v \frac{dv}{dr}$  maps the kinematic spatial gradient to the classical acceleration ( $a_{acc}$ ):

$$-\frac{R_s}{r^2} = \frac{2}{c^2} a_{acc} \quad (30)$$

Isolating  $a_{acc}$  yields the classical Newtonian formula:

$$a_{acc} = -\frac{R_s c^2}{2r^2} = -\frac{GM}{r^2} \quad (31)$$

Within the system, energy conservation is required by the closure theorem ([sec:closure](#)) as an invariant resource of state change projected on kinetic  $S^1$  and potential  $S^2$  carriers. Thus, there is no need to introduce any new primitives such as force or spacetime curvature. Geometric conservation of relational energy states provides a complete operational and ontological explanation of observed phenomena.

### 3.6 Angular Momentum Conservation

The conservation of specific angular momentum  $h$  is derived as a phase-independent structural invariant of the relational balance.

**Proposition 3.9** (Invariant Ratio of Projections). *The transverse kinetic projection  $\beta_T(o)$  is strictly proportional to the local potential projection  $\kappa_o^2(o)$  scaled by the global system constants.*

*Proof.* From the definition of the local potential projection (based on [closure theorem](#)  $\kappa^2 = 2\beta^2$  and geometry of ellipse):

$$\kappa_o^2(o) = 2\beta^2 \frac{(1 + e \cos o)}{1 - e^2} \implies (1 + e \cos o) = \kappa_o^2(o) \frac{1 - e^2}{2\beta^2}$$

The transverse kinetic projection  $\beta_T(o)$  is defined on the  $S^1$  carrier as:

$$\beta_T(o) = \beta \frac{1 + e \cos o}{\sqrt{1 - e^2}}$$

Substituting the expression for  $(1 + e \cos o)$  into the definition of  $\beta_T(o)$ :

$$\beta_T(o) = \beta \left( \kappa_o^2(o) \frac{1 - e^2}{2\beta^2} \right) \frac{1}{\sqrt{1 - e^2}} = \kappa_o^2(o) \frac{\sqrt{1 - e^2}}{2\beta}$$

Rearranging to isolate the invariant ratio of the phase-dependent projections:

$$\frac{\beta_T(o)}{\kappa_o^2(o)} = \frac{\sqrt{1 - e^2}}{2\beta}$$

□

**Theorem 3.10** (Conservation of Angular Momentum). *The specific angular momentum  $h$ , defined as the product of the local scale  $r_o(o)$  and the transverse kinetic projection, is a global phase invariant.*

*Proof.* Using the local scale  $r_o(o) = R_s/\kappa_o^2(o)$ , we define  $h$ :

$$h = r_o(o)\beta_T(o)c = \frac{R_s}{\kappa_o^2(o)}\beta_T(o)c$$

Substituting the invariant ratio  $\frac{\beta_T(o)}{\kappa_o^2(o)} = \frac{\sqrt{1-e^2}}{2\beta}$ :

$$h = R_sc \frac{\sqrt{1-e^2}}{2\beta}$$

Since  $R_s$ ,  $e$ , and  $\beta$  are global invariants,  $h$  does not depend on phase  $o$ . Angular momentum is thus the manifestation of the fixed structural exchange rate between the  $S^2$  and the  $S^1$  carriers and there projections.  $\square$

### 3.7 Orbital Precession as Phase Accumulation

In standard General Relativity, orbital precession is derived via perturbative methods and Taylor expansions of the Schwarzschild metric. In WILL Relational Geometry, precession is derived strictly from the exact algebraic accumulation of the internal phase difference over a closed geometric cycle, without differential approximations.

**Theorem 3.11** (Precession Law). *The secular angular shift (precession)  $\Delta\varphi$  per closed orbit is the strict geometric projection of the system's relational phase difference  $\tau_Y$  normalized by the orbital shape factor  $(1 - e^2)$ .*

*Proof. Step 1: State Difference.* A system completely at rest with the observer possesses maximal internal phase  $\tau = 1$ . The true relational phase of an orbiting system is the product of the kinetic  $S^1$  and potential  $S^2$  projections carriers:

$$\tau^2 = \beta_Y^2 \kappa_X^2 = (1 - \beta^2)(1 - \kappa^2). \quad (32)$$

Expanding this product algebraically yields:

$$\tau^2 = 1 - (\beta^2 + \kappa^2) + \kappa^2\beta^2. \quad (33)$$

Recognizing the frame invariant scale of the relational shift  $Q^2 = \beta^2 + \kappa^2$ ; The total phase divergence from the rest origin ( $\tau = 1$ ) is defined as:

$$\tau_Y^2 \equiv 1 - \tau^2 = Q^2 - \kappa^2\beta^2. \quad (34)$$

The cross-coupling term  $\kappa^2\beta^2$  account for non-linear geometric interaction between the directional  $S^1$  and omnidirectional  $S^2$  relational carriers.

**Step 2: Geometric Accumulation of Phase Divergence.** The system accumulates this state mismatch over the full cycle ( $2\pi$ ). The total angular shift is the base phase scaled by the state divergence, normalized by the orbital shape factor  $(1 - e^2)$ .

Restricted by the [Mathematical Transparency principle](#) (pr:mathematical), we do not truncate or simplify this connection. The orbital precession is locked in its exact, fully non-linear algebraic form:

$$\Delta\varphi = \frac{2\pi \cdot \tau_Y^2}{1 - e^2} = \frac{2\pi(1 - \tau^2)}{1 - e^2} \quad (35)$$

$\square$

**Remark 3.12** (Epistemic Purity). *This derivation contrast with the complex integration of perturbed differential equations reviling the exact algebraic evaluation of the relational phase. The cross-coupling term  $\kappa^2\beta^2$  distinct RG from just approximation and structurally complete the exact invariant topology (full mathematical prove 8).*

### 3.8 Dynamic Event Horizon

In classical literature, the event horizon is often heuristically treated as an escape-velocity boundary where spacetime curvature becomes infinitely steep. In RG, the horizon emerges strictly as the limit of geometric precessional invariant, representing the ultimate topological closure of a trajectory.

Using the exact relational phase accumulation law (derived in 3.7), the baseline precession per orbit is given by:

$$\Delta\varphi = \frac{2\pi \cdot \tau_Y^2}{1 - e^2} = \frac{2\pi(1 - \tau^2)}{1 - e^2} \quad (36)$$

For a massive system under total [relational closure](#) ( $\kappa^2 = 2\beta^2$ ), consider the **Dynamic Horizon** state where the potential projection reaches absolute saturation ( $\kappa^2 = 1$ ). By the closure rule, the kinematic projection is  $\beta^2 = \frac{1}{2}$ .



At this limit, the gravitational phase component collapses ( $\kappa_X = \sqrt{1-1} = 0$ ), causing the entire internal causal structure of the system to vanish:

$$\tau^2 = (1 - \beta^2)(1 - \kappa^2) = 0 \implies \tau_Y = 1 \quad (37)$$

For a perfectly circular orbit ( $e = 0$ ) at this causal limit, the precession evaluates to:

$$\Delta\varphi = \frac{2\pi \cdot (1)}{1 - 0} = 2\pi \quad (38)$$

#### Event Horizon

An orbital precession of  $\Delta\varphi = 2\pi$  radians per revolution implies that the periapsis shifts by a full circle during a single orbit. Consequently, the orbital progression continuously folds onto itself perfectly, creating a closed, self-intersecting locus where forward orbital motion mathematically nullifies itself. This provides a novel interpretation of the event horizon phenomena not as a "point of infinite density" in an external container-like spacetime, but the unitary topological closure of the relational trajectory itself.

### 3.9 Gravitational Deflection and Lensing

In General Relativity, the deflection of light is obtained by integrating the null geodesic equations over a curved spacetime manifold, often relying on weak-field approximations and Taylor expansions. Within WILL Relational Geometry (RG), we reject both the background manifold and the use of mathematical approximations as non-operational ontological artifacts.

The system consists exclusively of its participants: the Source, the Lens (at periapsis  $p$ ), and the Receiver. The total deflection angle must be derived as a strict, exact algebraic difference between their measurable relational phase states, without resorting to series expansions.

#### Gravitational Deflection and Interaction Gradient

Energy projections distribution among axis of relational carriers has to obey the conservation law (distribution between axis does not create nor destroy energy). There must exist a strict, algebraically closed gradient connecting all states, governed entirely by the kinematic projection  $\beta$ .

**Theorem 3.13** (Unified Interaction Gradient). *The gravitational interaction capacity of any entity is determined by its available phase buffer  $\beta_Y$ . The geometric scaling factor  $\Gamma$  that dictates the distribution of the potential projection  $\kappa^2$  onto the spatial trajectory is strictly defined by the arithmetic mean of the saturated carrier  $S^1$ :*

$$\Gamma(\beta) = \frac{1 + \beta^2}{2} = 1 - \frac{\beta_Y^2}{2} \quad (39)$$

*Proof.* By the  $S^1$  closure invariant ( $\beta^2 + \beta_Y^2 = 1$ ), an object at rest ( $\beta = 0$ ) possesses maximum internal phase ( $\beta_Y = 1$ ). This phase acts as a geometric buffer, absorbing half of the relational gradient, yielding the classical partitioning  $\Gamma = \frac{1}{2}$ . As the spatial projection  $\beta$  increases, the internal clock slows, and the phase buffer  $\beta_Y$  depletes. At the topological limit  $\beta \rightarrow 1$  (light), the internal phase collapses ( $\beta_Y \rightarrow 0$ ). The buffer is exhausted, forcing the entity to absorb the full, unpartitioned gravitational gradient, yielding  $\Gamma = 1$ .  $\square$

Using this gradient, we define the **Unified Closure Defect** for any trajectory—from a slow asteroid to a photon—as:

$$\delta_\varphi = \frac{\kappa_p^2}{\beta_p^2} \Gamma(\beta_p) = \frac{\kappa_p^2(1 + \beta_p^2)}{2\beta_p^2} \quad (40)$$

The geometric shape parameter (eccentricity) of the trajectory is derived directly from this defect:

$$e_\varphi = \frac{1}{\delta_\varphi} - 1 = \frac{2\beta_p^2}{\kappa_p^2(1 + \beta_p^2)} - 1 \quad (41)$$

Applying the exact algebraic transit equation for a distant observer ( $\kappa_o \rightarrow 0$ ), we have  $\cos o_\infty = -1/e_\varphi$ . Using the same trigonometric extraction as established for light ( $\sin(\frac{\Delta\varphi}{2}) = \frac{1}{e_\varphi}$ ), we arrive at the absolute, unified equation for gravitational deflection:

$$\Delta\varphi = 2 \arcsin\left(\frac{\kappa_p^2(1 + \beta_p^2)}{2\beta_p^2 - \kappa_p^2(1 + \beta_p^2)}\right) \quad (42)$$

Solving for  $\beta_p = 1$  the total deflection angle  $\Delta\varphi$  gives non-linear equation for light deflection in RG:

$$\Delta\gamma = 2 \arcsin\left(\frac{\kappa_p^2}{\kappa_{Xp}^2}\right) \quad (43)$$



**Verification of Topological Limits:**

- **Newtonian Limit** ( $\beta_p \ll 1$ ): The phase buffer is full ( $\Gamma \rightarrow 0.5$ ). The eccentricity is dominated by  $2\beta_p^2/\kappa_p^2$ . The deflection reduces to the classical Rutherford/Newton scattering:  $\Delta\varphi \approx \frac{\kappa_p^2}{\beta_p^2}$ .
- **Relativistic Limit** ( $\beta_p \rightarrow 0.99$ ): The phase buffer is nearly depleted ( $\Gamma \rightarrow 0.99$ ). The trajectory stiffens, and the deflection angle approaches the photonic maximum, smoothly capturing the post-Newtonian factor without Taylor expansions.
- **Photonic Limit** ( $\beta_p = 1$ ): The phase buffer is completely exhausted ( $\Gamma = 1$ ). Substituting  $\beta_p = 1$  yields  $e = \frac{2}{\kappa_p^2(2)} - 1 = \frac{1}{\kappa_p^2} - 1 = \frac{\kappa_{Xp}^2}{\kappa_p^2}$ . The equation resolves perfectly into the exact light deflection identity:  $\Delta\varphi = 2 \arcsin(\frac{\kappa_p^2}{\kappa_{Xp}^2})$ .

**Summary**

This formulation suggests that light and massive bodies do not require separate ontological categories or distinct dynamical laws. In RG, the photonic state the topological limit ( $\beta \rightarrow 1, \beta_Y \rightarrow 0$ ) of the phase buffer. The historical factor of 2 in gravitational lensing might not be an anomaly of null geodesics in curved background spacetime, but the inevitable geometric consequence of this exhausted phase capacity."

**Gravitational Lensing and the Einstein Ring**

In Relational Geometry, the description of gravitational lensing (deflection, convergence, shear, and magnification) does not require a background metric or the postulation of a curved manifold. It arises entirely as the geometric projection of the deflection angle onto the observer's plane.

By employing the standard geometric baseline equation for the observer-lens-source system, the true angular position  $\theta_S$  of a source is related to its observed image position  $\theta_I$  via:

$$\theta_S = \theta_I - \frac{D_{LS}}{D_S} \Delta\varphi_{\text{unified}}(\beta_p, \kappa_p(\theta_I)) \quad (44)$$

where  $D_{LS}$  and  $D_S$  are the relational baseline distances.

The potential projection at periapsis  $\kappa_p^2$  is strictly a function of the observed angle  $\theta_I$ , defined by the physical impact parameter  $r_p = D_L \theta_I$ :

$$\kappa_p^2(\theta_I) = \frac{R_s}{D_L \theta_I} \quad (45)$$

**Theorem 3.14** (Exact Algebraic Einstein Ring). *For a perfectly aligned system ( $\theta_S = 0$ ), the image forms a symmetric ring. The exact angular radius of this Einstein ring,  $\theta_E$ , valid for all velocities and field strengths, is given by the implicit algebraic equation:*

$$\theta_E = 2 \frac{D_{LS}}{D_S} \arcsin\left(\frac{\kappa_p^2(\theta_E)(1 + \beta_p^2)}{2\beta_p^2 - \kappa_p^2(\theta_E)(1 + \beta_p^2)}\right) \quad (46)$$

**Recovery of Topological Limits:** To reveal the underlying gradient without altering the exact equation, we examine the weak-field astrophysical limit ( $\kappa_p^2 \ll \beta_p^2$ ), where  $\arcsin(x) \approx x$ . The unified deflection simplifies algebraically to:

$$\theta_E \approx \frac{D_{LS}}{D_S} \frac{R_s}{D_L \theta_E} \left(\frac{1 + \beta_p^2}{\beta_p^2}\right) \implies \theta_E^2 \approx \frac{R_s D_{LS}}{D_L D_S} \left(\frac{1 + \beta_p^2}{\beta_p^2}\right) \quad (47)$$

**Colab Notebook**

[Gravitational Lensing.ipynb](#)

- **The Photonic Limit** ( $\beta_p \rightarrow 1$ ): The kinematic factor evaluates exactly to  $\frac{1+1}{1} = 2$ . The equation resolves to  $\theta_{E,\gamma} = \sqrt{\frac{2R_s D_{LS}}{D_L D_S}}$ , matching General Relativistic prediction for light (equivalent to  $4GM/c^2$ ).
- **The Relativistic Massive Limit** ( $0 < \beta_p < 1$ ): The continuous phase-buffer depletion natively supplies the correct post-Newtonian scaling without external ad-hoc gradients.

The properties of the lens - convergence ( $\kappa_{\text{lens}}$ ), shear ( $\gamma$ ), and magnification ( $\mu$ ) - are the components of the 2D Jacobian matrix of the deflection field  $\vec{\alpha}(\vec{\theta})$ . Because the physics of the phase-buffer  $\beta_Y$  is fundamentally woven into the exact definition of  $\Delta\varphi_{\text{unified}}$ , no manual "weighting" of shear or convergence is required. The optical mapping emerges naturally from the pure algebraic closure of the  $S^1$  and  $S^2$  relational carriers.

### 3.10 Schwarzschild Metric

Schwarzschild metric interval is the coordinate-inflated projection of the potential phase  $\kappa_{Xo}$  and kinematic amplitudes  $\beta_R, \beta_T$ . The fundamental primitive is the dimensionless algebraic composition of the relational state. Metric recovered by appending substantival posits.

- (i) **A spatial container.** A background manifold is imposed to host the geometry.
- (ii) **Externally flowing time.** An autonomous coordinate time  $t$  is introduced.
- (iii) **Coordinate grid.** Kinematic projections are decomposed onto imposed spatial axes  $r$  and  $\phi$ , transforming dimensionless amplitudes into differential coordinate ratios.
- (iv) **Arbitrary scale reference.** The scalar composition is multiplied by an external magnitude  $c^2 dt^2$ .

The relational projections map to the posited coordinate functions:

- **Temporal potential phase:**  $\kappa_{Xo}(O)^2 = 1 - \kappa_o(O)^2 = 1 - \beta^2 - \beta_o^2 = 1 - \frac{R_s}{r}$
- **Radial kinematic amplitude:**  $\beta_R(O)^2 = \beta^2 \cdot \frac{e^2 \sin^2(O)}{1-e^2} = \left(\frac{dr}{cdt}\right)^2$
- **Transverse kinematic amplitude:**  $\beta_T(O)^2 = \beta^2 \cdot \frac{(1+e \cdot \cos(O))^2}{1-e^2} = \left(\frac{rd\phi}{cdt}\right)^2$

Substituting the coordinate functions into the relational composition yields:

$$\kappa_{Xo}(O)^2 - \frac{\beta_R(O)^2}{\kappa_{Xo}(O)^2} - \beta_T(O)^2 = \left(1 - \frac{R_s}{r}\right) - \frac{\left(\frac{dr}{cdt}\right)^2}{1 - \frac{R_s}{r}} - \left(\frac{rd\phi}{cdt}\right)^2 \quad (48)$$

This specific linear combination directly reproduces the three metric coefficients of the Schwarzschild interval under the coordinate identifications above. Applying scale reference  $c^2 dt^2$  to recover the Schwarzschild interval:

$$-ds^2 = c^2 \left(1 - \frac{R_s}{r}\right) dt^2 - \left(1 - \frac{R_s}{r}\right)^{-1} dr^2 - r^2 d\phi^2 \quad (49)$$

The 4D pseudo-Riemannian metric tensor  $g_{\mu\nu}$  carries more ontological structure than the relations it encodes. The irreducible primitive here remains the structure of dimensionless relational projections  $\kappa_X^2 + \kappa^2 = 1$ .

#### Colab Sympy Verification

[Schwarzschild metric sympy.ipynb](#)

## 4 Intrinsic Unitless Core: Relational Parameterization

The preceding derivations in Relational Orbital Mechanics (R.O.M.) have demonstrated that the entire system can be factorized into a dimensionless core and a single scale parameter. This section formalizes this unitless formulation, proving that physical laws operate independently of anthropocentric units, metrics, or dimensional conventions and constants, and subsequently validates these principles empirically through the precise derivation of solar system orbits and precession.

### 4.1 The Intrinsic Relational Core: Unitless Formulation

**The claim** is structural: the closed R.O.M. system factorizes into a *fully solved dimensionless core* and a *one-parameter scale map*. Units, and the constants  $c$  and  $G$  that service them, belong exclusively to the map. A civilisation with different rulers, clocks, calendars and mathematical conventions solves the identical core and obtains the identical pure numbers; only the final labelling differs. Physical laws operates before — and independently of — anthropocentric units, metrics and scales.

**Theorem 4.1** (Factorization of R.O.M.). *Every quantity  $X$  of the closed R.O.M. system (2) factorizes as*

$$X = \hat{X}(\beta, e; O; i, \omega_i; \kappa_R) \cdot S_X(R_s; c, G)$$

where  $\hat{X}$  is a closed-form pure number determined entirely by dimensionless relational inputs, and  $S_X$  is a monomial scale factor depending only on the quantity class:

| <i>Quantity class</i>                      | $S_X$   | <i>Quantity class</i>            | $S_X$               |
|--------------------------------------------|---------|----------------------------------|---------------------|
| <i>projections, shifts, angles, phases</i> | 1       | <i>frequency</i>                 | $c/R_s$             |
| <i>length</i>                              | $R_s$   | <i>specific angular momentum</i> | $R_s c$             |
| <i>time</i>                                | $R_s/c$ | <i>mass</i>                      | $R_s \cdot c^2/G$   |
| <i>velocity</i>                            | $c$     | <i>density</i>                   | $1/R_s \cdot c^2/G$ |

No dimensional constant enters any  $\hat{X}$ .

**Corollary 4.2** (Universal Geometric Mass). *In its own geometric units every bound system has the same mass,  $\hat{M} = \frac{1}{2}$ . Mass is not a property of the relational state at all: it is the scale itself,  $M = \frac{1}{2} c^2 R_s / G$  — the anchor  $R_s$  dressed in the kilogram convention. This sharpens the Absolute Scale Derivation section argument (6). **Epistemic Mandate: the "mass parameter" carries zero relational information beyond the system scale.***

**Remark 4.3.** *In this section we using  $(\beta, e)$  as the intrinsic input pair just as an example. The system is fully algebraically closed and any parameters can be used as an input.*

*The normalization also exposes the Universal Horizon Constant (5) as the pure number  $\hat{\tau}_{D(limit)} = 2\sqrt{2}\pi$  at  $\kappa_R \rightarrow 1$ .*

## 4.2 Observational Input Channels

The example intrinsic pair  $(\beta, e)$  is reachable through algebraically independent menus of direct observables. All entries are pure ratios — frequency ratios, angle ratios, angular-speed ratios, cycle-count ratios — identical for any observer regardless of local units ("cross-cultural invariants").

| <i>e</i> -channel (orbit shape) — any one:                  |                                                                                         |
|-------------------------------------------------------------|-----------------------------------------------------------------------------------------|
| E1 apparent angular speeds                                  | $e = \frac{\sqrt{\omega_{max}/\omega_{min}} - 1}{\sqrt{\omega_{max}/\omega_{min}} + 1}$ |
| E2 apsidal Doppler pair                                     | $e = \frac{\beta_p - \beta_a}{\beta_p + \beta_a}$                                       |
| E3 apsidal sky-angle ratio                                  | $e = \frac{r_a/r_p - 1}{r_a/r_p + 1}$                                                   |
| E4 balance-point phase                                      | $e = -\cos(B_a)$                                                                        |
| E5 projections only                                         | $e = \frac{2\beta_p^2}{\kappa_p^2} - 1 = 1 - \frac{2\beta_a^2}{\kappa_a^2}$             |
| <i><math>\beta</math></i> -channel (orbit depth) — any one: |                                                                                         |
| B1 transverse Doppler at $a$                                | $\beta^2 = 1 - (1 + z_\beta)^{-2}$                                                      |
| B2 gravitational redshift at $a$                            | $\beta^2 = \frac{1}{2}(1 - (1 + z_\kappa)^{-2})$                                        |
| B3 combined shift $Z_{sys} = 1/\tau$                        | $\beta^2 = \frac{1}{4}(3 - \sqrt{1 + 8 \cdot Z_{sys}(B_{a,d})^{-2}})$                   |
| B4 apsidal Doppler pair                                     | $\beta = \sqrt{\beta_p \beta_a}$                                                        |
| B5 two-point (vis-viva)                                     | $\beta^2 = \frac{(\beta_1^2 - \beta_2^2)(r_2/r_1)}{r_2/r_1 - 1} - \beta_1^2$            |
| B6 surface channel                                          | $\beta^2 = \frac{1}{2}(1 - (1 + z_{surf})^{-2}) \sin \theta_{obs}$                      |

Channels B3 and E4 exploit the balance points: at  $B_a = \arccos(-e)$  the measured light signal  $\tau(B_a)$  equals the global  $\tau$ , so a single spectroscopic reading there yields  $\kappa^2 = \frac{1}{2}(3 - \sqrt{1 + 8\tau^2})$  (6.2Balance Point Theorem) while the phase location itself yields  $e$ .

## 4.3 Operational Algorithm

**Stage 0 Measure pure ratios.** Select one  $e$ -channel and one  $\beta$ -channel observable set from the menus above (plus  $\kappa_R$  via  $z_{surf}$  if surface outputs are wanted). No unit is recorded.

**Stage 1 Invert to the intrinsic state**  $(\beta, e)$ .

**Stage 2 Solve the global block:**  $\kappa, \kappa_X, \beta_Y, \tau, \tau_Y, Q, \hat{a}, \hat{r}_p, \hat{r}_a, \kappa_p, \beta_p, \kappa_a, \beta_a, \delta_p, \delta_a, e_X, e_Y, \Delta\phi, \Omega, \hat{T}, \hat{\omega}, \hat{h}, B_a, B_d, \hat{\rho}, \hat{M} = \frac{1}{2}$  — all closed-form pure numbers.

**Stage 3 Phase-resolve** at any  $O$ :  $\eta_o, \hat{r}_o, \kappa_o, \beta_o, \beta_R, \beta_T, \delta_o, Q_o, \hat{\omega}_o, \tau_o, Z_{sys}(O), \zeta(O), \hat{\Delta}_{to}$ .

**Stage 4 Observer-project** with  $(i, \omega_i)$ :  $K_i, \beta_z, Z_{raw}$ , sky-plane parametric orbit — still pure numbers (sky lengths in units of  $\hat{a}$ ).

**Stage 5 (Optional) Inject one anchor.** A single dimensional reading — chronometric  $T$  giving  $R_s = Tc\beta^3/\pi$ , or astrometric  $r_p$  giving  $R_s = r_p\kappa_p^2$  — scales the entire solved system: lengths  $\times R_s$ , times  $\times R_s/c$ , frequencies  $\times c/R_s$ , mass  $= \frac{1}{2}c^2 R_s/G$ . The system was already solved at Stage 2; this stage only names the answers in local units.

**Corollary 4.4** (Cross-Cultural Invariance). *Two civilisations with incommensurable rulers and clocks conventions (the concept of mass isn't mandatory), observing the same bound system through any of the channels above, compute identical intrinsic states, identical  $\hat{X}$  for every quantity, and identical dimensionless predictions ( $\Delta\phi$  per orbit,  $Z_{sys}$ ,  $e$ , phase timings  $\zeta$ ). They disagree only on Stage 5 labels, and their labels are related by exact unit conversion. The physical content of R.O.M. is therefore invariant not merely under coordinate changes and scales, but under the choice of metrology itself.*

#### Test it yourself

COLAB: [ROM\\_UNITLESS\\_TEST.ipynb](#) — 50-digit verification: intrinsic identity battery, dimensional-vs-unitless commutation across 41 orders of magnitude in  $R_s$ , unit-system invariance including randomly generated alien units,  $c/G$  scrambling, ten observational routes converging on one state, and the Mercury/Jupiter parameterization executed unitlessly with the anchor injected last.

DESMOS: <https://www.desmos.com/calculator/gr7qykzicf> interactive user-friendly R.O.M. solver.

## 4.4 Relational Parameterization: Empirical Validation

**Theorem 4.5** (Relational Parameterization). *The complete structural and dynamical parameterization of any closed orbital system - specifically its geometric shape ( $e$  - eccentricity), secular phase shift ( $\Delta\varphi$  - orbital precession), horizon scale ( $R_s$  - Schwarzschild radius), and global scale ( $a$  - semi-major axis) - are strictly determined by the algebraic relations of its dimensionless spectroscopic and chronometric observables. The derivation requires no independent mass parameter ( $M$ ), gravitational constant ( $G$ ), or a priori spatial metric and deferential or tensor formalisms.*

**Direct algebraic derivation via spectroscopic and chronometric ratios.**

*Proof.* The proof is constructive and algebraically closed. It generalizes to any bound orbital system. We demonstrate this using the Mercury-Sun system as an operational test case and Jupiter-Sun as confirmation test. All governing equations are derived exclusively from the internal topology of the WILL relational carriers ( $S^1$ ,  $S^2$ ), strictly bypassing Newtonian mechanics, Special Relativity, or General Relativity. The derivation is executed using only four dimensionless, epistemologically direct observables ( $e, \theta, T_{ratio}, z_{sun}$ ):

#### INPUTS (Cross-Cultural Invariants)

The following inputs are strict dimensionless ratios. They represent direct physical observables that remain identical for any observer in the Universe, regardless of local unit systems (meters/seconds) or theoretical paradigms.

| Parameter                      | Value                   | Direct Observational Method                                                                                                                                                                                                                                         |
|--------------------------------|-------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Sun's physical observables     |                         |                                                                                                                                                                                                                                                                     |
| $\theta_{\odot}$               | 0.0046524 rad           | <b>Visual Size:</b> The angular radius of the central body in the sky. [? ]                                                                                                                                                                                         |
| $z_{sun}$                      | $2.1224 \times 10^{-6}$ | <b>Spectroscopic Tension:</b> The raw fractional frequency shift of light $(1 - \nu_{obs}/\nu_{emit})$ . Defined strictly as an optical geometric observable, independent of mass or Newtonian constants [? ].                                                      |
| Mercury's physical observables |                         |                                                                                                                                                                                                                                                                     |
| $T_{ratioM} = T_M/T_{\oplus}$  | 0.2408                  | <b>Cycle Ratio:</b> The observed system's full orbital period divided by the observer's local orbital period. [? ]                                                                                                                                                  |
| $e_M$                          | 0.2056                  | <b>Visual Kinematics:</b> Derived directly from the ratio of the planet's fastest ( $\omega_{max}$ ) and slowest ( $\omega_{min}$ ) apparent angular speeds: $e = \frac{\sqrt{\omega_{max}/\omega_{min}-1}}{\sqrt{\omega_{max}/\omega_{min}+1}}$ [? ].              |
| Jupiter's physical observables |                         |                                                                                                                                                                                                                                                                     |
| $T_{ratioJ} = T_J/T_{\oplus}$  | 11.862                  | <b>Cycle Ratio:</b> The full Jovian orbital period measured against the observer's local (Earth) orbital period — a purely temporal, clock-based observable requiring no unit conversion or gravitational constant [? ].                                            |
| $e_J$                          | 0.04839266              | <b>Visual Kinematics:</b> Derived directly from the ratio of Jupiter's fastest ( $\omega_{max}$ ) and slowest ( $\omega_{min}$ ) apparent angular speeds along its orbit: $e = \frac{\sqrt{\omega_{max}/\omega_{min}-1}}{\sqrt{\omega_{max}/\omega_{min}+1}}$ [? ]. |

## Algebraic Derivation

### 1. Relational Scale Factor

The ratio of the Sun's radius to the planet's perihelion distance is derived entirely from visual size and cycle ratios:

$$R_{ratio} = \frac{\theta_{\odot}}{(T_M/T_{\oplus})^{2/3} \cdot (1 - e_M)} \approx 0.0151290$$

### 2. Potential Projection (at perihelion)

Translating the spectral shift at the Sun's surface ( $z_{sun}$ ) to the potential projection (3.1) at Mercury's perihelion ( $\kappa_p^2$ ) using the derived scale factor:

$$\kappa_p^2 = (1 - \frac{1}{(1 + z_{sun})^2}) \cdot R_{ratio} \approx 6.4219 \times 10^{-8}$$

### 3. Kinematic Projection (at perihelion)

Deriving orbital kinematics strictly from the potential and eccentricity (3.3), requiring no Doppler or radar data:

$$\beta_p^2 = \frac{\kappa_p^2}{2}(1 + e_M) \approx 3.8712 \times 10^{-8}$$

### 4. Global Orbital Invariants

Extracting the system's global kinetic projection ( $\beta$ ) via the invariant binding energy  $\beta^2 = \kappa_p^2 - \beta_p^2$  (via [Energy-Symmetry Law sec:energy-symmetry](#)):

$$\beta^2 \approx 2.5508 \times 10^{-8} \quad \text{and} \quad \kappa^2 = 2\beta^2 \approx 5.1016 \times 10^{-8}$$

(via [Closure theorem thm:closure](#)).

### 5. Exact Orbital Precession

The secular phase shift emerges (3.7) from the accumulation of the relational state divergence ( $\tau_Y^2 = 1 - (1 - \kappa^2)(1 - \beta^2)$ ) over a full cycle:

$$\Delta\varphi = \frac{2\pi \cdot \tau_Y^2}{1 - e_M^2} \approx 5.0203 \times 10^{-7} \text{ rad/orbit}$$

Converting this pure geometric phase shift to standard observational units yields **43.015 arcsec/century**, perfectly matching the observed precession of Mercury.

## 6. Absolute System Scale (Metric Translation)

To translate the geometric scale into legacy metric units (meters), we map Mercury's orbital period ( $T_M = 7,600,521.6$  s) and the speed of light ( $c$ ) to the global kinematic projection ( $\beta$ ):

$$R_{sRG} = T_M \cdot c \cdot \frac{\beta^3}{\pi} \approx 2954.8 \text{ m}$$

Standard GR value:

$$R_{sGR} = \frac{2GM_{sun}}{c^2} \approx 2953.33 \text{ m} \quad \text{where} \quad M_{sun} = 1.98847 \times 10^{30} \text{ kg}$$

Discrepancy:

$$\frac{R_{sGR} - R_{sRG}}{R_{sGR}} \times 100 = |0.0476|\%$$

This variance is strictly within the observational uncertainty limits of the input parameters ( $z_{sun}$  and  $\theta_\odot$ ).

**7. Semi-major Axis** (Standard value  $\approx 5.79 \times 10^{10}$  m):

$$a = \frac{R_{sRG}}{\kappa^2} \approx 5.792288 \times 10^{10} \text{ m}$$

**8. Perihelion Radius** (Standard value  $\approx 4.60 \times 10^{10}$  m):

$$r_p = \frac{R_{sRG}}{\kappa_p^2} \approx 4.601394 \times 10^{10} \text{ m}$$

## Universality of the Framework: The Jovian Extension

To rigorously demonstrate that this parameterization is not a localized artifact of the Mercury-Sun proximity, but a universal geometric property of bound systems, we apply the exact same algebraic chain to Jupiter. We replace the inputs with Jupiter to Earth orbital periods ratios ( $T_{ratioJ}$ ) and Jupiter's eccentricity ( $e_J$ ), keeping the central reference invariants ( $\theta_\odot$ ,  $z_{sun}$ ,  $T_\oplus$ ) strictly identical.

### Jovian Inputs:

- $T_J = 4332.589$  days [? ]
- $e_J = 0.04839266$  [? ]
- $T_{ratioJ} = T_J/T_\oplus \approx 11.862$  [? ]

### 1. Global Potential Projection ( $\kappa_J$ )

The local potential projection at Jupiter's semi-major axis reduces to a remarkably elegant relation, scaling the solar surface tension strictly through the inverse temporal cycle ratio:

$$\kappa_J = \sqrt{\left(1 - \frac{1}{(1 + z_{sun})^2}\right) \cdot \frac{\theta_\odot}{(T_{ratioJ})^{2/3}}} \approx 6.1623 \times 10^{-5}$$

### 2. Absolute System Scale Recovery ( $R_{sRGJ}$ )

Expanding the algebraic chain utilizing Jupiter's specific kinematic projection ( $\beta_J$ ) yields the invariant core scale:

$$R_{sRGJ} = \frac{T_J \cdot c}{\pi} \left[ \frac{1}{1 - e_J} \left(1 - \frac{1}{(1 + z_{sun})^2}\right) \frac{\theta_\odot}{T_{ratioJ}^{2/3}} - \frac{1}{2} \frac{1 + e_J}{1 - e_J} \left(1 - \frac{1}{(1 + z_{sun})^2}\right) \frac{\theta_\odot}{T_{ratioJ}^{2/3}} \right]^{3/2}$$

$$R_{sRGJ} \approx 2955.41 \text{ m}$$

## Jovian Orbital Recovery

Translating the recovered systemic scale to the spatial scale of Jupiter's orbit:

$$a_J = \frac{R_{sRGJ}}{\kappa_J^2} \approx 7.7827 \times 10^{11} \text{ m}$$

Observed value  $a_{JDATA} = 778.57 \times 10^9 \text{ m}$ .

$$\text{Discrepancy} = \frac{|a_{JDATA} - a_J|}{a_{JDATA}} \times 100 \approx 0.038\%$$

The reconstruction of the Schwarzschild radius ( $R_s$ ) and the orbital scale ( $a$ ) from disparate chronometric cycles ( $T, e$ ) demonstrates that there's a constant direct invariant relation between:

- **potential** geometry ( $\kappa^2$  - "spacetime structure")

- and system's **kinetic** ( $\beta^2$ - "energy dynamics") both seen as relational measure of state difference, rooting directly from [foundational unification principle](#) **SPACETIME  $\equiv$  ENERGY** (pr:unifying).

□

Test it yourself

DESMOS: <https://www.desmos.com/calculator/mjen4ms452>

COLAB: [Mercury-Sun\\_Relational\\_Parameterization.ipynb](#)

## Conclusion

All major parameters of gravitational systems are recovered through algebraic closure without mass ( $M$ ), gravitational constant ( $G$ ), metric manifold, differential formalism, or external temporal flow.

**Corollary 4.6** (Epistemic Mandate and Parameter Parsimony). *If a system's generative chain is algebraically complete without a specific parameter, that parameter is an emergent scaling constant rather than a fundamental primitive. Because the structural and dynamical states  $(e, \Delta\varphi, R_s, a)$  are closed using only dimensionless spectroscopic and chronometric ratios  $(e, \theta_\odot, T_{ratio}, z_{sun})$ , the variables  $G$  and  $M$  function as translation factors for legacy unit systems. Their retention is required exclusively for mapping invariant relational geometry onto the kilogram-based convention without increasing the system's predictive capacity.*

COLAB: M and G independence full test

[ROM\\_MASS\\_G\\_FREE\\_CLOSURE.ipynb](#)

## 5 Time-Density Invariant ( $\tau_D$ )

**Definition 5.1** (Time-Density Invariant). *The time-density invariant  $\tau_D$  establishes the systemic phase-space limit for any stable trajectory, defined strictly through the orbital period  $T$  and visual angular radius  $\theta$ :*

$$\tau_D = T \sin(\theta)^{\frac{3}{2}} \quad (50)$$

**Theorem 5.2** (Relational Density). *The structural density of the central body is determined by the time-density invariant  $\tau_D$ , bypassing the Newtonian spatial mass distribution.*

*Proof.* For any stable trajectory, geometry enforces the trigonometric identity  $\sin(\theta) = \frac{R_{sur}}{a}$ . The operational parameter  $\tau_D$  algebraically defines the theoretical orbital cycle at the physical surface  $R_{sur}$  of the central body:

$$\tau_D = \frac{2\pi R_{sur}}{\beta_{sur} c} \quad (51)$$

where  $\beta_{sur}$  is the local kinematic projection at the surface. [Relational density  \$\rho\$  as derived in WILL RG Part I](#) : operates via spatial projection limits:

$$\rho = \frac{\kappa^2 c^2}{8\pi G a^2} \quad (52)$$

Isolating the invariant  $\tau_D$  structures this relationship as the strict inverse determinant of the system's central density:

$$\tau_D = \sqrt{\frac{\pi}{G\rho}} \quad (53)$$

□

**Theorem 5.3** (Universal Horizon Constant). *At the event horizon boundary limit, the proportional ratio of the limiting time-density  $\tau_{D(limit)}$  to the system scale  $R_s$  yields a universal horizon constant, operating exclusively on the universal tempo  $c$  and geometric constant  $\pi$ .*

*Proof.* The systemic horizon scale  $R_s$  isolates mathematically from the time-density invariant and surface kinematic projection  $\beta_S$ :

$$R_s = \frac{c}{\pi} \tau_D \beta_S^3 \quad (54)$$

At the event horizon boundary limit ( $R_{surf} = R_s$ ), the potential projection reaches maximum ( $\kappa^2 = 1$ ) and the kinematic projection fixes via the closure theorem (2) ( $\beta_S^2 = \frac{1}{2}$ ). The time-density limit evaluates to:

$$\tau_{D(limit)} = \frac{2\sqrt{2}\pi R_s}{c} \quad (55)$$



The proportional ratio resolves to the universal geometric constant:

$$\boxed{\frac{\tau_{D(limit)}}{R_s} = \frac{2\sqrt{2}\pi}{c}} \quad (56)$$

□

**Remark 5.4** (Ontological Consequence). *The algebraic isolation of  $\frac{\tau_{D(limit)}}{R_s}$  demonstrates that the boundary mechanics of a spacetime well operate exclusively on geometric proportionality and kinematic cycles. The parameters  $G$  and  $M$  are completely eradicated from the limiting topological structure of the system.*

## 6 Algebraic Determination of the Horizon Scale

A classical critique posits that because  $R_s$  can be equated to  $\frac{2GM}{c^2}$ , this derivation is a superficial algebraic reparameterization of Newtonian mechanics. As shown in 9, 8, [sec:earth-gps](#) and [ROM\\_FULL\\_TEST.ipynb](#), - this objection contradicts mathematical results and reverses the causal and ontological hierarchy.

The terms  $G$  and  $M$  are epistemologically secondary composites. They function as anthropic translation factors required to map Relational Geometry onto an assumed absolute spacetime container.  $R_s$ , conversely, is a primary, coordinate-independent relational invariant determent by the tension of the system's potential difference.

The derivation isolates true dynamic proportionality using exclusively operational observables (chronometric phase  $T$ ) and intrinsic horizon scaling ( $a, R_s$ ). Relational Geometry does not inherit from Newtonian constructs; rather, classical mechanics collapses from Relational Geometry when an absolute spatial background is artificially imposed.  $R_s$  does not conceal mass; rather, classical mass is an anthropocentric proxy for the invariant relational horizon  $R_s$ .

### 6.1 Method A: Differential (Two-Point Method)

This method is suitable when the orbital period is unknown, but the trajectory can be traced geometrically.

**Theorem 6.1** (Two-Point Schwarzschild Scale). *Given measurements of geometric position ( $r$ ) and kinematic intensity ( $\beta$ ) at two arbitrary points along a trajectory:*

- *Radii:  $r_1, r_2$  (from astrometry)*
- *Intensities:  $\beta_1, \beta_2$  (from de-projected spectral line widths or proper motion)*

*the Schwarzschild radius is:*

$$\boxed{R_s = \frac{r_1 r_2}{r_2 - r_1} (\beta_1^2 - \beta_2^2)}. \quad (57)$$

*Proof.* We invoke the **Conservation of the Global Kinetic Amplitude** (Vis-Viva Theorem), which states that for any closed system, the difference between the local potential and local kinetic projections is a global invariant equal to  $\beta^2$ .

Therefore, at any two points 1 and 2:

$$\kappa_1^2 - \beta_1^2 = \kappa_2^2 - \beta_2^2. \quad (58)$$

Rearranging to group the projections:

$$\kappa_1^2 - \kappa_2^2 = \beta_1^2 - \beta_2^2. \quad (59)$$

Substituting the field identity  $\kappa^2 = R_s/r$ :

$$\frac{R_s}{r_1} - \frac{R_s}{r_2} = \beta_1^2 - \beta_2^2. \quad (60)$$

Factoring out the scale parameter  $R_s$ :

$$R_s \left( \frac{r_2 - r_1}{r_1 r_2} \right) = \beta_1^2 - \beta_2^2. \quad (61)$$

Solving for  $R_s$ :

$$R_s = \frac{\beta_1^2 - \beta_2^2}{\frac{r_2 - r_1}{r_1 r_2}} = \frac{r_1 r_2}{r_2 - r_1} (\beta_1^2 - \beta_2^2). \quad (62)$$

This formula extracts the system scale purely from geometric gradients. □



## 6.2 Method B: Geometric Resonance (Balance Point Method)

Using one of two orbital balance points:

- Ascending:  $B_a = \arccos(-e)$ ,
- Descending  $B_d = 2\pi - \arccos(-e)$ ,

when the system passes through its geometric balance points the instantaneous radius  $r$  equals the semi-major axis  $a$ . At this unique phases, the closure condition  $\kappa_o^2 = 2\beta_o^2$  is satisfied.

**Theorem 6.2** (Balance Point Formula). *Given the geometric scale  $a$  and the total light signal  $\tau$  measured at the balance point ( $r = a$ ):*

$$R_s = \frac{a}{2}(3 - \sqrt{1 + 8\tau^2(B_a)}). \quad (63)$$

$$B_a = \arccos(1 - \delta^{-1}) = \arccos(-e) = \arccos(1 - \frac{2\beta_p^2}{\kappa_p^2}) = \arccos(\frac{2\beta_a^2}{\kappa_a^2} - 1) \text{ (orbital balance point where } \kappa_o^2 = 2\beta_o^2 \text{ is true)}$$

*Proof. Step 1: Spacetime Factor Expansion.* At the balance point ( $r = a$ ), the closure condition implies  $\kappa^2 = R_s/a$  and  $\beta^2 = R_s/2a$ . The observable  $\tau$  is:

$$\tau^2 = (1 - \kappa^2)(1 - \beta^2) = (1 - \frac{R_s}{a})(1 - \frac{R_s}{2a}). \quad (64)$$

**Step 2: Exact Geometric Solution.** Expanding the product:

$$\tau^2 = 1 - \frac{R_s}{2a} - \frac{R_s}{a} + \frac{R_s^2}{2a^2} = 1 - \frac{3R_s}{2a} + \frac{R_s^2}{2a^2}. \quad (65)$$

Multiplying by  $2a^2$  to clear denominators and rearranging into standard quadratic form ( $Ax^2 + Bx + C = 0$ ):

$$R_s^2 - 3aR_s + 2a^2(1 - \tau^2) = 0. \quad (66)$$

Solving for  $R_s$  using the quadratic formula ( $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a_{coef}}$ ):

$$R_s = \frac{3a \pm \sqrt{(3a)^2 - 4(1)(2a^2(1 - \tau^2))}}{2}. \quad (67)$$

Simplifying the term under the radical:

$$9a^2 - 8a^2(1 - \tau^2) = 9a^2 - 8a^2 + 8a^2\tau^2 = a^2(1 + 8\tau^2). \quad (68)$$

Thus:

$$R_s = \frac{3a \pm a\sqrt{1 + 8\tau^2}}{2} = \frac{a}{2}(3 \pm \sqrt{1 + 8\tau^2}). \quad (69)$$

For stable orbits, we must select the negative root. □

**Remark 6.3** (Light Separation). *This unit-less factor in brackets is:*

- **Potential projection at semi-major axis:**  $\kappa^2 = \frac{1}{2}(3 - \sqrt{1 + 8\tau^2(B_a)})$

*A unique way to separate gravitational part from the light signal.*

## 6.3 Method C: Instantaneous (Arbitrary Phase Method)

Most generally, if the orbital geometry ( $a$ ) is known, the scale  $R_s$  can be derived from a **single epoch** observation at any arbitrary radius  $r_o$ .

**Theorem 6.4** (Arbitrary Phase Formula). *Given the orbital geometry ( $a, r_o$ ) and the single light observable  $\tau_o$ :*

$$R_s = \frac{r_o}{2(2a - r_o)}(4a - r_o - \sqrt{(4a - r_o)^2 - 8a(2a - r_o)(1 - \tau_o^2)}). \quad (70)$$

*Proof. Step 1: Relational Invariant Form.* We start with the global kinetic amplitude invariant  $\beta_{global}^2 = \kappa_o^2 - \beta_o^2$ . For a bound orbit, the global kinetic amplitude is  $\beta_{global}^2 = R_s/2a$ . We express the local kinematic intensity  $\beta_o^2$  at arbitrary radius  $r$  in terms of the local field intensity  $\kappa_o^2$ :

$$\kappa_o^2 - \beta_o^2 = \frac{R_s}{2a} \implies \beta_o^2 = \kappa_o^2 - \frac{R_s}{2a}. \quad (71)$$

Substituting  $\kappa_o^2 = R_s/r$ :

$$\beta_o^2 = R_s \left( \frac{1}{r} - \frac{1}{2a} \right). \quad (72)$$

**Step 2: Constraint via Observables.** We substitute the expressions for  $\kappa_o^2$  and  $\beta_o^2$  into the observable constraint  $\tau_o^2 = (1 - \kappa_o^2)(1 - \beta_o^2)$ :

$$\tau_o^2 = \left(1 - \frac{R_s}{r}\right) \left(1 - R_s \left[\frac{1}{r} - \frac{1}{2a}\right]\right). \quad (73)$$

Let  $A = \frac{1}{r}$  and  $B = \frac{1}{r} - \frac{1}{2a}$ . The equation becomes:

$$\tau_o^2 = (1 - AR_s)(1 - BR_s) = 1 - (A + B)R_s + AB R_s^2. \quad (74)$$

Rearranging into quadratic form:

$$(AB)R_s^2 - (A + B)R_s + (1 - \tau_o^2) = 0. \quad (75)$$

**Step 3: Coefficient Expansion.** We compute the coefficients explicitly:

$$AB = \frac{1}{r} \left( \frac{2a - r}{2ar} \right) = \frac{2a - r}{2ar^2} \quad (76)$$

$$A + B = \frac{1}{r} + \frac{2a - r}{2ar} = \frac{2a + 2a - r}{2ar} = \frac{4a - r}{2ar} \quad (77)$$

Multiplying the entire quadratic equation by  $2ar^2$  to clear denominators:

$$(2a - r)R_s^2 - r(4a - r)R_s + 2ar^2(1 - \tau_o^2) = 0. \quad (78)$$

**Step 4: Solution.** Solving for  $R_s$ :

$$R_s = \frac{r(4a - r) \pm \sqrt{r^2(4a - r)^2 - 4(2a - r)(2ar^2)(1 - \tau_o^2)}}{2(2a - r)}. \quad (79)$$

Factoring out  $r^2$  from the radical term  $\sqrt{r^2(\dots)} = r\sqrt{(\dots)}$ :

$$R_s = \frac{r}{2(2a - r)} ((4a - r) \pm \sqrt{(4a - r)^2 - 8a(2a - r)(1 - \tau_o^2)}). \quad (80)$$

For stable orbits, we select the negative root, yielding the algebraic link between the observed light phase and the system's geometric scale.  $\square$

## 7 Rotational Systems (Kerr Without Metric)

### Contextual Bounds

- **For a gravitationally closed (static) system**, the physical boundary is defined by the condition  $\kappa^2 = 1$ . The closure principle ( $\kappa^2 = 2\beta^2$ ) is what dictates that this corresponds to a kinetic state of  $\beta^2 = 1/2$ .
- **For a kinematically closed (maximally rotating) system**, the physical boundary is defined by the condition  $\beta^2 = 1$ . The same closure principle ( $\kappa^2 = 2\beta^2$ ) then necessitates that the corresponding gravitational state must be  $\kappa^2 = 2$ .

For rotating black holes, we establish the connection between relational kinetic projection and the Kerr metric by defining:

$$\beta = \frac{ac^2}{GM}, \quad \kappa = \sqrt{2}\beta$$

where:

- $\beta$  is the relational rotation parameter, with  $0 \leq \beta \leq 1$ ,
- $\kappa$  is related to the geometry and gravity,
- $R_s = \frac{2GM}{c^2}$  is the Schwarzschild radius,
- $a = \frac{J}{Mc}$  is the Kerr rotation parameter,
- $J$  is the angular momentum of the black hole,
- $M$  is the mass of the black hole.

We also derive a key invariant relationship:

$$a_{\max} = \frac{R_s}{2} = \beta_{\max}^2 r$$

This relationship holds when  $r = \frac{R_s}{2\beta^2}$ , providing an elegant connection between the parameters.

## Event Horizon

Using our approach, the inner and outer event horizons of the Kerr metric are expressed as:

$$r_{\pm} = \frac{R_s}{2}(1 \pm \beta_Y)$$

For the extreme case where  $\beta = 1$  (maximal rotation), the horizons merge at:

$$r_+ = r_- = \frac{R_s}{2}$$

This coincides with the minimum radius in our model predicted using maximum value of  $\kappa$  parameter  $\kappa_{max} = \sqrt{2}$ :

$$r_{\min} = \frac{1}{\kappa_{max}^2} R_s = \frac{1}{2} R_s$$

## Ergosphere

The radius of the ergosphere in our model is described as:

$$r_{\text{ergo}} = \frac{R_s}{2}(1 + \sqrt{1 - \beta^2 \cos^2 \theta})$$

This formulation correctly reproduces the key features of the ergosphere:

- At the equator ( $\theta = \pi/2$ ),  $r_{\text{ergo}} = R_s$  for any rotation parameter,
- At the poles ( $\theta = 0$ ),  $r_{\text{ergo}}$  coincides with the event horizon radius.

## Ring Singularity

Unlike the Schwarzschild metric with its point singularity, the Kerr metric features a ring singularity located at:

$$r = 0, \quad \theta = \frac{\pi}{2}$$

The "size" of this ring is proportional to  $a = \frac{GM}{c^2}\beta$ , reaching its maximum for extreme black holes ( $\beta = 1$ ).

## Naked Singularity

For  $\beta \leq 1$ , a naked singularity does not emerge, aligning with the cosmic censorship Principle. In our model, Energy Symmetry Law enforce constraint by limiting  $\beta$  to the range  $[0, 1]$ .

## 7.1 Chiral Asymmetry: The Algebraic Origin of Frame-Dragging

In standard General Relativity (GR), the chiral asymmetry of orbits around a rotating mass—commonly termed "frame-dragging" or the Lense-Thirring effect—is structurally derived from the off-diagonal  $g_{t\phi}$  component of the Kerr metric tensor. In Relational Orbital Mechanics (R.O.M.), this asymmetry is not a geometric deformation of a background manifold, but a strict algebraic consequence of linear phase superposition on the directed  $S^1$  carrier.

### Composite Kinematic Projection and DOF Conservation

Because the kinematic projection  $\beta$  operates on the 1-DOF relational carrier  $S^1$ , motions sharing the same directional axis must superimpose linearly prior to quadratic energy evaluation. For a test mass orbiting a spinning central body, the composite kinematic state is:

$$\beta_{\Sigma} = \beta_{\text{orb}} \pm \beta_{\text{spin}} \quad (81)$$

where (+) denotes prograde and (−) denotes retrograde motion.

To conserve dimensional degrees of freedom when projecting a 3D central body's spin onto the 1D carrier  $S^1$ , a 1/3 scaling factor is applied. The local operational spin projection scales the specific angular momentum  $a = J/(Mc)$  against the spatial depth  $r$ :

$$\beta_{\text{spin}} = \frac{a}{3r} \quad (82)$$

## Composite Kinematic State from the Relational Origin

By the [Relational Origin Principle](#), neither the orbiting test object nor the central body possesses an intrinsic velocity or an intrinsic spin. The only physically real quantity is the relational difference between the two bodies, registered as a single amplitude on the directed kinematic carrier  $S^1$ . There are therefore not two velocities to be vector-composed, but one  $S^1$  difference carrying two contributions of opposite chirality:

$$\beta_\Sigma = \beta_{\text{orb}} \pm \beta_{\text{spin}} \quad (83)$$

where (+) denotes prograde and (−) retrograde alignment of the spin contribution with the orbital difference. Because both are shares of the *same* difference, they superimpose linearly; squaring is applied only to the completed amplitude  $\beta_\Sigma$ , never to its individual shares.

## The Algebraic Symmetry Breaker

Applying the [relational closure theorem](#) ( $\kappa^2 = 2\beta^2$ ) to the composite state yields:

$$\kappa^2 = 2(\beta_{\text{orb}} \pm \beta_{\text{spin}})^2 = 2\beta_{\text{orb}}^2 + 2\beta_{\text{spin}}^2 \pm 4\beta_{\text{orb}}\beta_{\text{spin}} \quad (84)$$

The emergent cross-term  $\pm 4\beta_{\text{orb}}\beta_{\text{spin}}$  is the exact symmetry breaker. Because  $\kappa^2 = R_s/r$ , this mathematically mandates that a prograde orbit (+) requires a deeper potential projection (smaller radius) than a retrograde orbit (−) at an equivalent orbital velocity. This algebraic constraint is operationally indistinguishable from the spatial drag generated by the  $g_{t\phi}$  metric.

## Expansion of the Precessional Divergence

The internal causal order of the composite system evaluates via the relational phase factor  $\tau^2 = (1 - \kappa^2)(1 - \beta_\Sigma^2)$ . Expanding this algebraically and applying the strict directional boundary ( $\pm = 1$ ) isolates the chiral divergence:

$$\tau_{Y(\text{chiral})}^2 = \pm(6\beta_{\text{orb}}\beta_{\text{spin}} - 8\beta_{\text{orb}}^3\beta_{\text{spin}} - 8\beta_{\text{orb}}\beta_{\text{spin}}^3) \quad (85)$$

Substituting the local operational projection into the isolated chiral polynomial defines the phase shift per orbital cycle, normalized by the orbital shape ( $1 - e^2$ ):

$$\Delta\varphi_{\text{chiral}} = \frac{2\pi}{1 - e^2} [\pm(6\beta_{\text{orb}}\beta_{\text{spin}} - 8\beta_{\text{orb}}^3\beta_{\text{spin}} - 8\beta_{\text{orb}}\beta_{\text{spin}}^3)] \quad (86)$$

Colab Sympy Verification

[Kerr sympy.ipynb](#)

## 7.2 Legacy Translation: Lense-Thirring Equivalence

To verify structural compatibility with historical models, we isolate the primary geometric observable by ablating the higher-order geometric cross-coupling terms ( $\beta^4 \rightarrow 0$ ) and setting the orbit to circular ( $e = 0$ ). The equation simplifies algebraically to:

$$\Delta\varphi_{\text{ablated}} \simeq 12\pi\beta_{\text{orb}}\beta_{\text{spin}} \quad (87)$$

Substituting the operational spin projection  $\beta_{\text{spin}} = \frac{J}{3Mc r}$  and the kinematic orbital projection  $\beta_{\text{orb}} = \frac{v}{c}$  recovers the empirical post-Newtonian node precession:

$$\Delta\varphi_{\text{ablated}} \simeq 12\pi\left(\frac{v}{c}\right)\left(\frac{J}{3Mc r}\right) = \frac{4\pi v J}{Mc^2 r} \quad (88)$$

Applying the kinematic scaling identity  $\frac{v}{M} = \frac{G}{r v}$ :

$$\Delta\varphi_{\text{ablated}} \simeq \frac{4\pi G J}{c^2 r^2 v} \quad (89)$$

The Lense-Thirring effect emerges natively from the algebraic closure of the relational carriers, executed without invoking differential tensors or a curved pseudo-Riemannian manifold.

COLAB Empirical Verification

This Chiral Asymmetry when tested against LAGEOS-1 Satellite (Earth orbit) shows perfect agreement with observations and GR predictions: [LAGEOS I Lense-Thirring.ipynb](#)

### 7.3 Epistemological Conclusion

To generate chiral orbital asymmetry (frame-dragging), exact General Relativity must postulate a pseudo-Riemannian manifold with an off-diagonal metric tensor component,  $g_{t\phi}$ , to induce spatial torsion. R.O.M. reproduces the identical physical symmetry and precessional kinematics algebraically. Enforcing the [relational closure theorem](#) onto the composite one-dimensional carrier  $\beta_\Sigma = \beta_{\text{orb}} \pm \beta_{\text{spin}}$  yields the necessary quadratic expansion. The emergent cross-term:

$$\pm 4\beta_{\text{orb}}\beta_{\text{spin}} \quad (90)$$

mathematically mandates the chiral divergence. This algebraic constraint is operationally indistinguishable from the spatial drag generated by the  $g_{t\phi}$  metric. The geometric deformation of a background spacetime is therefore mathematically obsolete. The dimensional container and its differential tensors are eradicated from the ontology.

## 8 Mercury's Precession: GR 1PN as First-Order Approximation of RG

### 8.1 Statement of the Verification Problem

The standard first-order post-Newtonian expression for the perihelion advance of a bound two-body orbit is

$$\Delta\varphi_{\text{GR}} = \frac{3\pi R_s}{a(1-e^2)}, \quad (91)$$

with  $a$  the semi-major axis,  $e$  the eccentricity, and  $R_s = 2GM/c^2$  the Schwarzschild scale of the central body.

WILL Relational Geometry (WILL RG) expresses the secular angular shift per closed orbit as the strict geometric accumulation of the system's phase divergence from the rest origin:

$$\Delta\varphi_{\text{RG}} = \frac{2\pi\tau_Y^2}{1-e^2}, \quad \tau_Y^2 \equiv 1 - \tau^2 = 1 - (1 - \beta^2)(1 - \kappa^2) = \beta^2 + \kappa^2 - \beta^2\kappa^2, \quad (92)$$

evaluated at the orbital reference scale (semi-major axis  $a$ ), where the closure condition  $\beta^2(a) = \kappa^2(a)/2$  and the field relation  $\kappa^2(a) = R_s/a$  hold.

The question addressed in this section is whether (91) is the leading-order content of (92) in the small parameter  $R_s/a$ . The structure of (92) suggests an even stronger statement than in the time-dilation case ([sec:earth-gps](#)): the GR formula should be recoverable by removing a single explicit term — the cross-coupling  $\beta^2\kappa^2$  — from  $\tau_Y^2$ , with the residual being exactly  $-2\pi\beta^2\kappa^2/(1-e^2)$ .

### 8.2 Methodology

The verification is split into two independent parts, both executed in a single public notebook (Section 8.5).

**Part I — symbolic identity.** The exact WILL RG precession is expressed in closed form by substituting the [closure condition](#)  $\beta^2 = \kappa^2/2$  and  $\kappa^2 = R_s/a$  into (92). The result is simplified algebraically and subtracted from (91). The algebraic identity between the residual and the analytically predicted second-order term  $-2\pi\beta^2\kappa^2/(1-e^2)|_a$  is verified using `sympy.simplify`. As an independent cross-check, (92) is Taylor-expanded in a bookkeeping parameter  $\varepsilon$  that scales  $R_s$ , and the first-order coefficient is compared with (91).

**Part II — numerical comparison.** Both (91) and (92) are evaluated with 40-digit precision arithmetic (`mpmath`, `mp.dps = 40`) using Mercury's J2000 orbital elements:  $a = 5.7909050 \times 10^{10}$  m,  $e = 0.20563593$ , orbital period  $T = 87.9691$  d,  $GM_\odot = 1.32712440018 \times 10^{20}$  m<sup>3</sup>s<sup>-2</sup>,  $c = 299,792,458$  m s<sup>-1</sup>, giving  $R_s^\odot = 2953.25$  m. Per-orbit values are converted to arcseconds per Julian century using  $36,525$  d/ $T$  orbits per century. The numerical discrepancy  $\Delta\varphi_{\text{RG}} - \Delta\varphi_{\text{GR}}$  is compared with the analytically predicted second-order correction obtained in Part I.

### 8.3 Results

**Symbolic.** Substituting the closure and field relations into (92) yields the closed-form WILL RG precession

$$\Delta\varphi_{\text{RG}} = \frac{3\pi R_s}{a(1-e^2)} - \frac{\pi R_s^2}{a^2(1-e^2)}. \quad (93)$$

The first term is identically (91). The algebraic difference is therefore

$$\Delta\varphi_{\text{RG}} - \Delta\varphi_{\text{GR}} = -\frac{\pi R_s^2}{a^2(1-e^2)} \equiv -\frac{2\pi\beta^2\kappa^2}{1-e^2}\bigg|_a. \quad (94)$$

The identity  $[\Delta\varphi_{\text{RG}} - \Delta\varphi_{\text{GR}}] - [-2\pi\beta^2\kappa^2/(1-e^2)]|_a = 0$  vanishes identically in  $(R_s, a, e)$ . The Taylor expansion of (92) in  $R_s/a$  reproduces (91) at first order with zero residual.

**Structural observation.** In the time-dilation case the GR additive formula was the linear coefficient of a Taylor expansion of a multiplicative  $\tau$ -ratio. In the present case the correspondence is more direct: the GR first-order PN precession formula coincides exactly with the WILL RG expression *after a single explicit cancellation*, namely  $\beta^2\kappa^2 \rightarrow 0$  in  $\tau_Y^2 = 1 - (1 - \beta^2)(1 - \kappa^2)$ . The omitted term encodes the second-order coupling between the kinematic  $S^1$  and gravitational  $S^2$  projection carriers of the WILL framework.

**Numerical.** With Mercury's parameters listed above:

| quantity                                                              | value                              |
|-----------------------------------------------------------------------|------------------------------------|
| $\Delta\varphi_{\text{RG}}$ (exact, per orbit)                        | $5.01867565337 \times 10^{-7}$ rad |
| $\Delta\varphi_{\text{GR}}$ (first-order PN, per orbit)               | $5.01867573869 \times 10^{-7}$ rad |
| $\Delta\varphi_{\text{RG}} - \Delta\varphi_{\text{GR}}$ (per orbit)   | $-8.531 \times 10^{-15}$ rad       |
| $-\pi R_s^2/[a^2(1 - e^2)]$ (predicted, per orbit)                    | $-8.531 \times 10^{-15}$ rad       |
| residual                                                              | $\sim 10^{-41}$ rad                |
| $\Delta\varphi_{\text{RG}}$ (per century)                             | 42.9807844914''                    |
| $\Delta\varphi_{\text{GR}}$ (per century)                             | 42.9807852220''                    |
| $\Delta\varphi_{\text{RG}} - \Delta\varphi_{\text{GR}}$ (per century) | $-7.306 \times 10^{-7}$ ''         |

The numerical discrepancy  $\Delta\varphi_{\text{RG}} - \Delta\varphi_{\text{GR}}$  matches the analytically derived second-order term (94) to within the limit set by the 40-digit floating-point arithmetic. The relative magnitude of the term omitted by the first-order GR formula is  $\sim 1.7 \times 10^{-8}$  of the total precession.

## 8.4 Interpretation

Equation (93) establishes that the standard first-order post-Newtonian formula (91) is the leading-order content of the WILL RG exact expression (92) in  $R_s/a$ , and that the omitted contribution is exactly the cross-coupling term  $-2\pi\beta^2\kappa^2/(1 - e^2)$  at the semi-major axis. This is an algebraic identity in  $(R_s, a, e)$ , not a numerical near-match.

For Mercury, the second-order term in (93) contributes at the level of  $-7.3 \times 10^{-7}$ '' century $^{-1}$ , i.e.  $\sim 1.7 \times 10^{-8}$  of the total precession of  $\approx 42.98$ '' century $^{-1}$ . This deviation is well below the current precision of perihelion measurements and does not, by itself, constitute a distinguishing observation between the two formulations in the Solar System regime. The sign of the correction is fixed: WILL RG predicts a slightly smaller precession than the first-order post-Newtonian formula.

The scope of the present statement is restricted to the relation between the WILL RG closed-form expression (92) and the *first-order post-Newtonian* GR precession formula (91). A comparison against fully non-linear geodesic integration in the exact Schwarzschild geometry — which itself receives positive-definite corrections beyond the leading  $3\pi R_s/[a(1 - e^2)]$  term — is a separate calculation and is not addressed in this section.

## 8.5 Reproducibility

The complete symbolic and numerical verification reported in this section is contained in a single public Google Colab notebook, comprising the two scripts used to produce (93), (94), and the numerical table above:

Colab Notebook

[Mercury precession RG 1st order=GR.ipynb](#)

The notebook has no external dependencies beyond `sympy` and `mpmath` and is self-contained. All input parameters are listed explicitly in the numerical script; any change in  $a$ ,  $e$ , or  $GM_\odot$  rescales both  $\Delta\varphi_{\text{RG}}$  and  $\Delta\varphi_{\text{GR}}$  consistently and does not affect the symbolic identity (94).

## 9 S2 Star (Sgr A\*): R.O.M. vs GR and Instrumental Systematics

We apply Relational Orbital Mechanics to the 24-year astrometric and radial-velocity dataset of the S2 star orbiting the Galactic Centre: 46 astrometric epochs (1995–2018) and 82 radial velocities from six instruments, 174 raw observables in total. The R.O.M. model is confronted with two General Relativity references. The first is the standard first post-Newtonian parametrisation with a time-linear pericentre advance  $\omega(t) = \omega_0 + \dot{\omega}(t - T_0)$ . The second is a direct numerical integration of the 1PN equations of motion in harmonic gauge,

$$\ddot{\vec{r}} = -\frac{GM}{r^2} \left[ \left( 1 + \frac{v^2}{c^2} - \frac{4GM}{c^2 r} \right) \hat{n} - \frac{4\dot{r}}{c^2} \vec{v} \right], \quad (95)$$

which contains no parametrisation choice of any kind: the trajectory is whatever GR produces. The integration is validated independently of the fits: trajectories agree to  $5 \times 10^{-4} \mu\text{as}$  between tolerance settings and to  $1.2 \times 10^{-2} \mu\text{as}$  between integration methods, the 1PN energy returns to its value at successive pericentre passages to  $1.6 \times 10^{-12}$  relative, and the measured apsidal advance reproduces  $6\pi GM/c^2 p$  to  $1.2 \times 10^{-4}$  (12.40' per orbit). All three models carry the same

nine free parameters ( $a_{\text{as}}, e, i, \omega, \Omega, P, T_0, V_0, K$ ), zero free precession parameters, identical single-step Rømer correction, identical data and error model.

In R.O.M. the pericentre advance and the system scale are algebraically locked to the closure invariant:

$$f_{\text{prec}} = \frac{3\beta^2 - 2\beta^4}{1 - e^2}, \quad \beta = \frac{K\sqrt{1 - e^2}}{c \sin i}, \quad Z_{\text{sys}} = \frac{1}{\tau_o} = \frac{1}{\sqrt{1 - \kappa_o^2} \sqrt{1 - \beta_o^2}}, \quad R_s = \frac{Tc\beta^3}{\pi}, \quad a = \frac{R_s}{2\beta^2}. \quad (96)$$

The GR 1PN advance, expressed through the same fitted observables, is  $\Delta\omega_{\text{1PN}}/2\pi = 3(K/c)^2/\sin^2 i$  per orbit; at S2's velocity amplitude the two magnitudes agree to  $3 \times 10^{-5}$  relative, so any difference between the fits originates in trajectory geometry, not in the size of the advance.

## Bare nine-parameter comparison and its resolution

| Model (9 orbital parameters each)                      | $\chi^2$ bare | $\chi^2$ + frame & RV offsets | $\chi^2/\text{dof}$     |
|--------------------------------------------------------|---------------|-------------------------------|-------------------------|
| WILL R.O.M. (closure-locked, phase-coupled)            | 728.6         | 228.3                         | 4.42 $\rightarrow$ 1.46 |
| GR 1PN, time-linear $\omega(t)$                        | 800.8         | 225.1                         | 4.85 $\rightarrow$ 1.44 |
| GR 1PN, direct integration of motion in harmonic gauge | 809.3         | 227.8                         | 4.90 $\rightarrow$ 1.46 |

Table 3: S2 fit quality. Bare fits use orbital parameters only; the right column adds the standard calibration model of published S2 analyses (astrometric zero point  $x_0, y_0$ , reference-frame drift  $v_{x0}, v_{y0}$ , five inter-instrument radial-velocity offsets), applied identically to every model and profiled exactly. dof = 165 and 156 respectively.

Taken at face value, the bare comparison favours R.O.M. by  $\Delta\chi^2 = 80.7$  against integrated GR with equal parameter count. This number is arithmetically genuine and is not a property of the GR implementation: the time-linear baseline reproduces the integrated trajectory to  $36 \mu\text{as}$  RMS on noise-free synthetic data, well inside the  $574 \mu\text{as}$  median astrometric error, so replacing it by the exact integration changes the gap by only 8.5 units. The number is nevertheless not a statement about gravitation. A reduced  $\chi^2$  of 4.4–4.9 rejects all three bare models at high significance: no published analysis of this dataset fits absolute sky positions without a reference-frame solution, because the infrared frame is known to carry zero-point and drift uncertainty at the mas and  $0.1 \text{ mas yr}^{-1}$  level. When the standard linear calibration terms are granted identically to every model, all three converge to the same frame solution (offset  $\approx 1 \text{ mas}$ , drift  $0.08\text{--}0.21 \text{ mas yr}^{-1}$ , instrument offsets  $10\text{--}95 \text{ km s}^{-1}$ ) and the gap closes completely:  $\Delta\chi^2 = -0.5$  over 156 degrees of freedom. Relational Geometry and General Relativity are statistically indistinguishable on the S2 orbit.

## Injection–recovery: mechanism, directionality, non-circularity

The origin of the bare-fit difference was established by controlled injection. Synthetic observables were generated at the real epochs with the real error model, from a known truth, and refitted by both nine-parameter models.

| Injected truth                                 | Fit configuration      | $\Delta\chi^2$ (GR – R.O.M.) |
|------------------------------------------------|------------------------|------------------------------|
| Integrated GR, no systematics                  | bare, 9 par.           | –12                          |
| Integrated GR + frame drift, data direction    | bare, 9 par.           | +50                          |
| Integrated GR + frame drift, rotated direction | bare, 9 par.           | –24                          |
| R.O.M. + frame drift                           | with calibration model | $+18.7 \pm 11.2$             |
| Integrated GR + frame drift                    | with calibration model | $-5.8 \pm 11.4$              |
| <b>Real S2 data</b>                            | with calibration model | <b>–0.5</b>                  |

Table 4: Injection–recovery results (six noise realisations per calibrated row; injected drift in the data direction:  $x_0 = -1.0$ ,  $y_0 = +0.9 \text{ mas}$ ,  $v_{x0} = +0.08$ ,  $v_{y0} = +0.20 \text{ mas yr}^{-1}$ ). The real data fall on the GR-truth expectation and  $1.7\sigma$  below the R.O.M.-truth expectation.

Three facts follow. First, R.O.M. holds no generic advantage: on clean GR truth it loses, and it loses again when the injected drift points in a rotated direction. Second, the full observed advantage is manufactured by injecting a purely instrumental drift into purely GR data; the bare  $\Delta\chi^2$  therefore measures alignment between R.O.M.'s trajectory family and this dataset's particular frame drift, not relational dynamics. Third, the calibration model is not a signal eraser and is not circular: when R.O.M. is the injected truth its advantage survives the identical procedure at  $+18.7 \pm 11.2$ , the frame parameters are profiled freely inside each model's own fit with no GR quantity entering the R.O.M. fit anywhere, and R.O.M.'s own fit independently demands the same frame solution, improving by 500  $\chi^2$  units when granted it.

## Rigidity, not flexibility

The absorption is a structural property of the closure, not added freedom. Fitted to noise-free integrated-GR observables, the R.O.M. family cannot approach the GR trajectory below a  $440 \mu\text{as}$  RMS floor, so it is not a superset of GR and possesses no surplus fitting capacity; it carries zero free precession parameters, and the closure  $\kappa^2 = 2\beta^2$  removes degrees



of freedom rather than adding them. Relative to the time-linear family it owns exactly one rigid distortion mode: the apsidal advance enters as  $\omega_{\text{shift}} = f_{\text{prec}} O$ , locked to the true phase. The osculating apsis of the integrated GR orbit itself advances in the same phase-locked manner, accumulating 67% of each orbit's advance within six months of pericentre against 7% for a linear ramp, so phase-locked advance is native to GR's osculating dynamics in the Damour–Deruelle quasi-Keplerian sense; the single-eccentricity rosette is a third curve, distinct from both the time-linear approximation and the exact trajectory. That single rigid mode, projected through 24 years of epochs, happens to point along the documented frame drift of this dataset. A one-mode family absorbing a one-direction systematic by alignment is the precise mechanism by which a less flexible model outperformed a more accurate one, and it is the reason an explicit one-line drift term cancels the effect exactly.

## What the S2 test establishes

At the precision of current S2 data, R.O.M. is observationally equivalent to General Relativity. It reaches this equivalence from a closed dimensionless algebra: the advance magnitude matches GR 1PN to  $3 \times 10^{-5}$ , the full physical scale follows from chronometric and spectroscopic ratios alone through  $R_s = Tc\beta^3/\pi$ , with  $G$  and  $M$  entering only as a posteriori unit conversions, and the complete fit is evaluated in closed form. The genuine point of departure between the frameworks is of order  $\beta^4$ , where  $f_{\text{prec}} = (3\beta^2 - 2\beta^4)/(1 - e^2)$  separates from the GR second post-Newtonian structure; at S2's  $\beta^2 \approx 4 \times 10^{-5}$  this lies four orders of magnitude below the sensitivity of the dataset. The falsifiable frontier of the closure law is therefore not stellar orbits at the Galactic Centre but regimes where  $\beta^4$  terms are resolved: relativistic pulsar binaries, horizon-scale orbits, and gravitational-wave inspirals.

### Reproduction and MCMC Analysis

All fits, the 1PN integration, the calibration profiling, and the injection–recovery experiments are reproduced by the scripts in [ROMvsGR S2 study.ipynb](https://github.com/ROMvsGR/S2_study.ipynb). Full MCMC Analysis [https://willrg.com/MCMC\\_S2.html](https://willrg.com/MCMC_S2.html)

## 10 Relational Derivation of the L1 Equilibrium Point

The calculation of the Sun–Earth L1 equilibrium point is performed strictly through relational parameters derived from direct observables, independent of mass  $M$  or the gravitational constant  $G$ .

### 10.1 Observational Primitives

The derivation utilises only the following epistemologically direct quantities (identical in form to those used for Mercury and S2 in earlier sections):

- $\theta_{\odot}$ : angular radius of the Sun ( $0.0046524$  rad).
- $z_{\odot}$ : spectroscopic fractional frequency shift at the solar surface ( $2.1224 \times 10^{-6}$ ).
- $T_E$ : sidereal orbital period of the Earth ( $3.15576 \times 10^7$  s).
- $T_m$ : sidereal orbital period of the Moon ( $2.3606 \times 10^6$  s).
- $R_m$ : mean Earth–Moon distance ( $3.844 \times 10^8$  m).

### 10.2 Structural Parameters from Relational Projections

Using the absolute-scale formulae established in Section 6 (Method A) and the global kinematic relation  $R_s = Tc\beta^3/\pi$ :

$$R_{s\odot} = \frac{T_E}{\pi} c \sqrt{\frac{1}{2} \left(1 - \frac{1}{(1+z_{\odot})^2}\right) \sin(\theta_{\odot})}^3, \quad (97)$$

$$\beta_m = \frac{2\pi R_m}{T_m c}, \quad (98)$$

$$R_{sE} = \frac{T_m}{\pi} c \beta_m^3, \quad (99)$$

$$R_E = \frac{R_{s\odot}}{\left(1 - \frac{1}{(1+z_{\odot})^2}\right) \sin(\theta_{\odot})}. \quad (100)$$

These expressions recover the Earth–Sun distance  $R_E \approx 1.496 \times 10^{11}$  m and the Earth/Sun scale ratio  $\mu = R_{sE}/R_{s\odot} \approx 3.003 \times 10^{-6}$  directly from the observables, without any reference to Newtonian mass.

### 10.3 Earth–L1 Distance Derivation

Let  $R_{L1}$  be the distance from Earth to the L1 point and define the normalised separation

$$\alpha = \frac{R_E - R_{L1}}{R_E}.$$

Synchronous motion at L1 requires the kinematic projection  $\beta_{L1}$  to satisfy the orbital-period constraint. Enforcing the closure theorem  $\kappa^2 = 2\beta^2$  together with energy symmetry yields the exact quintic

$$\alpha^5 - 2\alpha^4 + \alpha^3 + \left(\frac{R_{sE}}{R_{s\odot}} - 1\right)\alpha^2 + 2\alpha - 1 = 0.$$

In the limit  $R_{sE} \ll R_{s\odot}$  the physical root is

$$R_{L1} \approx R_E \sqrt[3]{\frac{R_{sE}}{3R_{s\odot}}}.$$

Substituting the numerical values obtained above gives

$$R_{L1} \approx 1.498 \times 10^9 \text{ m}$$

(with more precise ephemeris constants the value rises to  $1.501 \times 10^9 \text{ m}$ ). Both figures lie within the combined uncertainty of the input observables  $z_{\odot}$  and  $\theta_{\odot}$ .

#### Result

The relational derivation reproduces the observed Earth–L1 distance

$$R_{L1} \approx 1.5 \times 10^9 \text{ m}$$

to better than 0.3% without invoking  $G$ ,  $M$ , or a background metric. The same algebraic chain that recovers Mercury’s precession and the S2 orbit now also locates the collinear equilibrium point of the Sun–Earth–test-mass system.

#### Test it yourself

Full symbolic + numerical verification (including the exact quintic root): [L1 Relational Derivation.ipynb](#)

This section closes the demonstration that all major gravitational observable - two-body orbits, precession, light deflection, and now the three-body L1 point - emerges as an inevitable algebraic consequence of the Foundational Methodological Principles.

Proceed to next section of WILL Relational Geometry paper: [Density, Mass, and Pressure](#)