

# A Recombining Trinomial Option Pricing Model with Time-Dependent Volatility

## 1 Introduction

This work develops a recombining trinomial tree framework for option pricing under time-dependent (piecewise-constant) volatility. The objective is to construct a numerically efficient and stable pricing model that supports both European and American options and can be calibrated to observed market prices. The resulting framework provides additional flexibility over constant-volatility models while retaining analytical transparency and fast computation.

## 2 Model Setup

### 2.1 Time Discretisation

Let  $S_0$  denote the initial underlying price,  $r$  the continuously compounded risk-free rate, and  $T$  the maturity. For the sake of notational simplicity, we assume a zero dividend yield (if this was not the case, we could just replace the risk-free rate  $r$  with  $b = r - q$ ). The time interval  $[0, T]$  is divided into  $N$  equal steps of length:

$$\Delta t = \frac{T}{N}, \quad t_i = i\Delta t.$$

### 2.2 Volatility Structure

The instantaneous volatility is modelled as a deterministic function of time  $\sigma(t)$ . For numerical implementation, it is discretised as

$$\sigma_i = \sigma(t_i), \quad i = 0, 1, \dots, N-1.$$

In practice, the volatility term structure is assumed to be piecewise constant:

$$\sigma(t) = \sigma_m, \quad t \in [\tau_{m-1}, \tau_m), \quad m = 1, \dots, M.$$

The model parameters are therefore collected in the vector

$$\Theta = (\sigma_1, \sigma_2, \dots, \sigma_M).$$

This specification corresponds to a local volatility model in time, rather than a full spot-dependent local volatility surface.

### 2.3 Trinomial Price Dynamics

At each time step, the underlying price evolves according to a trinomial rule:

$$S \longrightarrow \begin{cases} Su & (\text{up}), \\ S & (\text{middle}), \\ Sd & (\text{down}). \end{cases}$$

**Grid Construction.** To ensure the tree remains recombining (i.e.,  $S_{i,j} = S_0 u^j$ ) under time-dependent volatility, we construct a fixed spatial grid based on a reference volatility  $\sigma_{\text{ref}}$  (this will be the average of  $\Theta$ ). The up and down factors are defined as constant across all time steps:

$$u = \exp\left(\sigma_{\text{ref}}\sqrt{2\Delta t}\right), \quad d = \frac{1}{u}.$$

A generic node price is given by

$$S_{i,j} = S_0 u^j, \quad j \in \{-i, -i+1, \dots, i-1, i\}.$$

The time-dependent nature of the volatility  $\sigma_i$  is incorporated not by changing the grid spacing  $u$ , but by adjusting the transition probabilities  $\{p_{u,i}, p_{m,i}, p_{d,i}\}$  at each step  $i$ . This approach preserves the computational efficiency of the recombining lattice while correctly capturing the local variance  $\sigma_i^2 \Delta t$ .

## 3 Risk-Neutral Probability Construction

Define the one-step multiplicative return

$$X_{i+1} = \begin{cases} u & \text{with probability } p_{u,i}, \\ 1 & \text{with probability } p_{m,i}, \\ d & \text{with probability } p_{d,i}. \end{cases}$$

Then

$$S_{i+1} = S_i X_{i+1}.$$

The probabilities are determined by matching the first two moments under the risk-neutral measure.

### Normalisation

$$p_{u,i} + p_{m,i} + p_{d,i} = 1.$$

### Risk-Neutral Mean

No-arbitrage requires

$$\mathbb{E}[S_{i+1} \mid S_i] = S_i e^{r\Delta t},$$

which implies

$$up_{u,i} + p_{m,i} + dp_{d,i} = M, \quad M := e^{r\Delta t}.$$

## Local Variance

The local variance condition is imposed as

$$\text{Var}[X_{i+1}] = \sigma_i^2 \Delta t.$$

Let

$$\mathbb{E}[X_{i+1}^2] = u^2 p_{u,i} + p_{m,i} + d^2 p_{d,i}.$$

Then

$$u^2 p_{u,i} + p_{m,i} + d^2 p_{d,i} = \sigma_i^2 \Delta t + M^2 =: V.$$

## 4 Explicit Probability Formulas

The previous conditions lead to the linear system

$$\begin{pmatrix} 1 & 1 & 1 \\ u & 1 & d \\ u^2 & 1 & d^2 \end{pmatrix} \begin{pmatrix} p_{u,i} \\ p_{m,i} \\ p_{d,i} \end{pmatrix} = \begin{pmatrix} 1 \\ M \\ V \end{pmatrix}.$$

Since the coefficient matrix depends only on  $(u, d)$ , the system admits a closed-form solution. The exact analytical probabilities are

$$p_{u,i} = \frac{V - M(d+1) + d}{(u-1)(u-d)},$$

$$p_{d,i} = \frac{V - M(u+1) + u}{(d-1)(d-u)},$$

$$p_{m,i} = 1 - p_{u,i} - p_{d,i}.$$

These expressions are used directly in the implementation for improved computational efficiency. For reasonable choices of  $\Delta t$  and  $\sigma_i$ , the resulting probabilities remain within the admissible range.

## 5 Option Valuation

### Terminal Payoff

At maturity  $t_N$ ,

$$V_{N,j} = \begin{cases} \max(S_{N,j} - K, 0) & \text{Call,} \\ \max(K - S_{N,j}, 0) & \text{Put.} \end{cases}$$

## Backward Induction

For  $i = N - 1, \dots, 0$ ,

$$V_{i,j}^{\text{cont}} = e^{-r\Delta t} (p_{u,i} V_{i+1,j+1} + p_{m,i} V_{i+1,j} + p_{d,i} V_{i+1,j-1}).$$

For European options,

$$V_{i,j} = V_{i,j}^{\text{cont}}.$$

For American options,

$$V_{i,j} = \max(V_{i,j}^{\text{cont}}, V_{i,j}^{\text{intr}}).$$

The option price at inception is  $V_{0,0}$ .

## 6 Volatility Calibration

Given market option prices  $V_{\text{market}}(K_k, T_k)$ , the calibration problem is formulated as

$$E(\Theta) = \sum_k w_k (V_{\text{model}}(K_k, T_k; \Theta) - V_{\text{market}}(K_k, T_k))^2 + \lambda \sum_{m=1}^{M-1} (\sigma_{m+1} - \sigma_m)^2.$$

with calibration weights satisfying

$$\sum_k w_k = 1.$$

The calibrated volatility term structure is obtained as

$$\Theta^* = \arg \min_{\Theta} E(\Theta),$$

using a derivative-free optimisation algorithm.

## 7 Conclusion

The proposed model combines a recombining trinomial lattice with time-dependent volatility, analytical probability construction, and a flexible calibration framework. This results in an efficient and robust approach for pricing and calibrating European and American options while maintaining convergence to the Black–Scholes–Merton model in the continuous-time limit.