

Regularized Equality-Constrained Quadratic Programming

Anton Xue and Nikolai Matni

1 Introduction

We look at regularized approximations of equality-constrained quadratic programming. In particular, how much does the optimal solution deviate when a regularizer is introduced?

2 Background

Stephen Boyd and Lieven Vandenberghe [1].

2.1 Equality Constrained Quadratic Programming

A quadratic program with equality constraints [1] is

$$\begin{aligned} & \text{minimize} && \frac{1}{2}x^\top Qx \end{aligned} \tag{1}$$

$$\begin{aligned} & \text{subject to} && Ax = b \end{aligned} \tag{2}$$

with variable in $x \in \mathbb{R}^n$, where $Q \succeq 0$ and the constraint $A \in \mathbb{R}^{m \times n}$ is, for our purposes, fat and full rank.

The ℓ^2 regularized version for $\lambda > 0$ looks like

$$\begin{aligned} & \text{minimize} && \frac{1}{2}x^\top Qx + \lambda x^\top x \end{aligned} \tag{3}$$

$$\begin{aligned} & \text{subject to} && Ax = b \end{aligned} \tag{4}$$

2.2 Karush-Kuhn-Tucker Conditions

The Lagrangian for (1) is of form

$$L(x, p) = \frac{1}{2}x^\top Qx + p^\top (Ax - b)$$

for which the optimality conditions [1] are

$$Qx^* + A^\top p^* = b, \quad Ax^* = b$$

or more compcatctly expressed:

$$\begin{bmatrix} Q & A^\top \\ A & 0 \end{bmatrix} \begin{bmatrix} x^* \\ p^* \end{bmatrix} = \begin{bmatrix} 0 \\ b \end{bmatrix} \quad (5)$$

On the other hand the Lagrangian of the regularized problem (3) is

$$L_\lambda(x, p) = \frac{1}{2}x^\top (Q + \lambda I)x + p^\top (Ax - b)$$

which has the KKT conditions

$$\begin{bmatrix} Q + \lambda I & A^\top \\ A & 0 \end{bmatrix} \begin{bmatrix} x_\lambda^* \\ p_\lambda^* \end{bmatrix} = \begin{bmatrix} 0 \\ b \end{bmatrix} \quad (6)$$

2.3 Matrix Inversion

Consider a symmetric matrix

$$Q = \begin{bmatrix} A & B \\ B^\top & C \end{bmatrix}$$

when $A \succ 0$, define the Schur complement $S = C - B^\top A^{-1}B$. Then

$$\begin{bmatrix} A & B \\ B^\top & C \end{bmatrix}^{-1} = \begin{bmatrix} A^{-1} + A^{-1}BS^{-1}B^\top A^{-1} & -A^{-1}BS^{-1} \\ -S^{-1}B^\top A^{-1} & S^{-1} \end{bmatrix}$$

3 Bounding Norms

Noting that by linearity

$$\begin{bmatrix} Q & A^\top \\ A & 0 \end{bmatrix} \begin{bmatrix} x^* \\ p^* \end{bmatrix} + \begin{bmatrix} \lambda I & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} x^* \\ p^* \end{bmatrix} = \begin{bmatrix} Q + \lambda I & A^\top \\ A & 0 \end{bmatrix} \begin{bmatrix} x^* \\ p^* \end{bmatrix} = \begin{bmatrix} \lambda x^* \\ b \end{bmatrix}$$

and so through subtracting equations,

$$\begin{bmatrix} Q + \lambda I & A^\top \\ A & 0 \end{bmatrix} \begin{bmatrix} x^* - x_\lambda^* \\ p^* - p_\lambda^* \end{bmatrix} = \begin{bmatrix} \lambda x^* \\ 0 \end{bmatrix}$$

Given our assumptions on $Q + \lambda I \succ 0$ and A is full rank. For ease of notation, define $\Lambda = Q + \lambda I$. The Schur complement is then $S = -\Lambda^{-1}A^\top$, and

$$\begin{bmatrix} x^* - x_\lambda^* \\ p^* - p_\lambda^* \end{bmatrix} = \begin{bmatrix} \Lambda^{-1} + \Lambda^{-1}A^\top S^{-1}A\Lambda^{-1} & -\Lambda^{-1}A^\top S^{-1} \\ -S^{-1}A\Lambda^{-1} & S^{-1} \end{bmatrix} \begin{bmatrix} \lambda x^* \\ 0 \end{bmatrix}$$

In other words:

$$x^* - x_\lambda^* = (\Lambda^{-1} - \Lambda^{-1}A^\top(A\Lambda^{-1}A^\top)^{-1}A\Lambda^{-1})\lambda x^*$$

To simplify notation slightly, use $\Gamma = \Lambda^{-1}$ because it looks like an upside-down L . Then

$$x^* - x_\lambda^* = (\Gamma - \Gamma A^\top(A\Gamma A^\top)^{-1}A\Gamma)\lambda x^*$$

Theorem 1. *Without loss of generality, assume that*

$$Q = \begin{bmatrix} q_1 & & \\ & \ddots & \\ & & q_n \end{bmatrix}, \quad q_1 \geq \dots \geq q_n > 0.$$

Then for $\lambda \leq q_n$,

$$\|x^* - x_\lambda^*\| \leq \left(\frac{\lambda}{q_n + \lambda} - \frac{\lambda}{q_1 + \lambda} \right) \|x^*\|$$

Proof. First note that

$$\Lambda = \begin{bmatrix} q_1 + \lambda & & \\ & \ddots & \\ & & q_n + \lambda \end{bmatrix}, \quad \Gamma = \Lambda^{-1} = \begin{bmatrix} \frac{1}{q_1 + \lambda} & & \\ & \ddots & \\ & & \frac{1}{q_n + \lambda} \end{bmatrix}$$

We seek the bound the RHS of the inequality

$$\|x^* - x_\lambda^*\| \leq \|\Gamma - \Gamma A^\top(A\Gamma A^\top)^{-1}A\Gamma\| \cdot \lambda \|x^*\|$$

The primary challenge here is correctly *lower bounding* the $\Gamma A^\top(A\Gamma A^\top)^{-1}A\Gamma$ term. For this we *upper bound* the $A\Gamma A^\top$ term. Letting

$$A = U\Sigma V^\top = U \begin{bmatrix} \Sigma_1 & 0 \end{bmatrix} \begin{bmatrix} V_1^\top \\ V_2^\top \end{bmatrix}, \quad \Sigma_1 = \begin{bmatrix} \sigma_1 & & \\ & \ddots & \\ & & \sigma_m \end{bmatrix}$$

be an SVD of A ; an upper bound of $A\Gamma A^\top$ is

$$A\Gamma A^\top = U\Sigma_1 V_1^\top \begin{bmatrix} \frac{1}{q_1 + \lambda} & & \\ & \ddots & \\ & & \frac{1}{q_n + \lambda} \end{bmatrix} V_1 \Sigma_1 U^\top \preceq \frac{1}{q_n + \lambda} U\Sigma_1^2 U^\top$$

Consequently, a lower bound for $\Gamma A^\top(A\Gamma A^\top)^{-1}A\Gamma$ is

$$\Gamma A^\top(A\Gamma A^\top)^{-1}A\Gamma \succeq \Gamma V_1 \Sigma_1 U^\top \left(\frac{1}{q_n + \lambda} U\Sigma_1^2 U^\top \right)^{-1} U\Sigma_1 V_1^\top \Gamma = (q_n + \lambda) \Gamma V_1 V_1^\top \Gamma \succeq \frac{q_n + \lambda}{q_1 + \lambda} \Gamma$$

Then using this,

$$\Gamma - \Gamma A^\top(A\Gamma A^\top)^{-1}A\Gamma \preceq \Gamma - \frac{q_n + \lambda}{q_1 + \lambda} \Gamma \preceq \left(\frac{1}{q_n + \lambda} - \frac{1}{q_1 + \lambda} \right) I$$

for which we derive the desired bound

$$\|x^* - x_\lambda^*\| \leq \|\Gamma - \Gamma A^\top(A\Gamma A^\top)^{-1}A\Gamma\| \cdot \lambda \|x^*\| \leq \left(\frac{\lambda}{q_n + \lambda} - \frac{\lambda}{q_1 + \lambda} \right) \|x^*\|$$

□

4 Optimal Value Deviation

One challenge with merely bounding $\|x^* - x_\lambda^*\|$ is that this norm difference may grow unbounded. Consider an instance where the objective Q has a non-trivial kernel: we may have

$$\|\Pi_{\ker Q} x^*\| \gg \|\Pi_{\ker Q} x_\lambda^*\|$$

where $\Pi_{\ker Q}$ is the projection onto $\ker Q$; this is possible because the regularized problems would force $\Pi_{\ker Q} x_\lambda^*$ to be close to the origin rather than hiding far away from zero inside $\ker Q$. Instead, another way of examining sub-optimality is to consider the difference

$$\frac{1}{2}(x_\lambda^*)^\top Q x_\lambda^* - \frac{1}{2}(x^*)^\top Q x^* > 0$$

which we know holds because x_λ^* is no more optimal than x^* .

Theorem 2 ([2]). *If X is an $n \times q$ matrix contained in the column-space of an $n \times n$ symmetrical matrix V , then*

$$(V + XX^\top)^\dagger = V^\dagger - V^\dagger X(I + X^\top V^\dagger X)^{-1} X^\top V^\dagger$$

where “to be in the column-space” means the same as $VV^\dagger X = X$, and in a more general case $(I - VV^\dagger)X \neq 0$.

Theorem 3. *Define*

$$Z = I - A^\dagger A, \quad \hat{x} = A^\dagger b$$

so that solutions take form $\hat{x} + Zv$, then a bound on the optimality gap is

$$\frac{1}{2}(x_\lambda^*)^\top Q x_\lambda^* - \frac{1}{2}(x^*)^\top Q x^* \leq O\left(\frac{\lambda q_0}{q_0 - \lambda}\right)$$

where q_0 is the smallest non-zero eigenvalue of Q and $\lambda < q_0$.

Proof. Let $\hat{x} = A^\dagger b$, and $\Lambda = Q + \lambda I$, then

$$x^* = \hat{x} - Zu, \quad x_\lambda^* = \hat{x} - Zy$$

where

$$Z = I - A^\dagger A, \quad u = (Z^\top M Z)^\dagger Z^\top Q \hat{x}, \quad y = (Z^\top \Lambda Z)^\dagger Z^\top \Lambda \hat{x}$$

which can be derived from examining the optimality conditions of the quadratic problem. However, because $\hat{x} \in \text{ran } A^\top = (\ker A)^\perp$, there is a further simplification as $Z^\top \hat{x} = 0$:

$$y = (Z^\top \Lambda Z)^\dagger Z^\top (Q + \lambda I) \hat{x} = (Z^\top \Lambda Z)^\dagger Z^\top Q \hat{x}$$

Comparing the objective values attained by x_λ^\star and x^\star , we have

$$\begin{aligned}
& \frac{1}{2}(x_\lambda^\star)^\top Q x_\lambda^\star - \frac{1}{2}(x^\star)^\top Q x^\star \\
&= \frac{1}{2}(\hat{x} + Zy)^\top Q(\hat{x} + Zy) - \frac{1}{2}(\hat{x} + Zu)^\top Q(\hat{x} + Zu) \\
&= (\hat{x})^\top QZ(y - u) + \frac{1}{2}y^\top Z^\top QZy - \frac{1}{2}u^\top Z^\top QZu \\
&= (\hat{x})^\top QZ(y - u) + \frac{1}{2}\left(\|Q^{1/2}Zy\|^2 - \|Q^{1/2}Zu\|^2\right) \\
&= (\hat{x})^\top QZ(y - u) + \frac{1}{2}\left[(\|Q^{1/2}Zu\| + \|Q^{1/2}Zy\|)(\|Q^{1/2}Zy\| - \|Q^{1/2}Zu\|)\right] \\
&\leq (\hat{x})^\top QZ(y - u) + \frac{1}{2}\left[(\|Q^{1/2}Zu\| + \|Q^{1/2}Zy\|) \cdot \|Q^{1/2}Z(y - u)\|\right] \quad (\text{Reverse triangle})
\end{aligned}$$

Which simplifies to

$$\frac{1}{2}(x_\lambda^\star)^\top Q x_\lambda^\star - \frac{1}{2}(x^\star)^\top Q x^\star \leq \left(\|Q^{1/2}\hat{x}\| + \frac{1}{2}\|Q^{1/2}Zu\| + \frac{1}{2}\|Q^{1/2}Zy\|\right) \cdot \|Q^{1/2}Z(y - u)\| \quad (7)$$

We now apply to Theorem 2 to $Q^{1/2}Zy$: identifying the mapping $V \mapsto Z^\top QZ$ and $X \mapsto \sqrt{\lambda}Z$, first show that X lies in the column space of V . For this, consider the eigen decomposition

$$V = FDF^\top, \quad F = \begin{bmatrix} F_1 & F_2 \end{bmatrix}, \quad D = \text{diag}(d_1, \dots, d_k, 0, \dots, 0), \quad X = \sqrt{\lambda}F_1$$

where F is orthogonal. Then $D^\dagger = \text{diag}(d_1^{-1}, \dots, d_k^{-1}, 0, \dots, 0)$ and so

$$VV^\dagger X = (FDF^\top)(FD^\dagger F^\top)\sqrt{\lambda}F_1 = \sqrt{\lambda}F_1 = X$$

thus, Theorem 2 applies and

$$\begin{aligned}
Q^{1/2}Zy &= Q^{1/2}Z(Z^\top(Q + \lambda I)Z)^\dagger Z^\top Q\hat{x} \\
&= Q^{1/2}Z[V^\dagger - V^\dagger X(I + X^\top V^\dagger X)^{-1}X^\top V^\dagger]Z^\top Q\hat{x}
\end{aligned}$$

With this, we now bound the terms that appear in (7). For the easy one:

$$\|Q^{1/2}\hat{x}\| = \|Q^{1/2}A^\dagger b\| \leq \lambda_{\max}(Q)^{1/2} \frac{1}{\sigma_{\min}(A)} \|b\| \quad (8)$$

For the inside second term, accounting for the fact that Z is a projection,

$$\|Q^{1/2}Zu\| = \|Q^{1/2}ZV^\dagger Z^\top QA^\dagger b\| \leq \lambda_{\max}(Q)^{3/2} \frac{1}{q_0 \sigma_{\min}(A)} \|b\| \quad (9)$$

where q_0 is the smallest non-zero eigenvalue of Q . For the third inside term:

$$\begin{aligned}
& \|Q^{1/2}Zy\| \\
&= \|Q^{1/2}Z[V^\dagger - V^\dagger X(I + X^\top V^\dagger X)^{-1}X^\top V^\dagger]Z^\top QA^\dagger b\| \\
&\leq \|Q^{1/2}ZV^\dagger Z^\top QA^\dagger b\| + \|Q^{1/2}ZV^\dagger X(I + X^\top V^\dagger X)^{-1}X^\top V^\dagger Z^\top QA^\dagger b\| \\
&\leq \|Q^{1/2}ZV^\dagger Z^\top QA^\dagger b\| + \|Q\|^{3/2}\|V^\dagger\|^2 \cdot \|X\|^2 \cdot \|A^\dagger b\| \cdot \|(I + X^\top V^\dagger X)^{-1}\| \\
&\leq \frac{\lambda_{\max}(Q)^{3/2} \cdot \|b\|}{q_0 \sigma_{\min}(A)} + \frac{\lambda_{\max}(Q)^{3/2} \cdot \|b\| \lambda}{q_0^2 \sigma_{\min}(A)} \cdot \|(I + X^\top V^\dagger X)^{-1}\| \\
&\leq \frac{\lambda_{\max}(Q)^{3/2} \cdot \|b\|}{q_0 \sigma_{\min}(A)} + \frac{\lambda_{\max}(Q)^{3/2} \cdot \|b\| \lambda}{q_0^2 \sigma_{\min}(A)} \sum_{l=0}^{\infty} \|X^\top V^\dagger X\|^l \\
&\leq \frac{\lambda_{\max}(Q)^{3/2} \cdot \|b\|}{q_0 \sigma_{\min}(A)} + \frac{\lambda_{\max}(Q)^{3/2} \cdot \|b\| \lambda}{q_0^2 \sigma_{\min}(A)} \sum_{l=0}^{\infty} (\lambda/q_0)^l \\
&\leq \frac{\lambda_{\max}(Q)^{3/2} \cdot \|b\|}{q_0 \sigma_{\min}(A)} + \frac{\lambda_{\max}(Q)^{3/2} \cdot \|b\| \lambda}{q_0^2 \sigma_{\min}(A)} \left(1 - \frac{\lambda}{q_0}\right)^{-1}
\end{aligned}$$

which requires $\lambda < q_0$ for convergence. Finally for the right outer term:

$$\|Q^{1/2}Z(y - u)\| = \|Q^{1/2}ZV^\dagger X(I + X^\top V^\dagger X)^{-1}X^\top V^\dagger Z^\top QA^\dagger b\| \leq \frac{\lambda_{\max}(Q)^{3/2}\|b\|\lambda}{q_0^2 \sigma_{\min}(A)} \left(1 - \frac{\lambda}{q_0}\right)^{-1}$$

Putting these together, we find that

$$\begin{aligned}
& \frac{1}{2}(x_\lambda^*)^\top Qx_\lambda^* - \frac{1}{2}x^*Qx^* \\
&\leq \left(\|Q^{1/2}\hat{x}\| + \frac{1}{2}\|Q^{1/2}Zu\| + \frac{1}{2}\|Q^{1/2}Zy\|\right) \cdot \|Q^{1/2}Z(y - u)\| \\
&\leq \max\{\lambda_{\max}(Q)^2, \lambda_{\max}(Q)^3\} \cdot \max\{q_0^{-2}, q_0^{-3}, q_0^{-4}\} \cdot \left(\frac{\|b\|}{\sigma_{\min}(A)}\right)^2 O\left(\frac{\lambda q_0}{q_0 - \lambda}\right)
\end{aligned}$$

□

References

- [1] Stephen P Boyd and Lieven Vandenberghe. *Convex Optimization*. Cambridge University Press, 2004.
- [2] Pavel Kovanic. “On the pseudoinverse of a sum of symmetric matrices with applications to estimation”. In: *Kybernetika* 15.5 (1979), pp. 341–348.