# Constrained Linear Quadratic Regular Optimization

#### Anton Xue

#### 1 Introduction

We look at constrained linear quadratic regulator (LQR) optimization.

## 2 Background

Relevant background on linear systems, LQR, and matrix magic.

#### 2.1 Linear Systems and the Linear Quadratic Regulator

Linear system dynamics

$$x_{t+1} = Ax_t + Bu_t, \qquad (x_t)_{t=0}^{\infty} \in \mathbb{R}^n, \quad (u_t)_{t=0}^{\infty} \in \mathbb{R}^p,$$

Finite horizion linear quadratic regular (LQR) cost function

$$J = \sum_{t=0}^{T} x_t^{\top} Q x_t + u_t^{\top} R u_t, \qquad Q \succeq 0, \quad R \succ 0,$$

which is finite when the system dynamics  $(x_t, u_t)$  tends towards the origin. Infinite-horizon LQR average-cost function

$$J = \limsup_{T \to \infty} \frac{1}{T} \sum_{t=0}^{T-1} x_t^{\top} Q x_t + u_t^{\top} R u_t, \qquad Q \succeq 0, \quad R \succ 0,$$

which allows for systems whose stability is not necessarily at the origin. Both versions of LQR admit an optimal feedback control sequence that is linear, i.e., u = Kt, with respect to the algebraic Riccati equation in P where

$$P = Q + A^{T}PA - A^{T}PB(R + B^{T}PB)^{-1}B^{T}PA, \qquad K = -(R + B^{T}PB)^{-1}B^{T}PA$$

where the solution for  $P \succ 0$ , if it exists, is unique.

#### 2.2 Affine Systems

Given the affine system

$$x_{t+1} = Ax_t + Bu_t + c$$

we can reformulate it as a linear system

$$z_{t+1} = \begin{bmatrix} x_{t+1} \\ 1 \end{bmatrix} = \begin{bmatrix} A & c \\ 0 & 1 \end{bmatrix} \begin{bmatrix} x_t \\ 1 \end{bmatrix} + \begin{bmatrix} B \\ 0 \end{bmatrix} u_t = \tilde{A}z_t + \tilde{B}u_t$$
 (1)

#### 2.3 Completing the Square

For the matrix case with variable in x from Wikipedia [2],

$$x^{\top}Qx + p^{\top}x + r = (x - h)^{\top}Q(x - h) + k, \qquad h = -\frac{1}{2}Q^{-1}p, \qquad k = r - \frac{1}{4}p^{\top}Q^{-1}p$$
 (2)

where Q is assumed to be invertible.

#### 3 Some Results with Affine Constraints

**Theorem 1.** An infinite horizon average-cost LQR problem with quadratic-plus-affine terms

minimize 
$$\limsup_{T \to \infty} \frac{1}{T} \sum_{t=0}^{T-1} x_t^{\top} Q x_t + q^{\top} x_t + q_0 + u_t^{\top} R u_t + r^{\top} u_t + r_0$$
subject to 
$$x_{t+1} = A x_t + B u_t, \text{ for all } t$$

admits an equivalent infinite horizon average-cost LQR problem without affine terms.

*Proof.* Applying (2) we can write the objective as

$$\limsup_{T \to \infty} \frac{1}{T} \sum_{t=0}^{T-1} (x_t - h)^{\top} Q(x_t - h) + (u_t - l)^{\top} R(u_t - l) + k,$$

with coefficients

$$h = -\frac{1}{2}Q^{-1}q, \qquad l = -\frac{1}{2}R^{-1}r, \qquad k = q_0 + r_0 - \frac{1}{4}(q^{\top}Qq + r^{\top}Rr)$$

Observe that k is constant with respect to the variables  $(x_t, u_t)$  and the objective is equivalently

$$k + \limsup_{T \to \infty} \frac{1}{T} \sum_{t=0}^{T-1} (x_t - h)^{\top} Q(x_t - h) + (u_t - l)^{\top} R(u_t - l).$$

Define the change-of-variables  $\tilde{x}_t = x_t - h$  and  $\tilde{u}_t = u_t - l$  for all t, which induces the affine system

$$\tilde{x}_{t+1} = A\tilde{x}_t + B\tilde{u}_t + (Ah + Bl - h)$$

and after the appropriate change of variables to recover a linear system as in (1),

$$z_{t+1} = \begin{bmatrix} A & (Ah + Bl - h) \\ 0 & 1 \end{bmatrix} \begin{bmatrix} \tilde{x}_t \\ 1 \end{bmatrix} + \begin{bmatrix} B \\ 0 \end{bmatrix} \tilde{u}_t = \tilde{A}z_t + \tilde{B}\tilde{u}_t.$$

For this, the reduced problem is then

minimize 
$$k + \limsup_{T \to \infty} \frac{1}{T} \sum_{t=0}^{T-1} z_t^{\top} \begin{bmatrix} Q & 0 \\ 0 & 0 \end{bmatrix} z_t + \tilde{u}_t^{\top} R \tilde{u}_t$$
  
subject to  $z_{t+1} = \begin{bmatrix} A & (Ah + Bl - h) \\ 0 & 1 \end{bmatrix} z_t + \begin{bmatrix} B \\ 0 \end{bmatrix} \tilde{u}_t$ , for all  $t$ 

with variables in  $(z_t)_{t=0}^{\infty}$  and  $(u_t)_{t=0}^{\infty}$ . Barring the k offset in the objective, this is now equivalent to an infinite horizon average-cost LQR problem with no affine terms.

Because quadratic-plus-affine terms can be eliminated to yield a problem with purely quadratic cost, Theorem 1 implies that quadratic systems also have an optimal feedback control sequence that is linear for the equivalent system with trajectory in  $(z_t, \tilde{u}_t)_{t=0}^{\infty}$ . But since  $(z_t, \tilde{u}_t)$  are linear transformations of the original trajectory  $(x_t, u_t)$ , the optimal feedback policy for the original dynamics is also linear.

**Theorem 2.** The infinite-horizon average-cost LQR problem with affine constraints

minimize 
$$\limsup_{T \to \infty} \frac{1}{T} \sum_{t=0}^{T-1} x_t^{\top} Q x_t + u_t^{\top} R u_t$$
subject to 
$$x_{t+1} = A x_t + B u_t, \text{ for all } t$$
$$C x_t = d, \text{ for all } t$$

where C is fat and full rank, admits an optimal feedback control policy that is linear if:

- There exists  $\hat{u}$  such that  $\hat{x} = A\hat{x} + B\hat{u}$ , where  $\hat{x} = C^{\dagger}d$ .
- $\operatorname{ran} AN \subseteq \operatorname{ran} B$ , where  $\operatorname{ran} N = \ker C$ .

*Proof.* Supposing the assumptions, we can rewrite our system dynamics as

Defining the following for compactness,

$$J(z_t, v_t) = \begin{bmatrix} \hat{x} \\ Nz_t \end{bmatrix}^{\top} \begin{bmatrix} Q & 0 \\ 0 & Q \end{bmatrix} \begin{bmatrix} \hat{x} \\ Nz_t \end{bmatrix} + \begin{bmatrix} 0 \\ 2Q\hat{x} \end{bmatrix} \begin{bmatrix} \hat{x} \\ Nz_t \end{bmatrix} + \begin{bmatrix} \hat{u} \\ v_t \end{bmatrix}^{\top} \begin{bmatrix} R & 0 \\ 0 & R \end{bmatrix} \begin{bmatrix} \hat{u} \\ v_t \end{bmatrix} + \begin{bmatrix} 0 \\ 2R\hat{x} \end{bmatrix} \begin{bmatrix} \hat{u} \\ v_t \end{bmatrix}$$

makes it so that

$$x_t^{\top} Q x_t + u_t^{\top} R u_t = (\hat{x} + N z_t)^{\top} Q (\hat{x} + N z_t) + (\hat{u} + v_t)^{\top} R (\hat{u} + v_t) = J(z_t, v_t)$$

and that the objective can now be written as

minimize 
$$\limsup_{T \to \infty} \frac{1}{T} \sum_{t=0}^{T-1} J(z_t, v_t)$$
  
subject to (3)

with variables in  $(z_t)_{t=0}^{\infty}$  and  $(v_t)_{t=0}^{\infty}$ . Since J is a quadratic-plus-affine decomposition of the state and control costs, we know that this infinite-horizon average-cost LQR problem has an optimal feedback controller that is linear.

### References

- [1] Stephen Boyd. "Ee363: Linear dynamical systems". In: Lecture. Stanford Univer., Winter Quarter 9 (2008). URL: https://stanford.edu/class/ee363/lectures/stoch\_lqr.pdf.
- [2] Wikipedia contributors. Completing the square Wikipedia, The Free Encyclopedia. [Online; accessed 14-May-2020]. 2020. URL: https://en.wikipedia.org/w/index.php?title=Completing\_the\_square&oldid=953923455.