
Smoothing of cosmic-ray background models

ERLANGEN CENTRE FOR ASTROPARTICLE PHYSICS

Gammapy Coding Sprint 15.07.2019, Erlangen Andreas Specovius










Spectromorphological bg template






- Bg from archival H.E.S.S. data
- Predict the residual hadronic bg
 rate per solidangle, energy and time
 as a function of x, y and energy
- Subject to statistical fluctuations
 - Get rid of them
 - -> smooth
 - Ensure:
 - Preserve spatial curvature
 - Positive rates





Remove statistical fluctuations





 Total Variation denoising: Minimizes L1 norm of data gradient

skimage.restoration.denoise tv chambolle

2-dim Gaussian Filter:
 Convolution with gaussian filter

scipy.ndimage.gaussian_filter

Kernel Density Estimation:
 Probability density estimation using gaussian kernel

scipy.stats.gaussian kde


• 2-dim 3rd order **Spline**


scipy.signal.cspline2d(data, par)


 Spherical harmonics (analytical description)


Preserve the shape















Real raw data











































Smoothing of cosmic-ray background models

ERLANGEN CENT Questions?
FOR ASTROPARTICLE

Gammapy Coding Sprint 15.07.2019, Erlangen Andreas Specovius







3D template analysis



