

- ▶ This talk is about regression methods in which the dependent variable takes nonnegative integer or count values.
- ▶ The dependent variable is usually the number of times an event occurs.

Overview

Some examples of event counts are:

- ▶ number of claims per year on a particular car owners insurance policy,
- ▶ number of workdays missed due to sickness of a dependent in a one-year period,
- ▶ number of papers published per year by a researcher.

Poisson Distribution: Prussian Cavalry

- ▶ The classic Poisson example is the data set of von Bortkiewicz (1898), for the chance of a Prussian cavalryman being killed by the kick of a horse.
- ▶ Ten army corps were observed over 20 years, giving a total of 200 observations of one corps for a one year period. The period or module of observation is thus one year.

Poisson Distribution: Prussian Cavalry

- ▶ The total deaths from horse kicks were 122, and the average number of deaths per year per corps was thus $122/200 = 0.61$.
- ▶ In any given year, we expect to observe, well, not exactly 0.61 deaths in one corps
- ▶ Here, then, is the classic Poisson situation: a rare event, whose average rate is small, with observations made over many small intervals of time.

Overview

- ▶ Poisson regression is main technique used to model count variables.
- ▶ Sometimes conventional Poisson Regression is not an appropriate technique, and alternative or variant techniques are used instead.
- ▶ Negative Binomial regression is for modelling count variables, usually for over-dispersed count outcome variables.

Generalized Linear Models

- ▶ In statistics, the problem of modelling count variables is an example of generalized linear modelling.
- ▶ Generalized linear models are fit using the `glm()` function.
- ▶ The form of the `glm` function is

```
glm(formula, family=familytype(link=linkfunction),  
     data=dataname)
```

Generalized Linear Models

Family	Default Link Function
binomial	(link = "logit")
gaussian	(link = "identity")
Gamma	(link = "inverse")
inverse.gaussian	(link = " $1/\mu^2$ ")
poisson	(link = "log")
quasibinomial	(link = "logit")
quasipoisson	(link = "log")

Texts on GLMs

- ▶ Dobson, A. J. (1990) An Introduction to Generalized Linear Models. (*London: Chapman and Hall.*)
- ▶ Hastie, T. J. and Pregibon, D. (1992) Generalized linear models. Chapter 6 of Statistical Models in S eds J. M. Chambers and T. J. Hastie, Wadsworth & Brooks/Cole.
- ▶ McCullagh P. and Nelder, J. A. (1989) Generalized Linear Models. (*London: Chapman and Hall.*)
- ▶ Venables, W. N. and Ripley, B. D. (2002) Modern Applied Statistics with S. *New York: Springer.*

VGAM: Vector Generalized Linear and Additive Models

Author(s): Thomas W. Yee (t.yee@auckland.ac.nz)

License: GPL-2

URL: <http://www.stat.auckland.ac.nz/~yee/VGAM>

Vector generalized linear and additive models, and associated models (Reduced-Rank VGLMs, Quadratic RR-VGLMs, Reduced-Rank VGAMs).

This package fits many models and distribution by maximum likelihood estimation (MLE) or penalized MLE. Also fits constrained ordination models in ecology.