

# MedCore Pharmaceuticals — Clinical Dosage Report

Document ID: MCP-2025-447 | Classification: Confidential

## 1. Drug Overview

Drug Name: Veranidol (generic: veranidol hydrochloride)

Therapeutic Class: Non-opioid analgesic / anti-inflammatory

Approval Status: FDA Approved — March 2023

## 2. Dosage Guidelines

Adult Patients (18–64 years): 200mg orally, twice daily (every 12 hours). Maximum daily dose must not exceed 400mg. Treatment duration should not exceed 14 consecutive days without physician review.

Elderly Patients (65+ years): Reduce dose to 100mg orally, twice daily. Renal function must be monitored every 7 days. Do not exceed 200mg per day.

Pediatric Patients (under 18): Veranidol is strictly contraindicated for patients under 18 years of age.

## 3. Administration Instructions

Administer with food to reduce gastrointestinal side effects. Tablets must not be crushed or chewed. If a dose is missed, take as soon as remembered unless within 6 hours of the next scheduled dose.

## 4. Contraindications

Do not administer Veranidol to patients with: (a) known hypersensitivity to veranidol hydrochloride, (b) severe hepatic impairment (Child-Pugh Class C), (c) active peptic ulcer disease, (d) pregnancy in the third trimester.

## 5. Side Effects

Common (>10% incidence): nausea, headache, dizziness. Serious (report immediately): gastrointestinal bleeding, elevated liver enzymes (ALT >3x ULN), allergic reaction.

## 6. Storage

Store at 20–25°C (68–77°F). Keep away from moisture and direct sunlight. Discard unused medication after expiry date printed on packaging.