



Chemo-Scale™ - Chemo Sensitivity Test
PATIENT'S DETAILS AND CLINICAL HISTORY

PATIENT NAME	: Mr. A. B. C.	TUMOR TYPE	: Pancreatic ductal adenocarcinoma
AGE	: 64 Years	SPECIMEN TYPE	: Blood
GENDER	: Male	DATE OF COLLECTION	: 21.02.2022
ADDRESS	: ---	DATE OF ACCESSION	: 25.02.2022
REFERRED BY	: Dr. X. Y. Z.	DATE OF REPORT	: 07.03.2022

CHEMO-SCALE ASSAY RESULTS


THIS REPORT SHOULD BE READ AS A WHOLE AND USED AND ACTED UPON BY A REGISTERED MEDICAL PRACTITIONER ONLY.
 THIS IS NOT A PRESCRIPTION.


Chemo-Scale™ - Chemo Sensitivity Test

Chemosensitivity assay performed on circulating tumor and its associated cells indicates the effectiveness of chemotherapeutic drugs in descending order of efficacy.

RESPONSE TO DRUGS ALREADY ADMINISTERED TO PATIENT

Sr. No.	Drugs	% Cell Death
1	5-Fluorouracil/Capecitabine + Irinotecan + Oxaliplatin	Not tested
3	5-Fluorouracil/Capecitabine	<25
4	Irinotecan	42
5	Oxaliplatin	<25
2	Gemcitabine + Paclitaxel	Not tested
6	Gemcitabine	39
7	Paclitaxel	55

INTERPRETATION AND RECOMMENDATIONS

Among the panel of cytotoxic drugs tested, patient's tumor cells showed high response to Doxorubicin, Docetaxel, Etoposide, Paclitaxel and Cyclophosphamide (see graph on page 1).

METHODS AND LIMITATIONS

Circulating tumor and its associated cells were isolated from the submitted peripheral blood sample. Live cancer cells were tested against multiple chemotherapy agents. The number of drugs selected for testing depend on the number of circulating tumor associated cells isolated from the submitted sample.

A defined number of cells were incubated with different drugs with respective drug concentrations, mean peak plasma concentration and cell death events were measured. The extent of cell death was determined either using Varioskan LUX platform or by fluorescence-based staining of live / dead cells. Percent cell death was calculated to evaluate the response level of the drug.

Appropriate positive and negative controls were tested and evaluated in a similar manner simultaneously with the test sample and performed satisfactorily.

Analytical Validation of this assay shown sensitivity of 85.71% and specificity 99.99%.

This test was developed and its performance characteristics determined by Datar Cancer Genetics. It has not been cleared or approved by the U.S. Food and Drug Administration.

**THIS REPORT SHOULD BE READ AS A WHOLE AND USED AND ACTED UPON BY A REGISTERED MEDICAL PRACTITIONER ONLY.
THIS IS NOT A PRESCRIPTION.**


Chemo-Scale™ - Chemo Sensitivity Test

This laboratory is certified under the Clinical Laboratory Improvement Amendments (CLIA)-USA as qualified to perform high complexity clinical laboratory testing.

LIST OF DRUGS TESTED

5-Fluorouracil/Capecitabine, Bleomycin, Cabazitaxel, Carboplatin, Cisplatin, Cyclophosphamide, Dacarbazine, Dactinomycin, Docetaxel, Doxorubicin, Epirubicin, Eribulin, Etoposide, Gemcitabine, Ifosfamide, Irinotecan, Melphalan, Methotrexate, Mitomycin, Mitoxantrone, Oxaliplatin, Paclitaxel, Pemetrexed, Temozolomide, Topotecan, Trabectedin, Vinblastine, Vincristine, Vinorelbine.

DISCLAIMER

Chemo-scale test report is not intended to be a substitute for care by a licensed healthcare professional. The information provided in this report is not a substitute for medical advice, diagnosis, treatment or a full medical evaluation. All laboratory test results must be interpreted within the context of a patient's overall health and should be used along with other examinations or tests. Datar Cancer Genetic is not responsible for and does not accept the liability for any direct or indirect loss or damages arising from, or related to, the use of this information. This is not a prescription.

Cancer therapy selection, dosing, administration, and the management of related adverse events can be a complex process that should be handled by an experienced healthcare team. Clinicians must choose and verify treatment options based on the individual patient; drug dose modifications and supportive care interventions should be administered accordingly. The suggested treatment regimens may include both USFDA-approved and unapproved indications/regimens. Therapy Recommendations are a work in progress that may be refined as often as new significant data becomes available. Any clinician seeking to apply or consult DCGL's Therapy Recommendations is expected to use independent medical judgment in the context of individual clinical circumstances to determine any patient's care or treatment. DCGL makes no warranties of any kind whatsoever regarding their content, use, or application and disclaims any responsibility for their application or use in any way.

Results of Chemo-scale test are based on in-vitro testing of cancer cells under controlled laboratory conditions and may not exactly replicate in-vivo (within the patient's body).

This report should be read as a whole and used and acted upon by a registered medical practitioner only. This is not a prescription.

**THIS REPORT SHOULD BE READ AS A WHOLE AND USED AND ACTED UPON BY A REGISTERED MEDICAL PRACTITIONER ONLY.
THIS IS NOT A PRESCRIPTION.**


Chemo-Scale™ - Chemo Sensitivity Test
REFERENCES

- 1 Akolkar DB, Crook T, Page R, Patil D, Limaye S, Datta V, et al. Liquid biopsies to enable non-invasive real-time functional chemoresistance profiling in solid organ cancers. Journal of Clinical Oncology 38, no. 15_suppl (May 20, 2020) 3525-3525.
- 2 Crook T, Gaya A, Page R, Limaye S, Ranade A, Bhatt A, et al. Clinical utility of circulating tumor-associated cells to predict and monitor chemo-response in solid tumors. Cancer chemotherapy and pharmacology. 2021 Feb;87(2):197-205.
- 3 Nagarkar R, Patil D, Crook T, Datta V, Bhalerao S, Dhande S, et al. Encyclopedic tumor analysis for guiding treatment of advanced, broadly refractory cancers: results from the RESILIENT trial. Oncotarget. 2019 Sep 24;10(54):5605.
- 4 NCCN Clinical practice guidelines in oncology. Pancreatic Adenocarcinoma. Version 1.2022.
- 5 Schuster S, Akolkar D, Patil S, Patil D, Datta V, Srinivasan A, et al. In vitro functional interrogation of viable circulating tumor associated cells (C-TACs) for evaluating platin resistance. Annals of Oncology. 2019 Oct 1;30:v793.

****End of Report****

Dr. Rahul Gosavi
PhD. (Medical Microbiology)
Molecular Biologist

Dr. Darshana Patil
M.D. Pathology, Master in
Molecular Oncology (CNIO)
Medical Director

Dr. Ashwini Ghaisas
MRCOG, PGD-Clinical research, CCRG
Director - Application