

chronicity of exposure are unknown, but may be particularly relevant to occupationally exposed populations such as cashiers,⁵ who handle receipts 40 or more hours per week.

Limitations include the small volunteer sample and loss of participants in the second simulation. However, urinary BPA concentrations at baseline were similar in the full and smaller groups and similar to the US population (1.83 µg/L).⁶ A larger study is needed to confirm our findings and evaluate the clinical implications.

Acknowledgments

Funding/Support: This project was supported by grant 2 T42 OH008416-05 from the Harvard-NIOSH Education and Research Center.

Role of the Sponsor: The Harvard-NIOSH Education and Research Center had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

Additional Contributions: We thank Lori Torf (Harvard School of Public Health cafeteria) for providing us with access to the cafeteria cash registers for the simulations. We also thank the Hauser staff for assistance with storage of samples and supplies, Greg Wagner, MD (Harvard School of Public Health), for his advice and support at the start of the study, and Xiaoyun Ye, MS, Xiaoliu Zhou, MS, Tao Jia, MS, and Joshua Kramer (all 4 with the CDC) for technical assistance in measuring urinary bisphenol A. None of the individuals acknowledged were compensated for their contributions.

References

1. Ehrlich S, Williams PL, Missmer SA, et al. Urinary bisphenol A concentrations and early reproductive health outcomes among women undergoing IVF. *Hum Reprod*. 2012; 27(12):3583–3592. [PubMed: 23014629]
2. Braun JM, Kalkbrenner AE, Calafat AM, et al. Impact of early-life bisphenol A exposure on behavior and executive function in children. *Pediatrics*. 2011; 128(5):873–882. [PubMed: 22025598]
3. Carwile JL, Ye X, Zhou X, Calafat AM, Michels KB. Canned soup consumption and urinary bisphenol A: a randomized crossover trial. *JAMA*. 2011; 306(20):2218–2220. [PubMed: 22110104]
4. Biedermann S, Tschudin P, Grob K. Transfer of bisphenol A from thermal printer paper to the skin. *Anal Bioanal Chem*. 2010; 398(1):571–576. [PubMed: 20623271]
5. Braun JM, Kalkbrenner AE, Calafat AM, et al. Variability and predictors of urinary bisphenol A concentrations during pregnancy. *Environ Health Perspect*. 2011; 119(1):131–137. [PubMed: 21205581]
6. Centers for Disease Control and Prevention, National Center for Environmental Health. [January 25, 2014] Fourth national report on human exposure to environmental chemicals, updated tables. 2013 Sep. <http://www.cdc.gov/exposurereport/>.