



Figure 3. Bootstrapped mediation model documenting that perceptions of greater moral wrongfulness mediated the effect of anodal tDCS on reducing intentions to commit aggression. * $p < .05$, ** $p < .01$. Extended data are presented in Figure 3-1, available at <https://doi.org/10.1523/JNEUROSCI.3317-17.2018.f3-1>.

DLPFC influences intentions to commit aggression, they do not negate the involvement of other prefrontal areas, such as the ventromedial and anterior prefrontal cortex, or of nonprefrontal areas, including the temporal cortex. Future studies using complementary noninvasive neurostimulation approaches, such as transcranial magnetic stimulation and high-definition tDCS, may elucidate the anatomical specificity of this effect and the complexity of the functional neuroanatomy of violent behavior.

There has been increasing discussion of biological interventions on antisocial and aggressive behavior in both children and adults (Gesch et al., 2002; Raine et al., 2015; Hübner and White, 2016). Our initial findings, which are limited to intentions to commit aggression and moral judgment, require extensive replication. Nevertheless, among other etiological mechanisms, the role of biological factors on the development of antisocial behavior, including aggression, has been increasingly acknowledged (Raine, 2002; Glenn and Raine, 2014; Latvala et al., 2015). It has been suggested that treatment programs will be improved by considering biological mechanisms that potentially regulate aggression (Beauchaine et al., 2008). Thus, it can be argued that further investigation of basic science trials on tDCS may potentially offer a promising new biological approach for reducing aggression, which is a major public health problem and a feature of a variety of mental disorders, including antisocial personality disorder, intermittent explosive disorder, conduct disorder, and borderline personality disorder (American Psychiatric Association, 2013).

Conclusion

Understanding the etiology of aggression and the development of new interventions are paramount to a public health approach to violence reduction (Butchart et al., 2004; Slutkin, 2017). This first known application of prefrontal tDCS to study intentions to commit aggression takes a modest step toward advancing knowledge about the neural mechanisms that regulate aggression. Findings provide experimental evidence for the role of the prefrontal cortex on both physical and sexual assault, and suggest how the brain may, in theory, be amenable to change using a noninvasive tool with transient and relatively minor adverse effects (Poreisz et al., 2007; Fertonani et al., 2015). Nevertheless, a stronger evidence base that includes more consistent findings, documentation of long-term beneficial effects, and a comprehensive effort to rule out potentially aversive side effects is required before this technique can be considered in practice to reduce the perpetration of aggressive acts.

References

Akman D, Normandeau A, Sellin T, Wolfgang M (1968) Towards the measurement of criminality in Canada: a replication study. *Acta Criminologica* 1:135–260. [CrossRef](#)

- Alegria AA, Radua J, Rubia K (2016) Meta-analysis of fMRI studies of disruptive behavior disorders. *Am J Psychiatr* 173:1119–1130. [CrossRef Medline](#)
- American Psychiatric Association (2013) *Diagnostic and statistical manual of mental disorders* (5th ed.). Washington, DC: American Psychiatric Association.
- Anderson SW, Bechara A, Damasio H, Tranel D, Damasio AR (1999) Impairment of social and moral behavior related to early damage in human prefrontal cortex. *Nat Neurosci* 2:1032–1037. [CrossRef Medline](#)
- Armstrong TA, Boutwell BB (2012) Low resting heart rate and rational choice: integrating biological correlates of crime in criminological theories. *J Crim Jus* 40:31–39. [CrossRef](#)
- Bang H, Ni L, Davis CE (2004) Assessment of blinding in clinical trials. *Control Clin Trials* 25:143–156. [CrossRef Medline](#)
- Beauchaine TP, Neuhaus E, Brenner SL, Gatzke-Kopp L (2008) Ten good reasons to consider biological processes in prevention and intervention research. *Dev Psychopathol* 20:745–774. [Medline](#)
- Bechara A, Damasio AR, Damasio H, Anderson SW (1994) Insensitivity to future consequences following damage to human prefrontal cortex. *Cognition* 50:7–15. [CrossRef Medline](#)
- Brower MC, Price BH (2001) Neuropsychiatry of frontal lobe dysfunction in violent and criminal behaviour: a critical review. *J Neurol Neurosurg Psychiatry* 71:720–726. [CrossRef Medline](#)
- Brunoni AR, Nitsche MA, Bolognini N, Bikson M, Wagner T, Merabet L, Edwards DJ, Valero-Cabre A, Rotenberg A, Pascual-Leone A, Ferrucci R, Priori A, Boggio PS, Fregni F (2012) Clinical research with transcranial direct current stimulation (tDCS): challenges and future directions. *Brain Stimul* 5:175–195. [CrossRef Medline](#)
- Butchart A, Phinney A, Check P, Villaveces A (2004) Preventing violence: a guide to implementing recommendations of the World Report on Violence and Health. Geneva, Switzerland: Department of Injuries and Violence Prevention, World Health Organization.
- Chen CY, Raine A, Chou KH, Chen IY, Hung D, Lin CP (2016) Abnormal white matter integrity in rapists as indicated by diffusion tensor imaging. *BMC Neurosci* 17:45. [CrossRef Medline](#)
- Committee for Proprietary Medicinal Products (CPMP) (2004) Committee for Proprietary Medicinal Products (CPMP): points to consider on adjustment for baseline covariates. *Statistics in medicine* 23:701–709. [CrossRef Medline](#)
- Cui J, Tkachenko O, Gogel H, Kipman M, Preer LA, Weber M, Divatia SC, Demers LA, Olson EA, Buchholz JL, Bark JS, Rosso IM, Rauch SL, Killgore WD (2015) Microstructure of frontoparietal connections predicts individual resistance to sleep deprivation. *Neuroimage* 106:123–133. [CrossRef Medline](#)
- Dalwani M, Sakai JT, Mikulich-Gilbertson SK, Tanabe J, Raymond K, McWilliams SK, Thompson LL, Banich MT, Crowley TJ (2011) Reduced cortical gray matter volume in male adolescents with substance and conduct problems. *Drug Alcohol Depend* 118:295–305. [CrossRef Medline](#)
- Damasio AR (2000) A neural basis for sociopathy. *Arch Gen Psychiatry* 57:128–129. [CrossRef](#)
- Damasio H, Grabowski T, Frank R, Galaburda AM, Damasio AR (1994) The return of Phineas Gage: clues about the brain from the skull of a famous patient. *Science* 264:1102–1105. [CrossRef Medline](#)
- Dambacher F, Schuhmann T, Lobbestael J, Arntz A, Brugman S, Sack AT (2015a) No effects of bilateral tDCS over inferior frontal gyrus on response inhibition and aggression. *PLoS One* 10:e0132170. [CrossRef Medline](#)
- Dambacher F, Schuhmann T, Lobbestael J, Arntz A, Brugman S, Sack AT (2015b) Reducing proactive aggression through non-invasive brain stimulation. *Soc Cogn Affect Neurosci* 10:1303–1309. [CrossRef Medline](#)
- Dewall CN, Finkel EJ, Lambert NM, Slotter EB, Bodenhausen GV, Pond RS Jr,