



# Lecture-18

## Course: Applied Data Science

### Exponential Smoothing

By

**Dr. Sibarama Panigrahi**

Assistant Professor, Department of Computer Sc. & Engineering  
National Institute of Technology, Rourkela, Odisha, 769008, India

**Mobile No.:** +91-7377302566

**Email:** panigrahis[at]nitrkl[dot]ac[dot]in  
panigrahi[dot]sibarama[at]gmail[dot]com

# Exponential Smoothing

- Exponential smoothing was proposed in the late 1950s (Brown, 1959; Holt, 1957; Winters, 1960).
- Forecasts produced using exponential smoothing methods are weighted averages of past observations, with the weights decaying exponentially as the observations get older.
- In other words, the more recent the observation the higher the associated weight.

Brown, R. G. (1959). *Statistical forecasting for inventory control*. McGraw/Hill.

Holt, C. C. (1957). *Forecasting seasonals and trends by exponentially weighted averages* (ONR Memorandum No. 52). Carnegie Institute of Technology, Pittsburgh USA. Reprinted in the *International Journal of Forecasting*, 2004.

Winters, P. R. (1960). Forecasting sales by exponentially weighted moving averages. *Management Science*, 6(3), 324–342.

# Simple Exponential Smoothing

- The simplest of the exponentially smoothing methods is naturally called **simple exponential smoothing** (SES).
- This method is suitable for forecasting data with no clear trend or seasonal pattern.
- The naïve and the average as possible methods for forecasting such data.




Figure 8.1: Exports of goods and services from Algeria from 1960 to 2017.

Using the naïve method, all forecasts for the future are equal to the last observed value of the series,

$$\hat{y}_{T+h|T} = y_T,$$

Using the average method, all future forecasts are equal to a simple average of the observed data,

$$\hat{y}_{T+h|T} = \frac{1}{T} \sum_{t=1}^T y_t,$$

# Simple Exponential Smoothing

- The simplest of the exponentially smoothing methods is naturally called **simple exponential smoothing** (SES).
- This method is suitable for forecasting data with no clear trend or seasonal pattern.
- The naïve and the average as possible methods for forecasting such data.




Figure 8.1: Exports of goods and services from Algeria from 1960 to 2017.

We often want something between these two extremes. For example, it may be sensible to attach larger weights to more recent observations than to observations from the distant past. This is exactly the concept behind simple exponential smoothing. Forecasts are calculated using weighted averages, where the weights decrease exponentially as observations come from further in the past — the smallest weights are associated with the oldest observations:

$$\hat{y}_{T+1|T} = \alpha y_T + \alpha(1 - \alpha)y_{T-1} + \alpha(1 - \alpha)^2 y_{T-2} + \cdots, \quad (8.1)$$

# Simple Exponential Smoothing

## Forecast equation

$$\hat{y}_{T+1|T} = \alpha y_T + \alpha(1 - \alpha)y_{T-1} + \alpha(1 - \alpha)^2 y_{T-2} + \cdots,$$

where  $0 \leq \alpha \leq 1$ .

Weights assigned to observations for:				
Observation	$\alpha = 0.2$	$\alpha = 0.4$	$\alpha = 0.6$	$\alpha = 0.8$
$y_T$	0.2	0.4	0.6	0.8
$y_{T-1}$	0.16	0.24	0.24	0.16
$y_{T-2}$	0.128	0.144	0.096	0.032
$y_{T-3}$	0.1024	0.0864	0.0384	0.0064
$y_{T-4}$	$(0.2)(0.8)^4$	$(0.4)(0.6)^4$	$(0.6)(0.4)^4$	$(0.8)(0.2)^4$
$y_{T-5}$	$(0.2)(0.8)^5$	$(0.4)(0.6)^5$	$(0.6)(0.4)^5$	$(0.8)(0.2)^5$

For any  $\alpha$  between 0 and 1, the weights attached to the observations decrease exponentially as we go back in time, hence the name “exponential smoothing”. If  $\alpha$  is small (i.e., close to 0), more weight is given to observations from the more distant past. If  $\alpha$  is large (i.e., close to 1), more weight is given to the more recent observations. For the extreme case where  $\alpha = 1$ ,  $\hat{y}_{T+1|T} = y_T$ , and the forecasts are equal to the naïve forecasts.

**Source:** Forecasting Principles and Practice, 3<sup>rd</sup> Edition, Rob. J. Hyndman and George Athanasopoulos

# Simple Exponential Smoothing

## Component form

Forecast equation

$$\hat{y}_{t+h|t} = \ell_t$$

Smoothing equation

$$\ell_t = \alpha y_t + (1 - \alpha)\ell_{t-1}$$

- $\ell_t$  is the level (or the smoothed value) of the series at time  $t$ .
- $\hat{y}_{t+1|t} = \alpha y_t + (1 - \alpha)\hat{y}_{t|t-1}$
- $\hat{y}_{T+h|T} = \ell_T, h = 2, 3, \dots$

Iterate to get exponentially weighted moving average form.

## Weighted average form

$$\hat{y}_{T+1|T} = \sum_{j=0}^{T-1} \alpha(1 - \alpha)^j y_{T-j} + (1 - \alpha)^T \ell_0$$

# Simple Exponential Smoothing

- **Optimising smoothing parameters**


- Need to choose best values for  $\alpha$  and  $\ell_0$ .
- Similarly to regression, choose optimal parameters by minimising SSE:

$$\text{SSE} = \sum_{t=1}^T (y_t - \hat{y}_{t|t-1})^2.$$

- Unlike regression there is no closed form solution — use numerical optimization.
- For Algerian Exports example:
  - ▶  $\hat{\alpha} = 0.8400$
  - ▶  $\hat{\ell}_0 = 39.54$

**Source:** Forecasting Principles and Practice, 3<sup>rd</sup> Edition, Rob. J. Hyndman and George Athanasopoulos

# Simple Exponential Smoothing



Source: Forecasting Principles and Practice, 3<sup>rd</sup> Edition, Rob. J. Hyndman and George Athanasopoulos



# Holt's linear trend

## Component form

Forecast

$$\hat{y}_{t+h|t} = \ell_t + hb_t$$

Level

$$\ell_t = \alpha y_t + (1 - \alpha)(\ell_{t-1} + b_{t-1})$$


Trend

$$b_t = \beta^*(\ell_t - \ell_{t-1}) + (1 - \beta^*)b_{t-1},$$

- Two smoothing parameters  $\alpha$  and  $\beta^*$  ( $0 \leq \alpha, \beta^* \leq 1$ ).
- $\ell_t$  level: weighted average between  $y_t$  and one-step ahead forecast for time  $t$ , ( $\ell_{t-1} + b_{t-1} = \hat{y}_{t|t-1}$ )
- $b_t$  slope: weighted average of  $(\ell_t - \ell_{t-1})$  and  $b_{t-1}$ , current and previous estimate of slope.
- Choose  $\alpha, \beta^*, \ell_0, b_0$  to minimise SSE.


**Source:** Forecasting Principles and Practice, 3<sup>rd</sup> Edition, Rob. J. Hyndman and George Athanasopoulos

# Holt's linear trend



**Source:** Forecasting Principles and Practice, 3<sup>rd</sup> Edition, Rob. J. Hyndman and George Athanasopoulos

# Holt's linear trend



**Source:** Forecasting Principles and Practice, 3<sup>rd</sup> Edition, Rob. J. Hyndman and George Athanasopoulos

# Damped trend method

## Component form


$$\hat{y}_{t+h|t} = \ell_t + (\phi + \phi^2 + \dots + \phi^h)b_t$$

$$\ell_t = \alpha y_t + (1 - \alpha)(\ell_{t-1} + \phi b_{t-1})$$

$$b_t = \beta^*(\ell_t - \ell_{t-1}) + (1 - \beta^*)\phi b_{t-1}.$$

- Damping parameter  $0 < \phi < 1$ .
- If  $\phi = 1$ , identical to Holt's linear trend.
- As  $h \rightarrow \infty$ ,  $\hat{y}_{T+h|T} \rightarrow \ell_T + \phi b_T / (1 - \phi)$ .
- Short-run forecasts trended, long-run forecasts constant.

# Damped trend method



**Source:** Forecasting Principles and Practice, 3<sup>rd</sup> Edition, Rob. J. Hyndman and George Athanasopoulos

# Comparison: Australian Population

term	SES	Linear trend	Damped trend
$\alpha$	1.00	1.00	1.00
$\beta^*$		0.30	0.40
$\phi$			0.98
$\ell_0$	10.28	10.05	10.04
$b_0$		0.22	0.25
Training RMSE	0.24	0.06	0.07
Test RMSE	1.63	0.15	0.21
Test MASE	6.18	0.55	0.75
Test MAPE	6.09	0.55	0.74
Test MAE	1.45	0.13	0.18

**Source:** Forecasting Principles and Practice, 3<sup>rd</sup> Edition, Rob. J. Hyndman and George Athanasopoulos

# Holt-Winters additive method

Holt and Winters extended Holt's method to capture seasonality.

## Component form

$$\hat{y}_{t+h|t} = \ell_t + hb_t + s_{t+h-m(k+1)}$$

$$\ell_t = \alpha(y_t - s_{t-m}) + (1 - \alpha)(\ell_{t-1} + b_{t-1})$$

$$b_t = \beta^*(\ell_t - \ell_{t-1}) + (1 - \beta^*)b_{t-1}$$

$$s_t = \gamma(y_t - \ell_{t-1} - b_{t-1}) + (1 - \gamma)s_{t-m}$$

- $k = \text{integer part of } (h - 1)/m$ . Ensures estimates from the final year are used for forecasting.
- Parameters:  $0 \leq \alpha \leq 1$ ,  $0 \leq \beta^* \leq 1$ ,  $0 \leq \gamma \leq 1 - \alpha$  and  $m = \text{period of seasonality (e.g. } m = 4 \text{ for quarterly data)}$ .


# Holt-Winters additive method

- Seasonal component is usually expressed as
$$s_t = \gamma^*(y_t - \ell_t) + (1 - \gamma^*)s_{t-m}.$$
- Substitute in for  $\ell_t$ :
$$s_t = \gamma^*(1 - \alpha)(y_t - \ell_{t-1} - b_{t-1}) + [1 - \gamma^*(1 - \alpha)]s_{t-m}$$
- We set  $\gamma = \gamma^*(1 - \alpha)$ .
- The usual parameter restriction is  $0 \leq \gamma^* \leq 1$ , which translates to  $0 \leq \gamma \leq (1 - \alpha)$ .

**Source:** Forecasting Principles and Practice, 3<sup>rd</sup> Edition, Rob. J. Hyndman and George Athanasopoulos




# Exponential smoothing: seasonality



**Source:** Forecasting Principles and Practice, 3<sup>rd</sup> Edition, Rob. J. Hyndman and George Athanasopoulos

# Exponential smoothing: seasonality



**Source:** Forecasting Principles and Practice, 3<sup>rd</sup> Edition, Rob. J. Hyndman and George Athanasopoulos

# Holt-Winters multiplicative method

Seasonal variations change in proportion to the level of the series.

## Component form

$$\hat{y}_{t+h|t} = (\ell_t + hb_t)s_{t+h-m(k+1)}$$


$$\ell_t = \alpha \frac{y_t}{s_{t-m}} + (1 - \alpha)(\ell_{t-1} + b_{t-1})$$

$$b_t = \beta^*(\ell_t - \ell_{t-1}) + (1 - \beta^*)b_{t-1}$$

$$s_t = \gamma \frac{y_t}{(\ell_{t-1} + b_{t-1})} + (1 - \gamma)s_{t-m}$$

- $k$  is integer part of  $(h - 1)/m$ .
- Additive method:  $s_t$  in absolute terms — within each year  $\sum_i s_i \approx 0$ .
- Multiplicative method:  $s_t$  in relative terms — within each year  $\sum_i s_i \approx m$ .

# Example: Australian holiday tourism



**Source:** Forecasting Principles and Practice, 3<sup>rd</sup> Edition, Rob. J. Hyndman and George Athanasopoulos

# Holt-Winters damped method

Often the single most accurate forecasting method for seasonal data:

$$\hat{y}_{t+h|t} = [\ell_t + (\phi + \phi^2 + \dots + \phi^h)b_t]s_{t+h-m(k+1)}$$


$$\ell_t = \alpha(y_t/s_{t-m}) + (1 - \alpha)(\ell_{t-1} + \phi b_{t-1})$$

$$b_t = \beta^*(\ell_t - \ell_{t-1}) + (1 - \beta^*)\phi b_{t-1}$$

$$s_t = \gamma \frac{y_t}{(\ell_{t-1} + \phi b_{t-1})} + (1 - \gamma)s_{t-m}$$

**Source:** Forecasting Principles and Practice, 3<sup>rd</sup> Edition, Rob. J. Hyndman and George Athanasopoulos

# Holt-Winters with daily data



**Source:** Forecasting Principles and Practice, 3<sup>rd</sup> Edition, Rob. J. Hyndman and George Athanasopoulos

# Exponential smoothing methods

		Seasonal Component		
		N	A	M
Trend Component		(None)	(Additive)	(Multiplicative)
N	(None)	(N,N)	(N,A)	(N,M)
A	(Additive)	(A,N)	(A,A)	(A,M)
A <sub>d</sub>	(Additive damped)	(A <sub>d</sub> ,N)	(A <sub>d</sub> ,A)	(A <sub>d</sub> ,M)

**(N,N):** Simple exponential smoothing  
**(A,N):** Holt's linear method  
**(A<sub>d</sub>,N):** Additive damped trend method  
**(A,A):** Additive Holt-Winters' method  
**(A,M):** Multiplicative Holt-Winters' method  
**(A<sub>d</sub>,M):** Damped multiplicative Holt-Winters' method

**Source:** Forecasting Principles and Practice, 3<sup>rd</sup> Edition, Rob. J. Hyndman and George Athanasopoulos

# Exponential smoothing methods

Trend	Seasonal		
	N	A	M
N	$\hat{y}_{t+h t} = \ell_t$	$\hat{y}_{t+h t} = \ell_t + s_{t+h-m(k+1)}$	$\hat{y}_{t+h t} = \ell_t s_{t+h-m(k+1)}$
	$\ell_t = \alpha y_t + (1 - \alpha)\ell_{t-1}$	$\ell_t = \alpha(y_t - s_{t-m}) + (1 - \alpha)\ell_{t-1}$ $s_t = \gamma(y_t - \ell_{t-1}) + (1 - \gamma)s_{t-m}$	$\ell_t = \alpha(y_t/s_{t-m}) + (1 - \alpha)\ell_{t-1}$ $s_t = \gamma(y_t/\ell_{t-1}) + (1 - \gamma)s_{t-m}$
A	$\hat{y}_{t+h t} = \ell_t + hb_t$	$\hat{y}_{t+h t} = \ell_t + hb_t + s_{t+h-m(k+1)}$	$\hat{y}_{t+h t} = (\ell_t + hb_t)s_{t+h-m(k+1)}$
	$\ell_t = \alpha y_t + (1 - \alpha)(\ell_{t-1} + b_{t-1})$ $b_t = \beta^*(\ell_t - \ell_{t-1}) + (1 - \beta^*)b_{t-1}$	$\ell_t = \alpha(y_t - s_{t-m}) + (1 - \alpha)(\ell_{t-1} + b_{t-1})$ $b_t = \beta^*(\ell_t - \ell_{t-1}) + (1 - \beta^*)b_{t-1}$ $s_t = \gamma(y_t - \ell_{t-1} - b_{t-1}) + (1 - \gamma)s_{t-m}$	$\ell_t = \alpha(y_t/s_{t-m}) + (1 - \alpha)(\ell_{t-1} + b_{t-1})$ $b_t = \beta^*(\ell_t - \ell_{t-1}) + (1 - \beta^*)b_{t-1}$ $s_t = \gamma(y_t/(\ell_{t-1} + b_{t-1})) + (1 - \gamma)s_{t-m}$
Ad	$\hat{y}_{t+h t} = \ell_t + \phi_h b_t$	$\hat{y}_{t+h t} = \ell_t + \phi_h b_t + s_{t+h-m(k+1)}$	$\hat{y}_{t+h t} = (\ell_t + \phi_h b_t)s_{t+h-m(k+1)}$
	$\ell_t = \alpha y_t + (1 - \alpha)(\ell_{t-1} + \phi b_{t-1})$ $b_t = \beta^*(\ell_t - \ell_{t-1}) + (1 - \beta^*)\phi b_{t-1}$	$\ell_t = \alpha(y_t - s_{t-m}) + (1 - \alpha)(\ell_{t-1} + \phi b_{t-1})$ $b_t = \beta^*(\ell_t - \ell_{t-1}) + (1 - \beta^*)\phi b_{t-1}$ $s_t = \gamma(y_t - \ell_{t-1} - \phi b_{t-1}) + (1 - \gamma)s_{t-m}$	$\ell_t = \alpha(y_t/s_{t-m}) + (1 - \alpha)(\ell_{t-1} + \phi b_{t-1})$ $b_t = \beta^*(\ell_t - \ell_{t-1}) + (1 - \beta^*)\phi b_{t-1}$ $s_t = \gamma(y_t/(\ell_{t-1} + \phi b_{t-1})) + (1 - \gamma)s_{t-m}$

$\phi_h = \phi + \phi^2 + \dots + \phi^h$   
 $k$  is the integer part of  $(h - 1)/m$

**Source:** Forecasting Principles and Practice, 3<sup>rd</sup> Edition, Rob. J. Hyndman and George Athanasopoulos





For Your Valuable Time.